

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2003 (03.01.2003)

PCT

(10) International Publication Number
WO 03/000844 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number: **PCT/US02/19360**
- (22) International Filing Date: **18 June 2002 (18.06.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
- | | | |
|------------|-----------------------------|----|
| 60/300,508 | 22 June 2001 (22.06.2001) | US |
| 60/303,445 | 6 July 2001 (06.07.2001) | US |
| 60/305,405 | 13 July 2001 (13.07.2001) | US |
| 60/311,442 | 9 August 2001 (09.08.2001) | US |
| 60/314,821 | 24 August 2001 (24.08.2001) | US |
| 60/315,992 | 29 August 2001 (29.08.2001) | US |
| 60/378,205 | 3 May 2002 (03.05.2002) | US |

(71) Applicant (*for all designated States except US*): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GANDHI, Ameena, R.** [US/US]; 705 5th Avenue, San Francisco, CA 94118 (US). **KABLE, Amy, E.** [US/US]; 2345 Polk Street #4, San Francisco, CA 94109 (US). **SWARNAKAR, Anita** [CA/US]; 8 Locksley Avenue #5D, San Francisco, CA 94122 (US). **HAFALIA, April, J.A.** [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). **TRAN, Bao** [US/US]; 750 Salberg Avenue, Santa Clara, CA 95051 (US). **DUGGAN, Brendan, M.** [AU/US]; 243 Buena Vista Avenue # 306, Sunnyvale, CA 94086 (US). **WARREN, Bridget, A.** [US/US]; 2250 Homestead Court # 2, Los Altos, CA 94024 (US). **ISON, Craig, H.** [US/US]; 1242 Weathersfield Way, San Jose, CA 95118 (US). **HONCHELL, Cynthia, D.** [US/US]; 400 Laurel Street #203, San Carlos, CA 94070 (US). **NGUYEN, Danniell, B.** [US/US]; 1403 Ridgewood Drive, San Jose, CA 95118 (US). **LU, Dyung Aina, M.** [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **LEE, Ernestine, A.** [US/US]; 20523 Crow Creek Road, Castro Valley, CA 94552 (US). **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **FORSYTHE, Ian, J.** [US/US]; 308 Roble Avenue, Redwood City, CA 94061 (US). **BARROSO, Ines** [PT/GB]; 38 Eden Street, Cambridge, Cambridgeshire CB1 1EL (GB). **RAMKUMAR, Jayalaxmi** [IN/US];

34359 Maybird Circle, Fremont, CA 94555 (US). **GRIF-FIN, Jennifer, A.** [US/US]; 33691 Mello Way, Fremont, CA 94555 (US). **LI, Joana, X.** [US/US]; 1264 Geneva Avenue, San Francisco, CA 94112 (US). **YANG, Junming** [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US). **THANGAVELU, Kavitha** [IN/US]; 1950 Montecito Avenue #23, Mountain View, CA 94043 (US). **GIETZEN, Kimberly, J.** [US/US]; 691 Los Huecos Drive, San Jose, CA 95123 (US). **DING, Li** [CN/US]; 3353 Alma Street # 146, Palo Alto, CA 94306 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **BOROWSKY, Mark, L.** [US/US]; 122 Orchard Avenue, Redwood City, CA 94061 (US). **YAO, Monique, G.** [US/US]; 1189 Woodgate Drive, Carmel, IN 46033 (US). **WALIA, Narinder, K.** [US/US]; 890 Davis Street A # 205, San Leandro, CA 94577 (US). **MASON, Patricia, M.** [US/US]; 360 Clarke Lane, Morgan Hill, CA 95014 (US). **GURURAJAN, Rajagopal** [IN/US]; 5591 Dent Avenue, San Jose, CA 95118 (US). **LEE, Sally** [US/US]; 3643 26th Street, San Francisco, CA 94110 (US). **BECHA, Shanya, D.** [US/US]; 21062 Gary Drive # 117, Castro Valley, CA 94546 (US). **LEE, Soo Yeun** [KR/US]; 40 Westdale Avenue, Daly City, CA 94015 (US). **TRAN, Uyen, K.** [US/US]; 2638 Mabury Square, San Jose, CA 95133 (US). **ELLIOTT, Vicki, S.** [US/US]; 3770 Polton Place Way, San Jose, CA 95121 (US). **LUO, Wen** [CN/US]; 181 Cymbidium Circle, South San Francisco, CA 94080 (US). **SPRAGUE, William, W.** [US/US]; 611 13th Street # C, Sacramento, CA 95814 (US). **TANG, Y. Tom** [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **LU, Yan** [CN/US]; 3885 Corrina Way, Palo Alto, CA 94303 (US). **ZEBAJADIAN, Yeganeh** [IR/US]; 830 Junipero Serra Boulevard, San Francisco, CA 94127 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AI., AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),

[Continued on next page]

(54) Title: **PROTEIN MODIFICATION AND MAINTENANCE MOLECULES**

(57) Abstract: Various embodiments of the invention provide human proteinmodification and maintenance molecules (PMOD) and polynucleotides which identify and encode PMOD. Embodiments of the invention also provide expression vectors, host cells, antibodies, agonists, and antagonists. Other embodiments provide methods for diagnosing, treating, or preventing disorders associated with aberrant expression of PMOD.



WO 03/000844 A2



Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *without international search report and to be republished
upon receipt of that report*

PROTEIN MODIFICATION AND MAINTENANCE MOLECULES

TECHNICAL FIELD

The invention relates to novel nucleic acids, protein modification and maintenance molecules encoded by these nucleic acids, and to the use of these nucleic acids and proteins in the diagnosis, treatment, and prevention of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders. The invention also relates to the assessment of the effects of exogenous compounds on the expression of nucleic acids and protein modification and maintenance molecules.

BACKGROUND OF THE INVENTION

The cellular processes regulating modification and maintenance of protein molecules coordinate their function, conformation, stabilization, and degradation. Each of these processes is mediated by key enzymes or proteins such as kinases, phosphatases, proteases, protease inhibitors, isomerases, transferases, and molecular chaperones.

Kinases

Kinases catalyze the transfer of high-energy phosphate groups from adenosine triphosphate (ATP) to target proteins on the hydroxyamino acid residues serine, threonine, or tyrosine. Addition of a phosphate group alters the local charge on the acceptor molecule, causing internal conformational changes and potentially influencing intermolecular contacts. Reversible protein phosphorylation is the ubiquitous strategy used to control many of the intracellular events in eukaryotic cells. It is estimated that more than ten percent of proteins active in a typical mammalian cell are phosphorylated. Extracellular signals including hormones, neurotransmitters, and growth and differentiation factor can activate kinases, which can occur as cell surface receptors or as the activator of the final effector protein, but can also occur along the signal transduction pathway. Kinases are involved in all aspects of a cell's function, from basic metabolic processes, such as glycolysis, to cell-cycle regulation, differentiation, and communication with the extracellular environment through signal transduction cascades. Inappropriate phosphorylation of proteins in cells has been linked to changes in cell cycle progression and cell differentiation. Changes in the cell cycle have been linked to induction of apoptosis or cancer. Changes in cell differentiation have been linked to diseases and disorders of the reproductive system, immune system, and skeletal muscle.

There are two classes of protein kinases. One class, protein tyrosine kinases (PTKs), phosphorylates tyrosine residues, and the other class, protein serine/threonine kinases (STKs), phosphorylates serine and threonine residues. Some PTKs and STKs possess structural characteristics of both families and have dual specificity for both tyrosine and serine/threonine

residues. Almost all kinases contain a conserved 250-300 amino acid catalytic domain containing specific residues and sequence motifs characteristic of the kinase family. (Reviewed in Hardie, G. and Hanks, S. (1995) The Protein Kinase Facts Book, Vol I p.p. 17-20 Academic Press, San Diego, CA.).

5 Phosphatases

Phosphatases hydrolytically remove phosphate groups from proteins. Phosphatases are essential in determining the extent of phosphorylation in the cell and, together with kinases, regulate key cellular processes such as metabolic enzyme activity, proliferation, cell growth and differentiation, cell adhesion, and cell cycle progression. Protein phosphatases are characterized as either serine/threonine- or tyrosine-specific based on their preferred phospho-amino acid substrate. Some phosphatases (DSPs, for dual specificity phosphatases) can act on phosphorylated tyrosine, serine, or threonine residues. The protein serine/threonine phosphatases (PSPs) are important regulators of many cAMP-mediated hormone responses in cells. Protein tyrosine phosphatases (PTPs) play a significant role in cell cycle and cell signaling processes.

15 Proteases

Proteases cleave proteins and peptides at the peptide bond that forms the backbone of the protein or peptide chain. Proteolysis is one of the most important and frequent enzymatic reactions that occurs both within and outside of cells. Proteolysis is responsible for the activation and maturation of nascent polypeptides, the degradation of misfolded and damaged proteins, and the controlled turnover of peptides within the cell. Proteases participate in digestion, endocrine function, and tissue remodeling during embryonic development, wound healing, and normal growth. Proteases can play a role in regulatory processes by affecting the half life of regulatory proteins. Proteases are involved in the etiology or progression of disease states such as inflammation, angiogenesis, tumor dispersion and metastasis, cardiovascular disease, neurological disease, and bacterial, parasitic, and viral infections.

Proteases can be categorized on the basis of where they cleave their substrates. Exopeptidases, which include aminopeptidases, dipeptidyl peptidases, tripeptidases, carboxypeptidases, peptidyl-di-peptidases, dipeptidases, and omega peptidases, cleave residues at the termini of their substrates. Endopeptidases, including serine proteases, cysteine proteases, and metalloproteases, cleave at residues within the peptide. Four principal categories of mammalian proteases have been identified based on active site structure, mechanism of action, and overall three-dimensional structure. (See Beynon, R.J. and J.S. Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University Press, New York, NY, pp. 1-5.)

Serine Proteases

35 The serine proteases (SPs) are a large, widespread family of proteolytic enzymes that include

the digestive enzymes trypsin and chymotrypsin, components of the complement and blood-clotting cascades, and enzymes that control the degradation and turnover of macromolecules within the cell and in the extracellular matrix. Most of the more than 20 subfamilies can be grouped into six clans, each with a common ancestor. These six clans are hypothesized to have descended from at least four evolutionarily distinct ancestors. SPs are named for the presence of a serine residue found in the active catalytic site of most families. The active site is defined by the catalytic triad, a set of conserved asparagine, histidine, and serine residues critical for catalysis. These residues form a charge relay network that facilitates substrate binding. Other residues outside the active site form an oxyanion hole that stabilizes the tetrahedral transition intermediate formed during catalysis. SPs have a wide range of substrates and can be subdivided into subfamilies on the basis of their substrate specificity. The main subfamilies are named for the residue(s) after which they cleave: trypases (after arginine or lysine), aspases (after aspartate), chymases (after phenylalanine or leucine), metases (methionine), and serases (after serine) (Rawlings, N.D. and A.J. Barrett (1994) *Meth. Enz.* 244:19-61).

Most mammalian serine proteases are synthesized as zymogens, inactive precursors that are activated by proteolysis. For example, trypsinogen is converted to its active form, trypsin, by enteropeptidase. Enteropeptidase is an intestinal protease that removes an N-terminal fragment from trypsinogen. The remaining active fragment is trypsin, which in turn activates the precursors of the other pancreatic enzymes. Likewise, proteolysis of prothrombin, the precursor of thrombin, generates three separate polypeptide fragments. The N-terminal fragment is released while the other two fragments, which comprise active thrombin, remain associated through disulfide bonds.

The two largest SP subfamilies are the chymotrypsin (S1) and subtilisin (S8) families. Some members of the chymotrypsin family contain two structural domains unique to this family. Kringle domains are triple-looped, disulfide cross-linked domains found in varying copy number. Kringles are thought to play a role in binding mediators such as membranes, other proteins or phospholipids, and in the regulation of proteolytic activity (PROSITE PDOC00020). Apple domains are 90 amino-acid repeated domains, each containing six conserved cysteines. Three disulfide bonds link the first and sixth, second and fifth, and third and fourth cysteines (PROSITE PDOC00376). Apple domains are involved in protein-protein interactions. S1 family members include trypsin, chymotrypsin, coagulation factors IX-XII, complement factors B, C, and D, granzymes, kallikrein, and tissue- and urokinase-plasminogen activators. The subtilisin family has members found in the eubacteria, archaeobacteria, eukaryotes, and viruses. Subtilisins include the proprotein-processing endopeptidases kexin and furin and the pituitary prohormone convertases PC1, PC2, PC3, PC6, and PACE4 (Rawlings and Barrett, *supra*).

SPs have functions in many normal processes and some have been implicated in the etiology

or treatment of disease. Enterokinase, the initiator of intestinal digestion, is found in the intestinal brush border, where it cleaves the acidic propeptide from trypsinogen to yield active trypsin (Kitamoto, Y. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91: 7588-7592). Prolylcarboxypeptidase, a lysosomal serine peptidase that cleaves peptides such as angiotensin II and III and [des-Arg⁹] bradykinin, shares sequence homology with members of both the serine carboxypeptidase and prolylendopeptidase families (Tan, F. et al. (1993) *J. Biol. Chem.* 268:16631-16638). The protease neuropsin may influence synapse formation and neuronal connectivity in the hippocampus in response to neural signaling (Chen, Z.-L. et al. (1995) *J Neurosci* 15:5088-5097). Tissue plasminogen activator is useful for acute management of stroke (Zivin, J.A. (1999) *Neurology* 53:14-9) and myocardial infarction (Ross, A.M. (1999) *Clin Cardiol* 22:165-71). Some receptors (PAR, for proteinase-activated receptor), highly expressed throughout the digestive tract, are activated by proteolytic cleavage of an extracellular domain. The major agonists for PARs, thrombin, trypsin, and mast cell tryptase, are released in allergy and inflammatory conditions. Control of PAR activation by proteases has been suggested as a promising therapeutic target (Vergnolle, N. (2000) *Aliment. Pharmacol. Ther.* 14:257-266; Rice, K.D. et al. (1998) *Curr. Pharm. Des.* 4:381-396). Prostate-specific antigen (PSA) is a kallikrein-like serine protease synthesized and secreted exclusively by epithelial cells in the prostate gland. Serum PSA is elevated in prostate cancer and is the most sensitive physiological marker for monitoring cancer progression and response to therapy. PSA can also identify the prostate as the origin of a metastatic tumor. (Brawer, M. K. and Lange, P. H. (1989) *Urology* 33:11-16).

The kallikreins are a subfamily of serine proteases. KLK14 is a kallikrein gene located within the human kallikrein locus at 19q13.4. KLK14 is approximately 5.4 kb in length and transcribes two alternative transcripts present only in prostate and skeletal muscle. In prostate, KLK14 is expressed by both benign and malignant glandular epithelial cells, thus exhibiting an expression pattern similar to that of two other prostatic kallikreins, KLK2 and KLK3, which encode K2 and prostate-specific antigen, respectively (Hooper, J.D. et al. (2001) *Genomics* 73:117-122).

The signal peptidase is a specialized class of SP found in all prokaryotic and eukaryotic cell types that serves in the processing of signal peptides from certain proteins. Signal peptides are amino-terminal domains of a protein which direct the protein from its ribosomal assembly site to a particular cellular or extracellular location. Once the protein has been exported, removal of the signal sequence by a signal peptidase and posttranslational processing, e.g., glycosylation or phosphorylation, activate the protein. Signal peptidases exist as multi-subunit complexes in both yeast and mammals. The canine signal peptidase complex is composed of five subunits, all associated with the microsomal membrane and containing hydrophobic regions that span the membrane one or more times (Shelness, G.S. and G. Blobel (1990) *J. Biol. Chem.* 265:9512-9519).

Some of these subunits serve to fix the complex in its proper position on the membrane while others contain the actual catalytic activity.

Another family of proteases which have a serine in their active site are dependent on the hydrolysis of ATP for their activity. These proteases contain proteolytic core domains and regulatory ATPase domains which can be identified by the presence of the P-loop, an ATP/GTP-binding motif (PROSITE PDOC00803). Members of this family include the eukaryotic mitochondrial matrix proteases, Clp protease and the proteasome. Clp protease was originally found in plant chloroplasts but is believed to be widespread in both prokaryotic and eukaryotic cells. The gene for early-onset torsion dystonia encodes a protein related to Clp protease (Ozelius, L.J. et al. (1998) Adv. Neurol. 78:93-105).

The proteasome is an intracellular protease complex found in some bacteria and in all eukaryotic cells, and plays an important role in cellular physiology. Proteasomes are associated with the ubiquitin conjugation system (UCS), a major pathway for the degradation of cellular proteins of all types, including proteins that function to activate or repress cellular processes such as transcription and cell cycle progression (Ciechanover, A. (1994) Cell 79:13-21). In the UCS pathway, proteins targeted for degradation are conjugated to ubiquitin, a small heat stable protein. The ubiquitinated protein is then recognized and degraded by the proteasome. The resultant ubiquitin-peptide complex is hydrolyzed by a ubiquitin carboxyl terminal hydrolase, and free ubiquitin is released for reutilization by the UCS. Ubiquitin-proteasome systems are implicated in the degradation of mitotic cyclic kinases, oncoproteins, tumor suppressor genes (p53), cell surface receptors associated with signal transduction, transcriptional regulators, and mutated or damaged proteins (Ciechanover, *supra*). This pathway has been implicated in a number of diseases, including cystic fibrosis, Angelman's syndrome, and Liddle syndrome (reviewed in Schwartz, A.L. and A. Ciechanover (1999) Ann. Rev. Med. 50:57-74). A murine proto-oncogene, Unp, encodes a nuclear ubiquitin protease whose overexpression leads to oncogenic transformation of NIH3T3 cells. The human homologue of this gene is consistently elevated in small cell tumors and adenocarcinomas of the lung (Gray, D.A. (1995) Oncogene 10:2179-2183). Ubiquitin carboxyl terminal hydrolase is involved in the differentiation of a lymphoblastic leukemia cell line to a non-dividing mature state (Maki, A. et al. (1996) Differentiation 60:59-66). In neurons, ubiquitin carboxyl terminal hydrolase (PGP 9.5) expression is strong in the abnormal structures that occur in human neurodegenerative diseases (Lowe, J. et al. (1990) J. Pathol. 161:153-160). The proteasome is a large (~2000 kDa) multisubunit complex composed of a central catalytic core containing a variety of proteases arranged in four seven-membered rings with the active sites facing inwards into the central cavity, and terminal ATPase subunits covering the outer port of the cavity and regulating substrate entry (for review, see Schmidt, M. et al. (1999) Curr. Op. Chem. Biol. 3:584-591).

Cysteine Proteases

Cysteine proteases (CPs) are involved in diverse cellular processes ranging from the processing of precursor proteins to intracellular degradation. Nearly half of the CPs known are present only in viruses. CPs have a cysteine as the major catalytic residue at the active site where catalysis proceeds via a thioester intermediate and is facilitated by nearby histidine and asparagine residues. A glutamine residue is also important, as it helps to form an oxyanion hole. Two important CP families include the papain-like enzymes (C1) and the calpains (C2). Papain-like family members are generally lysosomal or secreted and therefore are synthesized with signal peptides as well as propeptides. Most members bear a conserved motif in the propeptide that may have structural significance (Karrer, K.M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:3063-3067). Three-dimensional structures of papain family members show a bilobed molecule with the catalytic site located between the two lobes. Papains include cathepsins B, C, H, L, and S, certain plant allergens and dipeptidyl peptidase (for a review, see Rawlings, N.D. and A.J. Barrett (1994) *Meth. Enz.* 244:461-486).

Some CPs are expressed ubiquitously, while others are produced only by cells of the immune system. Of particular note, CPs are produced by monocytes, macrophages and other cells which migrate to sites of inflammation and secrete molecules involved in tissue repair. Overabundance of these repair molecules plays a role in certain disorders. In autoimmune diseases such as rheumatoid arthritis, secretion of the cysteine peptidase cathepsin C degrades collagen, laminin, elastin and other structural proteins found in the extracellular matrix of bones. Bone weakened by such degradation is also more susceptible to tumor invasion and metastasis. Cathepsin L expression may also contribute to the influx of mononuclear cells which exacerbates the destruction of the rheumatoid synovium (Keyszer, G.M. (1995) *Arthritis Rheum.* 38:976-984).

Calpains are calcium-dependent cytosolic endopeptidases which contain both an N-terminal catalytic domain and a C-terminal calcium-binding domain. Calpain is expressed as a proenzyme heterodimer consisting of a catalytic subunit unique to each isoform and a regulatory subunit common to different isoforms. Each subunit bears a calcium-binding EF-hand domain. The regulatory subunit also contains a hydrophobic glycine-rich domain that allows the enzyme to associate with cell membranes. Calpains are activated by increased intracellular calcium concentration, which induces a change in conformation and limited autolysis. The resultant active molecule requires a lower calcium concentration for its activity (Chan S.L. and Mattson M.P. (1999) *J. Neurosci. Res.* 58:167-190). Calpain expression is predominantly neuronal, although it is present in other tissues. Several chronic neurodegenerative disorders, including ALS, Parkinson's disease and Alzheimer's disease are associated with increased calpain expression (Chan and Mattson, *supra*). Calpain-mediated breakdown of the cytoskeleton has been proposed to contribute to brain damage resulting from head

injury (McCracken E. et al. (1999) *J. Neurotrauma* 16:749-61). Calpain-3 is predominantly expressed in skeletal muscle, and is responsible for limb-girdle muscular dystrophy type 2A (Minami, N. et al. (1999) *J. Neurol. Sci.* 171:31-37).

Another family of thiol proteases is the caspases, which are involved in the initiation and execution phases of apoptosis. A pro-apoptotic signal can activate initiator caspases that trigger a proteolytic caspase cascade, leading to the hydrolysis of target proteins and the classic apoptotic death of the cell. Two active site residues, a cysteine and a histidine, have been implicated in the catalytic mechanism. Caspases are among the most specific endopeptidases, cleaving after aspartate residues. Caspases are synthesized as inactive zymogens consisting of one large (p20) and one small (p10) subunit separated by a small spacer region, and a variable N-terminal prodomain. This prodomain interacts with cofactors that can positively or negatively affect apoptosis. An activating signal causes autoproteolytic cleavage of a specific aspartate residue (D297 in the caspase-1 numbering convention) and removal of the spacer and prodomain, leaving a p10/p20 heterodimer. Two of these heterodimers interact via their small subunits to form the catalytically active tetramer. The long prodomains of some caspase family members have been shown to promote dimerization and auto-processing of procaspases. Some caspases contain a "death effector domain" in their prodomain by which they can be recruited into self-activating complexes with other caspases and FADD protein associated death receptors or the TNF receptor complex. In addition, two dimers from different caspase family members can associate, changing the substrate specificity of the resultant tetramer. Endogenous caspase inhibitors (inhibitor of apoptosis proteins, or IAPs) also exist. All these interactions have clear effects on the control of apoptosis (reviewed in Chan and Mattson, *supra*; Salveson, G.S. and V.M. Dixit (1999) *Proc. Nat. Acad. Sci. USA* 96:10964-10967).

Caspases have been implicated in a number of diseases. Mice lacking some caspases have severe nervous system defects due to failed apoptosis in the neuroepithelium and suffer early lethality. Others show severe defects in the inflammatory response, as caspases are responsible for processing IL-1b and possibly other inflammatory cytokines (Chan and Mattson, *supra*). Cowpox virus and baculoviruses target caspases to avoid the death of their host cell and promote successful infection. In addition, increases in inappropriate apoptosis have been reported in AIDS, neurodegenerative diseases and ischemic injury, while a decrease in cell death is associated with cancer (Salveson and Dixit, *supra*; Thompson, C.B. (1995) *Science* 267:1456-1462).

Aspartyl proteases

Aspartyl proteases (APs) include the lysosomal proteases cathepsins D and E, as well as chymosin, renin, and the gastric pepsins. Most retroviruses encode an AP, usually as part of the pol polyprotein. APs, also called acid proteases, are monomeric enzymes consisting of two domains, each domain containing one half of the active site with its own catalytic aspartic acid residue. APs

are most active in the range of pH 2–3, at which one of the aspartate residues is ionized and the other neutral. The pepsin family of APs contains many secreted enzymes, and all are likely to be synthesized with signal peptides and propeptides. Most family members have three disulfide loops, the first ~5 residue loop following the first aspartate, the second 5-6 residue loop preceding the second aspartate, and the third and largest loop occurring toward the C terminus. Retropepsins, on the other hand, are analogous to a single domain of pepsin, and become active as homodimers with each retropepsin monomer contributing one half of the active site. Retropepsins are required for processing the viral polyproteins.

APs have roles in various tissues, and some have been associated with disease. Renin mediates the first step in processing the hormone angiotensin, which is responsible for regulating electrolyte balance and blood pressure (reviewed in Crews, D.E. and S.R. Williams (1999) *Hum. Biol.* 71:475-503). Abnormal regulation and expression of cathepsins are evident in various inflammatory disease states. Expression of cathepsin D is elevated in synovial tissues from patients with rheumatoid arthritis and osteoarthritis. The increased expression and differential regulation of the cathepsins are linked to the metastatic potential of a variety of cancers (Chambers, A.F. et al. (1993) *Crit. Rev. Oncol.* 4:95-114).

Metalloproteases

Metalloproteases require a metal ion for activity, usually manganese or zinc. Most zinc-dependent metalloproteases share a common sequence in the zinc-binding domain. The active site is made up of two histidines which act as zinc ligands and a catalytic glutamic acid C-terminal to the first histidine. Proteins containing this signature sequence are known as the metzincins and include aminopeptidase N, angiotensin-converting enzyme, neurolysin, the matrix metalloproteases and the adamalysins (ADAMS). An alternate sequence is found in the zinc carboxypeptidases, in which all three conserved residues – two histidines and a glutamic acid – are involved in zinc binding.

A number of the neutral metalloendopeptidases, including angiotensin converting enzyme and the aminopeptidases, are involved in the metabolism of peptide hormones. High aminopeptidase B activity, for example, is found in the adrenal glands and neurohypophyses of hypertensive rats (Prieto, I. Et al. (1998) *Horm. Metab. Res.* 30:246-248). Oligopeptidase M/neurolysin can hydrolyze bradykinin as well as neurotensin (Serizawa, A. et al. (1995) *J. Biol. Chem* 270:2092-2098). Neurotensin is a vasoactive peptide that can act as a neurotransmitter in the brain, where it has been implicated in limiting food intake (Tritos, N.A. et al. (1999) *Neuropeptides* 33:339-349).

The matrix metalloproteases (MMPs) are a family of at least 23 enzymes that can degrade components of the extracellular matrix (ECM). They are Zn^{+2} endopeptidases with an N-terminal catalytic domain. Nearly all members of the family have a hinge peptide and C-terminal domain which can bind to substrate molecules in the ECM or to inhibitors produced by the tissue (TIMPs, for

tissue inhibitor of metalloprotease; Campbell, I.L. et al. (1999) Trends Neurosci. 22:285). The presence of fibronectin-like repeats, transmembrane domains, or C-terminal hemopexinase-like domains can be used to separate MMPs into collagenase, gelatinase, stromelysin and membrane-type MMP subfamilies. In the inactive form, the Zn^{+2} ion in the active site interacts with a cysteine in the pro-sequence. Activating factors disrupt the Zn^{+2} -cysteine interaction, or "cysteine switch," exposing the active site. This partially activates the enzyme, which then cleaves off its propeptide and becomes fully active. MMPs are often activated by the serine proteases plasmin and furin. MMPs are often regulated by stoichiometric, noncovalent interactions with inhibitors; the balance of protease to inhibitor, then, is very important in tissue homeostasis (reviewed in Yong, V.W. et al. (1998) Trends Neurosci. 21:75).

MMPs are implicated in a number of diseases including osteoarthritis (Mitchell, P. et al. (1996) J. Clin. Inv. 97:761), atherosclerotic plaque rupture (Sukhova, G.K. et al. (1999) Circulation 99:2503), aortic aneurysm (Schneiderman, J. et al. (1998) Am. J. Path. 152:703), non-healing wounds (Saarialho-Kere, U.K. et al. (1994) J. Clin. Inv. 94:79), bone resorption (Blavier, L. and J.M. Delaisse (1995) J. Cell Sci. 108:3649), age-related macular degeneration (Steen, B. et al. (1998) Invest. Ophthalmol. Vis. Sci. 39:2194), emphysema (Finlay, G.A. et al. (1997) Thorax 52:502), myocardial infarction (Rohde, L.E. et al. (1999) Circulation 99:3063) and dilated cardiomyopathy (Thomas, C.V. et al. (1998) Circulation 97:1708). MMP inhibitors prevent metastasis of mammary carcinoma and experimental tumors in rat, and Lewis lung carcinoma, hemangioma, and human ovarian carcinoma xenografts in mice (Eccles S.A. et al. (1996) Cancer Res. 56:2815; Anderson et al. (1996) Cancer Res. 56:715-718; Volpert, O.V. et al. (1996) J. Clin. Invest. 98:671; Taraboletti, G. et al. (1995) JNCI 87:293; Davies, B. et al. (1993) Cancer Res. 53:2087). MMPs may be active in Alzheimer's disease. A number of MMPs are implicated in multiple sclerosis, and administration of MMP inhibitors can relieve some of its symptoms (reviewed in Yong, supra).

Another family of metalloproteases is the ADAMs, for A Disintegrin and Metalloprotease Domain, which they share with their close relatives the adamalysins, snake venom metalloproteases (SVMPs). ADAMs combine features of both cell surface adhesion molecules and proteases, containing a prodomain, a protease domain, a disintegrin domain, a cysteine rich domain, an epidermal growth factor repeat, a transmembrane domain, and a cytoplasmic tail. The first three domains listed above are also found in the SVMPs. The ADAMs possess four potential functions: proteolysis, adhesion, signaling and fusion. The ADAMs share the metzincin zinc binding sequence and are inhibited by some MMP antagonists such as TIMP-1.

ADAMs are implicated in such processes as sperm-egg binding and fusion, myoblast fusion, and protein-ectodomain processing or shedding of cytokines, cytokine receptors, adhesion proteins and other extracellular protein domains (Schlöndorff, J. and C.P. Blobel (1999) J. Cell. Sci.

112:3603-3617). The Kuzbanian protein cleaves a substrate in the NOTCH pathway (possibly NOTCH itself), activating the program for lateral inhibition in *Drosophila* neural development. Two ADAMs, TACE (ADAM 17) and ADAM 10, are proposed to have analogous roles in the processing of amyloid precursor protein in the brain (Schlöndorff and Blobel, supra). TACE has also been identified as the TNF activating enzyme (Black, R.A. et al. (1997) Nature 385:729). TNF is a pleiotropic cytokine that is important in mobilizing host defenses in response to infection or trauma, but can cause severe damage in excess and is often overproduced in autoimmune disease. TACE cleaves membrane-bound pro-TNF to release a soluble form. Other ADAMs may be involved in a similar type of processing of other membrane-bound molecules.

- 10 The ADAMTS sub-family has all of the features of ADAM family metalloproteases and contain an additional thrombospondin domain (TS). The prototypic ADAMTS was identified in mouse, found to be expressed in heart and kidney and upregulated by proinflammatory stimuli (Kuno, K. et al. (1997) J. Biol. Chem. 272:556). To date eleven members are recognized by the Human Genome Organization (HUGO; <http://www.gene.ucl.ac.uk/users/hester/adamts.html#Approved>).
- 15 Members of this family have the ability to degrade aggrecan, a high molecular weight proteoglycan which provides cartilage with important mechanical properties including compressibility, and which is lost during the development of arthritis. Enzymes which degrade aggrecan are thus considered attractive targets to prevent and slow the degradation of articular cartilage (See, e.g., Tortorella, M.D. (1999) Science 284:1664; Abbaszade, I. (1999) J. Biol. Chem. 274:23443). Other members are reported to have antiangiogenic potential (Kuno et al., supra) and/or procollagen processing (Colige, A. et al. (1997) Proc.Natl. Acad. Sci. USA 94:2374).

- 20 All members of the MDC family of integral membrane proteins contain a metalloproteinase-like domain, a disintegrin-like domain and a cysteine-rich domain. They have been identified in a wide range of mammalian tissues and many are abundantly expressed in the male reproductive tract. A number of MDC proteins (fertilin alpha, fertilin beta, tMDC I, tMDC II and tMDC III) are localized to spermatogenic cells and processed as spermatozoa pass through the epididymis, yielding proteins that retain their disintegrin domain on mature spermatozoa. Fertilin beta and tMDC I have been implicated in egg recognition, mediated by a disintegrin-integrin interaction (Frayne, J. et al. (1998) J. Reprod. Fertil. Suppl. 53:149-155).

- 30 Examples of manganese metalloenzymes include aminopeptidase P and human proline dipeptidase (PEPD). Aminopeptidase P can degrade bradykinin, a nonapeptide activated in a variety of inflammatory responses. Aminopeptidase P has been implicated in coronary ischemia/reperfusion injury. Administration of aminopeptidase P inhibitors has been shown to have a cardioprotective effect in rats (Ersahin, C. et al (1999) J. Cardiovasc. Pharmacol. 34:604-611).

- 35 Protease inhibitors

Protease inhibitors and other regulators of protease activity control the activity and effects of proteases. Protease inhibitors have been shown to control pathogenesis in animal models of proteolytic disorders (Murphy, G. (1991) *Agents Actions Suppl.* 35:69-76). In patients with HIV disease protease inhibitors have been shown to be effective in preventing disease progression and

5 reducing mortality (Barry, M. et al. (1997) *Clin. Pharmacokinet.* 32:194-209). Low levels of the cystatins, low molecular weight inhibitors of the cysteine proteases, correlate with malignant progression of tumors. (Calkins, C. et al. (1995) *Biol. Biochem. Hoppe Seyler* 376:71-80). The cystatin superfamily of protease inhibitors is characterized by a particular pattern of linearly arranged and tandemly repeated disulfide loops (Kellermann, J. et al. (1989) *J. Biol. Chem.* 264:14121-14128).

10 An example of a representative of a structural prototype of a novel family among the cystatin superfamily is human alpha 2-HS glycoprotein (AHSG), a plasma protein synthesized in liver and selectively concentrated in bone matrix, dentine, and other mineralized tissues (Triffitt, J.T. (1976) *Calcif. Tissue Res.* 22:27-33), which is also classified as belonging to the fetuin family. Fetuins are characterized by the presence of 2 N-terminally located cystatin-like repeats and a unique C-terminal

15 domain which is not present in other proteins of the cystatin superfamily (PROSITE PDOC00966). AHSG has been reported to be involved in bone formation and resorption as well as immune responses (Yang, F. et al. (1992) 1130:149-156; Lee, C.C. et al. (1987) *PNAS USA* 84:4403-4407; Nakamura, O. et al. (1999) *Biosci. Biotechnol. Biochem.* 63:1383-1391). Additionally, AHSG has been implicated in infertility associated with endometriosis (Mathur, S.P. (2000) *Am. J. Reprod.*

20 *Immunol.* 44:89-95; Mathur, S.P. et al. (1999) *Autoimmunity* 29:121-127) and inhibition of osteogenesis (Binkert, C. et al. (1999) *J. Biol. Chem.* 274:28514-28520). Decreased serum levels of AHSG have been detected in patients with acute leukemias, chronic granulocyte and myelomonocyte leukemias, lymphomas, myelofibrosis, multiple myeloma, metastatizing solid tumors, systemic lupus erythematosus, rheumatoid arthritis, acute alcoholic hepatitis, fatty liver, chronic active hepatitis,

25 liver cirrhosis, acute and chronic pancreatitis, and Crohn's disease (Kalabay, L. et al. (1992) *Orv. Hetil.* 133:1553-1554; 1559-1560). Serpins are inhibitors of mammalian plasma serine proteases. Many serpins serve to regulate the blood clotting cascade and/or the complement cascade in mammals. Sp32 is a positive regulator of the mammalian acrosomal protease, acrosin, that binds the proenzyme, proacrosin, and thereby aides in packaging the enzyme into the acrosomal matrix (T.

30 Baba et al. (1994) *J. Biol. Chem.* 269:10133-10140). The Kunitz family of serine protease inhibitors are characterized by one or more "Kunitz domains" containing a series of cysteine residues that are regularly spaced over approximately 50 amino acid residues and form three intrachain disulfide bonds. Members of this family include aprotinin, tissue factor pathway inhibitor (TFPI-1 and TFPI-2), inter- α -trypsin inhibitor (ITI), and bikunin. (Marlor, C.W. et al. (1997) *J. Biol. Chem.* 272:12202-

35 12208.) Members of this family are potent inhibitors (in the nanomolar range) against serine

proteases such as kallikrein and plasmin. has clinical utility in reduction of perioperative blood loss. ITI has been found to inactivate human trypsin, chymotrypsin, neutrophil elastase and cathepsin G (Morii, M. et al. (1985) Biol. Chem. Hoppe Seyler 366:19-21); and is suspected of playing a key role in the biology of the extracellular matrix and in the pathophysiology of chronic bronchopulmonary diseases or lung cancer progression (Cuvelier, A. et al. (2000) Rev. Mal. Respir. 17:437-446).

Eppin (Epididymal protease inhibitor) is a family of protease inhibitors expressed in the epididymis and testis. Two eppin isoforms contain both Kunitz-type and WAP-type four disulfide core protease inhibitor consensus sequences. Eppin-1 is expressed only in the testis and epididymis; Eppin-2 is expressed only in the epididymis and Eppin-3 only in the testis (Richardson, R.T. et al. (2001) Gene 270:93-102).

Human cystatin C is a potent inhibitor of cysteine proteases. Further, it has amyloidogenic properties. It refolds to produce very tight two-fold symmetric dimers while retaining the secondary structure of the monomeric form. The structure suggests a mechanism for its aggregation in the brain arteries of elderly people with amyloid angiopathy. A more severe 'conformational disease' is associated with the L68Q mutant of human cystatin C, which causes massive amyloidosis, cerebral hemorrhage, and death in young adults (Janowski, R. et al. (2001) Nat. Struct. Biol. 8(4):316-20).

A major portion of all proteins synthesized in eukaryotic cells are synthesized on the cytosolic surface of the endoplasmic reticulum (ER). Before these immature proteins are distributed to other organelles in the cell or are secreted, they must be transported into the interior lumen of the ER where post-translational modifications are performed. These modifications include protein folding and the formation of disulfide bonds, and N-linked glycosylations.

Protein Isomerases

Protein folding in the ER is aided by two principal types of protein isomerases, protein disulfide isomerase (PDI), and peptidyl-prolyl isomerase (PPI). PDI catalyzes the oxidation of free sulfhydryl groups in cysteine residues to form intramolecular disulfide bonds in proteins. PPI, an enzyme that catalyzes the isomerization of certain proline imidic bonds in oligopeptides and proteins, is considered to govern one of the rate limiting steps in the folding of many proteins to their final functional conformation. The cyclophilins represent a major class of PPI that was originally identified as the major receptor for the immunosuppressive drug cyclosporin A (Handschumacher, R.E. et al. (1984) Science 226: 544-547).

Protein Glycosylation

The glycosylation of most soluble secreted and membrane-bound proteins by oligosaccharides linked to asparagine residues in proteins is also performed in the ER. This reaction is catalyzed by a membrane-bound enzyme, oligosaccharyl transferase. Although the exact purpose of this "N-linked" glycosylation is unknown, the presence of oligosaccharides tends to make a

glycoprotein resistant to protease digestion. In addition, oligosaccharides attached to cell-surface proteins called selectins are known to function in cell-cell adhesion processes (Alberts, B. et al. (1994) Molecular Biology of the Cell Garland Publishing Co., New York, NY. p.608). "O-linked" glycosylation of proteins also occurs in the ER by the addition of N-acetylgalactosamine to the hydroxyl group of a serine or threonine residue followed by the sequential addition of other sugar residues to the first. This process is catalyzed by a series of glycosyltransferases each specific for a particular donor sugar nucleotide and acceptor molecule (Lodish, H. et al. (1995) Molecular Cell Biology, W. H. Freeman and Co., New York, NY pp.700-708). In many cases, both - and O-linked oligosaccharides appear to be required for the secretion of proteins or the movement of plasma membrane glycoproteins to the cell surface. For example, one of the glycosyltransferases in the dolichol pathway, dolichol phosphate mannose synthase,, is required in N:-glycosylation, O-mannosylation, and glycosylphosphatidylinositol membrane anchoring of protein (Tomita, S. et al. (1998) J. Biol. Chem. 9249-9254). Thus, in many cases, both N- and O-linked oligosaccharides appear to be required for the secretion of proteins or the movement of plasma membrane glycoproteins to the cell surface.

An additional glycosylation mechanism operates in the ER specifically to target lysosomal enzymes to lysosomes and prevent their secretion. Lysosomal enzymes in the ER receive an N-linked oligosaccharide, like plasma membrane and secreted proteins, but are then phosphorylated on one or two mannose residues. The phosphorylation of mannose residues occurs in two steps, the first step being the addition of an N-acetylglucosamine phosphate residue by N-acetylglucosamine phosphotransferase, and the second the removal of the N-acetylglucosamine group by phosphodiesterase. The phosphorylated mannose residue then targets the lysosomal enzyme to a mannose 6-phosphate receptor which transports it to a lysosome vesicle (Lodish et al. supra, pp. 708-711).

Chaperones

Molecular chaperones are proteins that aid in the proper folding of immature proteins and refolding of improperly folded ones, the assembly of protein subunits, and in the transport of unfolded proteins across membranes. Chaperones are also called heat-shock proteins (hsp) because of their tendency to be expressed in dramatically increased amounts following brief exposure of cells to elevated temperatures. This latter property most likely reflects their need in the refolding of proteins that have become denatured by the high temperatures. Chaperones may be divided into several classes according to their location, function, and molecular weight, and include hsp60, TCP1, hsp70, hsp40 (also called DnaJ), and hsp90. For example, hsp90 binds to steroid hormone receptors,

represses transcription in the absence of the ligand, and provides proper folding of the ligand-binding domain of the receptor in the presence of the hormone (Burstion, S.G. and A.R. Clarke (1995) *Essays Biochem.* 29:125-136). Hsp60 and hsp70 chaperones aid in the transport and folding of newly synthesized proteins. Hsp70 acts early in protein folding, binding a newly synthesized protein before it leaves the ribosome and transporting the protein to the mitochondria or ER before releasing the folded protein. Hsp60, along with hsp10, binds misfolded proteins and gives them the opportunity to refold correctly. All chaperones share an affinity for hydrophobic patches on incompletely folded proteins and the ability to hydrolyze ATP. The energy of ATP hydrolysis is used to release the hsp-bound protein in its properly folded state (Alberts, B. et al. supra, pp 214, 571-572).

Dipeptidyl-peptidase I, a lysosomal cysteine proteinase, is important in intracellular degradation of proteins and appears to be a central coordinator for activation of many serine proteinases in immune/inflammatory cells. The gene has been mapped to chromosomal region 11q14.1-q14.3. Dipeptidyl-peptidase I is expressed at high levels in lung, kidney, and placenta, and also at high levels in polymorphonuclear leukocytes and alveolar macrophages and their precursor cells (Rao, N.V. et al. (1997) *J. Biol. Chem.* 272:10260-10265).

IAP is a protein family that has baculovirus IAP repeat (BIR) domains and inhibits apoptosis. A human IAP family gene, Apollon, encodes a 530 kDa protein that contains a single BIR domain and a ubiquitin-conjugating enzyme domain. Apollon has been observed to protect cells from undergoing apoptosis and implicated in tumorigenesis and drug resistance (Chen, Z. et al. (1999) *Biochem. Biophys. Res. Commun.* 264:847-854).

The RTVL-H family is a medium repetitive family of endogenous retrovirus-like sequences found in the genomes of humans and other primates. Different subfamilies of RTVL-H elements are designated Type I, Type Ia, and Type II (Goodchild, N.L. (1993) *Virology* 196:778-788).

Lysyl Hydroxylases

Lysyl hydroxylase is an enzyme involved in collagen biosynthesis. Collagens are a family of fibrous structural proteins that are found in essentially all tissues. Collagens are the most abundant proteins in mammals, and are essential for the formation of connective tissue such as skin, bone, tendon, cartilage, blood vessels and teeth. Members of the collagen family can be distinguished from one another by the degree of cross-linking between collagen fibers and by the number of carbohydrate units (e.g., galactose or glucosylgalactose) attached to the collagen fibers. Hydroxylated lysine residues (hydroxylysine) are essential for stability of cross-linking and as attachment points for carbohydrate units.

The enzyme lysyl hydroxylase catalyzes the hydroxylation of lysine residues to form hydroxylysine. Lysyl hydroxylase targets the lysine residue of the sequence, X-lys-gly (lys = lysine, gly = glycine, and X = any amino acid residue). Three isoforms of lysyl hydroxylase have been

characterized, termed LH1 (or PLOD; procollagen-lysine, 2-oxoglutarate 5-dioxygenase), LH2 (or PLOD2), and LH3. The three enzymes share 60% sequence identity overall, with even higher similarity in the C-terminal region. In addition, there are regions in the middle of the molecule that have an identity of more than 80% (Valtavaara, M. et al. (1998) *J. Biol. Chem.* 273:12881-12886).

- 5 Diminished lysyl hydroxylase activity is involved in certain connective tissue disorders. In particular mutations, including a truncation and duplications within the coding region of the gene for PLOD, have been described in patients with type VI Ehlers-Danos syndrome (Hyland, J. et al. (1992) *Nature Genet.* 2:228-31; Hautala, T. et al. (1993) *Genomics* 15:399-404).

Ubiquitin-Associated Proteins

- 10 The ubiquitin conjugation system (UCS), is a major pathway for the degradation of cellular proteins of all types, including proteins that function to activate or repress cellular processes such as transcription, cell cycle progression, and immune recognition (Ciechanover, A. (1994) *Cell* 79:13-21). The process of ubiquitin conjugation and protein degradation involves several steps (Jentsch, S. (1992) *Annu. Rev. Genet.* 26:179-207). First ubiquitin (Ub), a small, heat stable protein is activated
15 by a ubiquitin-activating enzyme (E1) in an ATP dependent reaction which binds the C-terminus of Ub to the thiol group of an internal cysteine residue in E1. Activated Ub is then transferred to one of several Ub-conjugating enzymes (E2). Different ubiquitin-dependent proteolytic pathways employ structurally similar, but distinct ubiquitin-conjugating enzymes that are associated with recognition subunits which direct them to proteins carrying a particular degradation signal. E2 then transfers the
20 Ub molecule through its C-terminal glycine to a member of the ubiquitin-protein ligase family, E3. Next, E3 transfers the Ub molecule to the target protein. Additional Ub molecules may be added to the target protein forming a multi-Ub chain structure. The ubiquitinated protein is then recognized and degraded by the proteasome, an intracellular protease complex found in some bacteria and in all eukaryotic cells. The resultant ubiquitin-peptide complex is hydrolyzed by a ubiquitin carboxyl
25 terminal hydrolase, and free ubiquitin is released for reutilization by the UCS.

- Ubiquitin-proteasome systems are implicated in the degradation of mitotic cyclic kinases, oncoproteins, tumor suppressor genes (p53), cell surface receptors associated with signal transduction, transcriptional regulators, and mutated or damaged proteins (Ciechanover, supra). This pathway has been implicated in a number of diseases, including cystic fibrosis, Angelman's
30 syndrome, and Liddle syndrome (reviewed in Schwartz, A.L. and A. Ciechanover (1999) *Annu. Rev. Med.* 50:57-74). A murine proto-oncogene, Unp, encodes a nuclear ubiquitin protease whose overexpression leads to oncogenic transformation of NIH3T3 cells. The human homologue of this gene is consistently elevated in small cell tumors and adenocarcinomas of the lung (Gray, D.A. (1995) *Oncogene* 10:2179-2183). Ubiquitin carboxyl terminal hydrolase is involved in the
35 differentiation of a lymphoblastic leukemia cell line to a non-dividing mature state (Maki, A. et al.

(1996) Differentiation 60:59-66). In neurons, ubiquitin carboxyl terminal hydrolase (PGP 9.5) expression is strong in the abnormal structures that occur in human neurodegenerative diseases (Lowe, J. et al. (1990) J. Pathol. 161:153-160).

Additional ubiquitin-like proteins which also possess the ability to covalently modify other cellular proteins have been identified in recent years. (For review, see Yeh, E.T.H. et al. (2000) Gene 248:1-14; and Jentsch, S. and Pyrowolakis, G. (2000) Trends Cell Biol. 10:335-342.) These ubiquitin-like protein modifiers include the sentrins (also known as SUMO proteins), NEDD8, and Apg12. The conjugation pathways for these proteins closely resemble that for ubiquitin. For example, conjugation of sentrin requires the E1 heterodimer AOS1/UBA2, and a single E2 enzyme, UBC9. The recently discovered protein S3 may function as a sentrin ligase. The yeast protein Ulp1 is a sentrin hydrolase. Inactivation of Ulp1 in yeast results in severe cell cycle defects. In humans, seven sentrin specific proteases (SENP) have been identified, which range in size from 238 to 1112 amino acid residues (Yeh, supra). All human SENPs share a conserved C-terminal domain. The N-terminal regions may regulate cellular location and substrate specificity.

Sentrinization does not promote protein degradation as does ubiquitin. In some cases sentrinization appears to be important for stable localization of target proteins in nuclear bodies. Substrates for sentrinization include PML, a RING finger protein with tumor suppressor activity, HIPK2, a co-repressor for homeodomain transcription factors, and the tumor suppressor p53. I κ B α , a cytosolic inhibitor of NF κ B, a transcription factor involved in induction of inflammation associated proteins, is also a substrate for sentrinization. Sentrinized I κ B α cannot be ubiquitinated and is resistant to proteasomal degradation, suggesting links between the ubiquitin and sentrin pathways. Jentsch, supra).

Expression profiling

Microarrays are analytical tools used in bioanalysis. A microarray has a plurality of molecules spatially distributed over, and stably associated with, the surface of a solid support. Microarrays of polypeptides, polynucleotides, and/or antibodies have been developed and find use in a variety of applications, such as gene sequencing, monitoring gene expression, gene mapping, bacterial identification, drug discovery, and combinatorial chemistry.

One area in particular in which microarrays find use is in gene expression analysis. Array technology can provide a simple way to explore the expression of a single polymorphic gene or the expression profile of a large number of related or unrelated genes. When the expression of a single gene is examined, arrays are employed to detect the expression of a specific gene or its variants. When an expression profile is examined, arrays provide a platform for identifying genes that are tissue specific, are affected by a substance being tested in a toxicology assay, are part of a signaling

cascade, carry out housekeeping functions, or are specifically related to a particular genetic predisposition, condition, disease, or disorder.

Steroids affecting protein modification

Steroids are a class of lipid-soluble molecules, including cholesterol, bile acids, vitamin D, and hormones, that share a common four-ring structure based on cyclopentanoperhydrophenanthrene and that carry out a wide variety of functions. Cholesterol, for example, is a component of cell membranes that controls membrane fluidity. It is also a precursor for bile acids which solubilize lipids and facilitate absorption in the small intestine during digestion. Vitamin D regulates the absorption of calcium in the small intestine and controls the concentration of calcium in plasma.

Steroid hormones, produced by the adrenal cortex, ovaries, and testes, include glucocorticoids, mineralocorticoids, androgens, and estrogens. They control various biological processes by binding to intracellular receptors that regulate transcription of specific genes in the nucleus. Glucocorticoids, for example, increase blood glucose concentrations by regulation of gluconeogenesis in the liver, increase blood concentrations of fatty acids by promoting lipolysis in adipose tissues, modulate sensitivity to catecholamines in the central nervous system, and reduce inflammation. The principal mineralocorticoid, aldosterone, is produced by the adrenal cortex and acts on cells of the distal tubules of the kidney to enhance sodium ion reabsorption. Androgens, produced by the interstitial cells of Leydig in the testis, include the male sex hormone testosterone, which triggers changes at puberty, the production of sperm and maintenance of secondary sexual characteristics. Female sex hormones, estrogen and progesterone, are produced by the ovaries and also by the placenta and adrenal cortex of the fetus during pregnancy. Estrogen regulates female reproductive processes and secondary sexual characteristics. Progesterone regulates changes in the endometrium during the menstrual cycle and pregnancy.

Steroid hormones are widely used for fertility control and in anti-inflammatory treatments for physical injuries and diseases such as arthritis, asthma, and auto-immune disorders. Progesterone, a naturally occurring progestin, is primarily used to treat amenorrhea, abnormal uterine bleeding, or as a contraceptive. Endogenous progesterone is responsible for inducing secretory activity in the endometrium of the estrogen-primed uterus in preparation for the implantation of a fertilized egg and for the maintenance of pregnancy. It is secreted from the corpus luteum in response to luteinizing hormone (LH). The primary contraceptive effect of exogenous progestins involves the suppression of the midcycle surge of LH. At the cellular level, progestins diffuse freely into target cells and bind to the progesterone receptor. Target cells include the female reproductive tract, the mammary gland, the hypothalamus, and the pituitary. Once bound to the receptor, progestins slow the frequency of release of gonadotropin releasing hormone from the hypothalamus and blunt the pre-ovulatory LH surge, thereby preventing follicular maturation and ovulation. Progesterone has minimal estrogenic

and androgenic activity. Progesterone is metabolized hepatically to pregnanediol and conjugated with glucuronic acid.

Medroxyprogesterone (MAH), also known as 6 α -methyl-17-hydroxyprogesterone, is a synthetic progestin with a pharmacological activity about 15 times greater than progesterone. MAH is used for the treatment of renal and endometrial carcinomas, amenorrhea, abnormal uterine bleeding, and endometriosis associated with hormonal imbalance. MAH has a stimulatory effect on respiratory centers and has been used in cases of low blood oxygenation caused by sleep apnea, chronic obstructive pulmonary disease, or hypercapnia.

Mifepristone, also known as RU-486, is an antiprogesterone drug that blocks receptors of progesterone. It counteracts the effects of progesterone, which is needed to sustain pregnancy. Mifepristone induces spontaneous abortion when administered in early pregnancy followed by treatment with the prostaglandin, misoprostol. Further, studies show that mifepristone at a substantially lower dose can be highly effective as a postcoital contraceptive when administered within five days after unprotected intercourse, thus providing women with a "morning-after pill" in case of contraceptive failure or sexual assault. Mifepristone also has potential uses in the treatment of breast and ovarian cancers in cases in which tumors are progesterone-dependent. It interferes with steroid-dependent growth of brain meningiomas, and may be useful in treatment of endometriosis where it blocks the estrogen-dependent growth of endometrial tissues. It may also be useful in treatment of uterine fibroid tumors and Cushing's Syndrome. Mifepristone binds to glucocorticoid receptors and interferes with cortisol binding. Mifepristone also may act as an anti-glucocorticoid and be effective for treating conditions where cortisol levels are elevated such as AIDS, anorexia nervosa, ulcers, diabetes, Parkinson's disease, multiple sclerosis, and Alzheimer's disease.

Danazol is a synthetic steroid derived from ethinyl testosterone. Danazol indirectly reduces estrogen production by lowering pituitary synthesis of follicle-stimulating hormone and LH. Danazol also binds to sex hormone receptors in target tissues, thereby exhibiting anabolic, antiestrogenic, and weakly androgenic activity. Danazol does not possess any progestogenic activity, and does not suppress normal pituitary release of corticotropin or release of cortisol by the adrenal glands. Danazol is used in the treatment of endometriosis to relieve pain and inhibit endometrial cell growth. It is also used to treat fibrocystic breast disease and hereditary angioedema.

Corticosteroids are used to relieve inflammation and to suppress the immune response. They inhibit eosinophil, basophil, and airway epithelial cell function by regulation of cytokines that mediate the inflammatory response. They inhibit leukocyte infiltration at the site of inflammation, interfere in the function of mediators of the inflammatory response, and suppress the humoral immune response. Corticosteroids are used to treat allergies, asthma, arthritis, and skin conditions. Beclomethasone is a synthetic glucocorticoid that is used to treat steroid-dependent asthma, to relieve

symptoms associated with allergic or nonallergic (vasomotor) rhinitis, or to prevent recurrent nasal polyps following surgical removal. The anti-inflammatory and vasoconstrictive effects of intranasal beclomethasone are 5000 times greater than those produced by hydrocortisone. Budesonide is a corticosteroid used to control symptoms associated with allergic rhinitis or asthma. Budesonide has high topical anti-inflammatory activity but low systemic activity. Dexamethasone is a synthetic glucocorticoid used in anti-inflammatory or immunosuppressive compositions. It is also used in inhalants to prevent symptoms of asthma. Due to its greater ability to reach the central nervous system, dexamethasone is usually the treatment of choice to control cerebral edema. Dexamethasone is approximately 20-30 times more potent than hydrocortisone and 5-7 times more potent than prednisone. Prednisone is metabolized in the liver to its active form, prednisolone, a glucocorticoid with anti-inflammatory properties. Prednisone is approximately 4 times more potent than hydrocortisone and the duration of action of prednisone is intermediate between hydrocortisone and dexamethasone. Prednisone is used to treat allograft rejection, asthma, systemic lupus erythematosus, arthritis, ulcerative colitis, and other inflammatory conditions. Betamethasone is a synthetic glucocorticoid with antiinflammatory and immunosuppressive activity and is used to treat psoriasis and fungal infections, such as athlete's foot and ringworm.

The anti-inflammatory actions of corticosteroids are thought to involve phospholipase A₂ inhibitory proteins, collectively called lipocortins. Lipocortins, in turn, control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of the precursor molecule arachidonic acid. Proposed mechanisms of action include decreased IgE synthesis, increased number of β -adrenergic receptors on leukocytes, and decreased arachidonic acid metabolism. During an immediate allergic reaction, such as in chronic bronchial asthma, allergens bridge the IgE antibodies on the surface of mast cells, which triggers these cells to release chemotactic substances. Mast cell influx and activation, therefore, is partially responsible for the inflammation and hyperirritability of the oral mucosa in asthmatic patients. This inflammation can be retarded by administration of corticosteroids.

Toxicology Testing:

Array technology can provide a simple way to explore the expression of a single polymorphic gene or the expression profile of a large number of related or unrelated genes. When the expression of a single gene is examined, arrays are employed to detect the expression of a specific gene or its variants. When an expression profile is examined, arrays provide a platform for examining which genes are tissue specific, carry out housekeeping functions, are parts of a signaling cascade, or are specifically related to a particular genetic predisposition, condition, disease, or disorder.

The potential application of gene expression profiling is particularly relevant to improving diagnosis, prognosis, and treatment of disease. For example, both the levels and sequences expressed

in tissues from subjects with hyperlipidemia may be compared with the levels and sequences expressed in normal tissue.

Toxicity testing is a mandatory and time-consuming part of drug development programs in the pharmaceutical industry. A more rapid screen to determine the effects upon metabolism and to
5 detect toxicity of lead drug candidates may be the use of gene expression microarrays. For example, microarrays of various kinds may be produced using full length genes or gene fragments. These arrays can then be used to test samples treated with the drug candidates to elucidate the gene expression pattern associated with drug treatment. This gene pattern can be compared with gene expression patterns associated with compounds which produce known metabolic and toxicological
10 responses.

The human C3A cell line is a clonal derivative of HepG2/C3 (hepatoma cell line, isolated from a 15-year-old male with liver tumor), which was selected for strong contact inhibition of growth. The use of a clonal population enhances the reproducibility of the cells. C3A cells have many characteristics of primary human hepatocytes in culture: i) expression of insulin receptor and
15 insulin-like growth factor II receptor; ii) secretion of a high ratio of serum albumin compared with α -fetoprotein iii) conversion of ammonia to urea and glutamine; iv) metabolism of aromatic amino acids; and v) ability to proliferate in glucose-free and insulin-free medium. The C3A cell line is now well established as an *in vitro* model of the mature human liver (Mickelson et al. (1995) *Hepatology* 22:866-875; Nagendra et al. (1997) *Am. J. Physiol.* 272:G408-416).

Clofibrate is an hypolipidemic drug which lowers elevated levels of serum triglycerides. In
20 rodents, chronic treatment produces hepatomegaly and an increase in hepatic peroxisomes (peroxisome proliferation). Peroxisome proliferators (PPs) are a class of drugs which activate the PP-activated receptor in rodent liver, leading to enzyme induction, stimulation of S-phase, and a suppression of apoptosis (Hasmall and Roberts (1999) *Pharmacol. Ther.* 82:63-70). PPs include the
25 fibrate class of hypolipidemic drugs, phenobarbitone, thiazolidinediones, certain non-steroidal anti-inflammatory drugs, and naturally-occurring fatty acid-derived molecules (Gelman *et al.* (1999) *Cell. Mol. Life Sci.* 55:932-943). Clofibrate has been shown to increase levels of cytochrome P450 4A. It is also involved in transcription of β -oxidation genes as well as induction of PP-activated receptors (Kawashima *et al.* (1997) *Arch. Biochem. Biophys.* 347:148-154). Peroxisome proliferation that is
30 induced by both clofibrate and the chemically-related compound fenofibrate is mediated by a common inhibitory effect on mitochondrial membrane depolarization (Zhou and Wallace (1999) *Toxicol. Sci.* 48:82-89).

Dexamethasone and its derivatives, dexamethasone sodium phosphate and dexamethasone acetate, are synthetic glucocorticoids used as anti-inflammatory or immunosuppressive agents.
35 Dexamethasone has little to no mineralocorticoid activity and is usually selected for management of

cerebral edema because of its superior ability to penetrate the central nervous system. Glucocorticoids are naturally occurring hormones that prevent or suppress inflammation and immune responses when administered at pharmacological doses. Responses can include inhibition of leukocyte infiltration at the site of inflammation, interference in the function of mediators of inflammatory response, and suppression of humoral immune responses. The anti-inflammatory actions of corticosteroids are thought to involve phospholipase A₂ inhibitory proteins, collectively called lipocortins. The numerous adverse effects related to corticosteroid use usually depend on the dose administered and the duration of therapy. Proposed mechanisms of action include decreased IgE synthesis, increased number of β -adrenergic receptors on leukocytes, and decreased arachidonic acid metabolism. During an immediate allergic reaction, such as in chronic bronchial asthma, allergens bridge the IgE antibodies on the surface of mast cells, which triggers these cells to release chemotactic substances. Mast cell influx and activation, therefore, is partially responsible for the inflammation and hyperirritability of the oral mucosa in asthmatic patients. This inflammation can be retarded by administration of adrenocorticoids. As with other corticosteroids, the effects upon liver metabolism and hormone clearance mechanisms are important to understand the pharmacodynamics of a drug.

Cancer

Prostate cancer develops through a multistage progression ultimately resulting in an aggressive tumor phenotype. The initial step in tumor progression involves the hyperproliferation of normal luminal and/or basal epithelial cells. Androgen responsive cells become hyperplastic and evolve into early-stage tumors. Although early-stage tumors are often androgen sensitive and respond to androgen ablation, a population of androgen independent cells evolve from the hyperplastic population. These cells represent a more advanced form of prostate tumor that may become invasive and potentially become metastatic to the bone, brain, or lung.

Breast cancer develops through a multi-step process in which pre-malignant mammary epithelial cells undergo a relatively defined sequence of events leading to tumor formation. An early event in tumor development is ductal hyperplasia. Cells undergoing rapid neoplastic growth gradually progress to invasive carcinoma and become metastatic to the lung, bone, and potentially other organs. Several variables that may influence the process of tumor progression and malignant transformation include genetic factors, environmental factors, growth factors, and hormones. Based on the complexity of this process, it is critical to study a population of human mammary epithelial cells undergoing the process of malignant transformation, and to associate specific stages of progression with phenotypic and molecular characteristics.

Immune response proteins

Interleukin 12 (IL-12) is a pleiotropic cytokine produced by macrophages and B lymphocytes that can have multiple effects on T cells and natural killer (NK) cells. Effects include inducing

production of IFN- γ and TNF by resting and activated T and NK cells; enhancing the cytotoxic activity of resting NK and T cells, inducing and synergizing with IL-2 in the generation of lymphokine-activated killer (LAK) cells; acting as a comitogen to stimulate proliferation of resting T cells; and inducing proliferation of activated T and NK cells. Current evidence indicates that IL-12, produced by macrophages in response to infectious agents, is a central mediator of the cell-mediated immune response by its actions on the development, proliferation, and activities of TH1 cells. As the initiator of cell-mediated immunity, IL-12 may stimulate cell-mediated immune responses to microbial pathogens, metastatic cancers, and viral infections such as AIDS.

There is a need in the art for new compositions, including nucleic acids and proteins, for the diagnosis, prevention, and treatment of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders.

SUMMARY OF THE INVENTION

Various embodiments of the invention provide purified polypeptides, protein modification and maintenance molecules, referred to collectively as "PMOD" and individually as "PMOD-1," "PMOD-2," "PMOD-3," "PMOD-4," "PMOD-5," "PMOD-6," "PMOD-7," "PMOD-8," "PMOD-9," "PMOD-10," "PMOD-11," "PMOD-12," "PMOD-13," "PMOD-14," "PMOD-15," "PMOD-16," "PMOD-17," "PMOD-18," "PMOD-19," "PMOD-20," "PMOD-21," "PMOD-22," "PMOD-23," "PMOD-24," "PMOD-25," "PMOD-26," "PMOD-27," and "PMOD-28," and methods for using these proteins and their encoding polynucleotides for the detection, diagnosis, and treatment of diseases and medical conditions. Embodiments also provide methods for utilizing the purified protein modification and maintenance molecules and/or their encoding polynucleotides for facilitating the drug discovery process, including determination of efficacy, dosage, toxicity, and pharmacology. Related embodiments provide methods for utilizing the purified protein modification and maintenance molecules and/or their encoding polynucleotides for investigating the pathogenesis of diseases and medical conditions.

An embodiment provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. Another embodiment provides an isolated polypeptide comprising an amino acid sequence of SEQ ID NO:1-28.

Still another embodiment provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. In another embodiment, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-28. In an alternative embodiment, the polynucleotide is selected from the group consisting of SEQ ID NO:29-56.

Still another embodiment provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. Another embodiment provides a cell transformed with the recombinant polynucleotide. Yet another embodiment provides a transgenic organism comprising the recombinant polynucleotide.

Another embodiment provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Yet another embodiment provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an

amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28.

5 Still yet another embodiment provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, c) a polynucleotide complementary to the
10 polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In other embodiments, the polynucleotide can comprise at least about 20, 30, 40, 60, 80, or 100 contiguous nucleotides.

Yet another embodiment provides a method for detecting a target polynucleotide in a sample, said target polynucleotide being selected from the group consisting of a) a polynucleotide comprising
15 a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method
20 comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex. In a related embodiment, the
25 method can include detecting the amount of the hybridization complex. In still other embodiments, the probe can comprise at least about 20, 30, 40, 60, 80, or 100 contiguous nucleotides.

Still yet another embodiment provides a method for detecting a target polynucleotide in a sample, said target polynucleotide being selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, b) a
30 polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain
35 reaction amplification, and b) detecting the presence or absence of said amplified target

polynucleotide or fragment thereof. In a related embodiment, the method can include detecting the amount of the amplified target polynucleotide or fragment thereof.

Another embodiment provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and a pharmaceutically acceptable excipient. In one embodiment, the composition can comprise an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. Other embodiments provide a method of treating a disease or condition associated with decreased or abnormal expression of functional PMOD, comprising administering to a patient in need of such treatment the composition.

Yet another embodiment provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. Another embodiment provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. Yet another embodiment provides a method of treating a disease or condition associated with decreased expression of functional PMOD, comprising administering to a patient in need of such treatment the composition.

Still yet another embodiment provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. The method comprises a)

exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. Another embodiment provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. Yet another embodiment provides a method of treating a disease or condition associated with overexpression of functional
5 PMOD, comprising administering to a patient in need of such treatment the composition.

Another embodiment provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an
10 amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the
15 polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

Yet another embodiment provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, b) a polypeptide comprising a
20 naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28. The method comprises a) combining the
25 polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the
30 polypeptide.

Still yet another embodiment provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, b) detecting
35 altered expression of the target polynucleotide, and c) comparing the expression of the target

polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

Another embodiment provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide can comprise a fragment of a polynucleotide selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for full length polynucleotide and polypeptide embodiments of the invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog, and the PROTEOME database identification numbers and annotations of PROTEOME database homologs, for polypeptide embodiments of the invention. The probability scores for the matches between each polypeptide and its homolog(s) are also shown.

Table 3 shows structural features of polypeptide embodiments, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide embodiments, along with selected fragments of the polynucleotides.

Table 5 shows representative cDNA libraries for polynucleotide embodiments.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze polynucleotides and polypeptides, along with applicable descriptions, references, and threshold parameters.

Table 8 shows single nucleotide polymorphisms found in polynucleotide embodiments, along with allele frequencies in different human populations.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleic acids, and methods are described, it is understood that embodiments of the invention are not limited to the particular machines, instruments, materials, and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the invention.

As used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with various embodiments of the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"PMOD" refers to the amino acid sequences of substantially purified PMOD obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of PMOD. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other

compound or composition which modulates the activity of PMOD either by directly interacting with PMOD or by acting on components of the biological pathway in which PMOD participates.

An "allelic variant" is an alternative form of the gene encoding PMOD. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding PMOD include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as PMOD or a polypeptide with at least one functional characteristic of PMOD. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding PMOD, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide encoding PMOD. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent PMOD. Deliberate amino acid substitutions may be made on the basis of one or more similarities in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of PMOD is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" can refer to an oligopeptide, a peptide, a polypeptide, or a protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid. Amplification may be carried out using polymerase chain reaction (PCR) technologies or other nucleic acid amplification technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of PMOD. Antagonists may include proteins such as antibodies, anticalins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PMOD either by directly interacting with PMOD or by acting on components of the biological pathway in which PMOD participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind PMOD polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "aptamer" refers to a nucleic acid or oligonucleotide molecule that binds to a specific molecular target. Aptamers are derived from an *in vitro* evolutionary process (e.g., SELEX (Systematic Evolution of Ligands by EXponential Enrichment), described in U.S. Patent No. 5,270,163), which selects for target-specific aptamer sequences from large combinatorial libraries. Aptamer compositions may be double-stranded or single-stranded, and may include deoxyribonucleotides, ribonucleotides, nucleotide derivatives, or other nucleotide-like molecules. The nucleotide components of an aptamer may have modified sugar groups (e.g., the 2'-OH group of a ribonucleotide may be replaced by 2'-F or 2'-NH₂), which may improve a desired property, e.g., resistance to nucleases or longer lifetime in blood. Aptamers may be conjugated to other molecules, e.g., a high molecular weight carrier to slow clearance of the aptamer from the circulatory system. Aptamers may be specifically cross-linked to their cognate ligands, e.g., by photo-activation of a cross-linker. (See, e.g., Brody, E.N. and L. Gold (2000) J. Biotechnol. 74:5-13.)

The term "intramer" refers to an aptamer which is expressed *in vivo*. For example, a vaccinia virus-based RNA expression system has been used to express specific RNA aptamers at high levels in the cytoplasm of leukocytes (Blind, M. et al. (1999) Proc. Natl. Acad. Sci. USA 96:3606-3610).

The term "spiegelmer" refers to an aptamer which includes L-DNA, L-RNA, or other left-handed nucleotide derivatives or nucleotide-like molecules. Aptamers containing left-handed nucleotides are resistant to degradation by naturally occurring enzymes, which normally act on substrates containing right-handed nucleotides.

5 The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a polynucleotide having a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or
10 oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand,
15 and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

 The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic PMOD, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific
20 antibodies.

 "Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

 A "composition comprising a given polynucleotide" and a "composition comprising a given
25 polypeptide" can refer to any composition containing the given polynucleotide or polypeptide. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotides encoding PMOD or fragments of PMOD may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts
30 (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

 "Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been
35 assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer

program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

“Conservative amino acid substitutions” are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
10	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
15	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
20	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
25	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A “deletion” refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term “derivative” refers to a chemically modified polynucleotide or polypeptide.

Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A “detectable label” refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

“Differential expression” refers to increased or upregulated; or decreased, downregulated, or absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

“Exon shuffling” refers to the recombination of different coding regions (exons). Since an exon may represent a structural or functional domain of the encoded protein, new proteins may be assembled through the novel reassortment of stable substructures, thus allowing acceleration of the evolution of new protein functions.

A “fragment” is a unique portion of PMOD or a polynucleotide encoding PMOD which can be identical in sequence to, but shorter in length than, the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from about 5 to about 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:29-56 can comprise a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:29-56, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:29-56 can be employed in one or more embodiments of methods of the invention, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:29-56 from related polynucleotides. The precise length of a fragment of SEQ ID NO:29-56 and the region of SEQ ID NO:29-56 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-28 is encoded by a fragment of SEQ ID NO:29-56. A fragment of SEQ ID NO:1-28 can comprise a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-28. For example, a fragment of SEQ ID NO:1-28 can be used as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-28. The precise length of a fragment of SEQ ID NO:1-28 and the region of SEQ ID NO:1-28 to which the fragment

corresponds can be determined based on the intended purpose for the fragment using one or more analytical methods described herein or otherwise known in the art.

A "full length" polynucleotide is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using one or more computer algorithms or programs known in the art or described herein. For example, percent identity can be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms which can be used is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version

2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

5 *Open Gap: 5 and Extension Gap: 2 penalties*

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

10 Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length
15 supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid
20 sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative
25 substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of
30 polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise
35 comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version

2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

5 *Gap x drop-off: 50*

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term “humanized antibody” refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about

5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al.

- 5 (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC
10 concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions
15 will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acids by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex
20 may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid present in solution and another nucleic acid immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or polynucleotide
25 sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

30 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of PMOD which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of PMOD which is useful in any of the antibody production methods disclosed herein or known in the art.

35 The term "microarray" refers to an arrangement of a plurality of polynucleotides,

polypeptides, antibodies, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, antibody, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of PMOD. For example, modulation
5 may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of PMOD.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the
10 antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where
15 necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript
20 elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an PMOD may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of PMOD.

"Probe" refers to nucleic acids encoding PMOD, their complements, or fragments thereof,
25 which are used to detect identical, allelic or related nucleic acids. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide
30 by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme.

Primer pairs can be used for amplification (and identification) of a nucleic acid, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous
35 nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also

be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

- 5 Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs
10 can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

- Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to
15 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from
20 megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection
25 programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both
30 unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a nucleic acid that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA molecule, is composed of the same linear sequence of nucleotides as the reference DNA molecule with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing PMOD, nucleic acids encoding PMOD, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably at least about 75% free, and most preferably at least about 90% free from other components with which they are naturally associated.

5 A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells,
10 trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" or "expression profile" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods
15 well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term
"transformed cells" includes stably transformed cells in which the inserted DNA is capable of
20 replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the
25 art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. In another embodiment, the nucleic acid can be introduced by infection with a recombinant viral vector, such as a lentiviral vector (Lois, C. et al. (2002) Science 295:868-872). The term genetic manipulation does not include classical cross-breeding, or *in vitro* fertilization, but
30 rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided
35 in references such as Sambrook et al. (1989), *supra*.

A “variant” of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the “BLAST 2 Sequences” tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an “allelic” (as defined above), “splice,” “species,” or “polymorphic” variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotides that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass “single nucleotide polymorphisms” (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A “variant” of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the “BLAST 2 Sequences” tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

Various embodiments of the invention include new human protein modification and maintenance molecules (PMOD), the polynucleotides encoding PMOD, and the use of these compositions for the diagnosis, treatment, or prevention of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide embodiments of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is

denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown. Column 6 shows the Incyte ID numbers of physical, full length clones corresponding to polypeptide and polynucleotide embodiments. The full length clones encode polypeptides which have at least 95% sequence identity to the polypeptides shown in column 3.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database and the PROTEOME database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (GenBank ID NO:) of the nearest GenBank homolog and the PROTEOME database identification numbers (PROTEOME ID NO:) of the nearest PROTEOME database homologs. Column 4 shows the probability scores for the matches between each polypeptide and its homolog(s). Column 5 shows the annotation of the GenBank and PROTEOME database homolog(s) along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are protein modification and maintenance molecules. For example, SEQ ID NO:1 is 43% identical, from residue K223 to residue A774, to Arabidopsis thaliana ubiquitin-protein ligase 1 (GenBank ID g7108521) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.3e-85, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:1 also contains a HECT (ubiquitin-transferase) domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of

conserved protein family domains. (See Table 3.) Data from BLIMPS and BLAST analyses provide further corroborative evidence that SEQ ID NO:1 is a ubiquitin-protein ligase.

As another example, SEQ ID NO:5 is 38% identical, from residue E22 to residue K368, to Arabidopsis thaliana ubiquitin-specific protease 26 (GenBank ID g11993492) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $1.1e-79$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:5 also contains a ubiquitin carboxyl-terminal hydrolases 1 domain and a ubiquitin carboxyl-terminal hydrolases 2 domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and additional BLAST analyses provide further corroborative evidence that SEQ ID NO:5 is a ubiquitin carboxyl terminal hydrolase.

As another example, SEQ ID NO:7 is 91% identical, from residue P23 to residue S531, to a human carboxypeptidase N (GenBank ID g179936) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $1.7e-235$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:7 also contains leucine-rich repeat domains as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from MOTIFS analysis provides further corroborative evidence that SEQ ID NO:7 is a carboxypeptidase.

As another example, SEQ ID NO:10 is 46% identical, from residue R8 to residue S143, to mouse testatin, which is related to the cysteine protease inhibitors, cystatins (GenBank ID g3928491) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $1.4e-27$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:10 also contains a cystatin domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from further BLAST analysis provides corroborative evidence that SEQ ID NO:10 is a cysteine protease inhibitor.

As another example, SEQ ID NO:20 is 99% identical, from residue M17 to residue K267, to human kallikrein 14 (GenBank ID g13897995) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $2.1e-136$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:20 also contains a trypsin domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:20 is a serine protease.

As another example, SEQ ID NO:27 is 96% identical, from residue M1 to residue T242, to human putative mast cell mMCP-7-like II tryptase (GenBank ID g4336577) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.8e-130, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:27 also contains a trypsin domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:27 is a trypsin-like serine protease.

SEQ ID NO:2-4, SEQ ID NO:6, SEQ ID NO:8-9, SEQ ID NO:11-19, SEQ ID NO:21-26 and SEQ ID NO:28 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-28 are described in Table 7.

As shown in Table 4, the full length polynucleotide embodiments were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Column 1 lists the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:), the corresponding Incyte polynucleotide consensus sequence number (Incyte ID) for each polynucleotide of the invention, and the length of each polynucleotide sequence in basepairs. Column 2 shows the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences used to assemble the full length polynucleotide embodiments, and of fragments of the polynucleotides which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:29-56 or that distinguish between SEQ ID NO:29-56 and related polynucleotides.

The polynucleotide fragments described in Column 2 of Table 4 may refer specifically, for example, to Incyte cDNAs derived from tissue-specific cDNA libraries or from pooled cDNA libraries. Alternatively, the polynucleotide fragments described in column 2 may refer to GenBank cDNAs or ESTs which contributed to the assembly of the full length polynucleotides. In addition, the polynucleotide fragments described in column 2 may identify sequences derived from the ENSEMBL (The Sanger Centre, Cambridge, UK) database (*i.e.*, those sequences including the designation "ENST"). Alternatively, the polynucleotide fragments described in column 2 may be derived from the NCBI RefSeq Nucleotide Sequence Records Database (*i.e.*, those sequences including the designation "NM" or "NT") or the NCBI RefSeq Protein Sequence Records (*i.e.*, those sequences including the designation "NP"). Alternatively, the polynucleotide fragments described in column 2 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. For example, a polynucleotide sequence identified as FL_XXXXXX_N₁_N₂_YYYYY_N₃_N₄ represents a "stitched" sequence in which XXXXXX is the identification number of the cluster of sequences to which the algorithm was applied, and YYYYYY is

the number of the prediction generated by the algorithm, and $N_{1,2,3...}$, if present, represent specific exons that may have been manually edited during analysis (See Example V). Alternatively, the polynucleotide fragments in column 2 may refer to assemblages of exons brought together by an “exon-stretching” algorithm. For example, a polynucleotide sequence identified as

- 5 FLXXXXXX_gAAAAA_gBBBBB_1_N is a “stretched” sequence, with XXXXXX being the Incyte project identification number, gAAAAA being the GenBank identification number of the human genomic sequence to which the “exon-stretching” algorithm was applied, gBBBBB being the GenBank identification number or NCBI RefSeq identification number of the nearest GenBank protein homolog, and N referring to specific exons (See Example V). In instances where a RefSeq
- 10 sequence was used as a protein homolog for the “exon-stretching” algorithm, a RefSeq identifier (denoted by “NM,” “NP,” or “NT”) may be used in place of the GenBank identifier (*i.e.*, gBBBBB).

- Alternatively, a prefix identifies component sequences that were hand-edited, predicted from genomic DNA sequences, or derived from a combination of sequence analysis methods. The following Table lists examples of component sequence prefixes and corresponding sequence analysis
- 15 methods associated with the prefixes (see Example IV and Example V).

Prefix	Type of analysis and/or examples of programs
GNN, GFG, ENST	Exon prediction from genomic sequences using, for example, GENSCAN (Stanford University, CA, USA) or FGENES (Computer Genomics Group, The Sanger Centre, Cambridge, UK).
GBI	Hand-edited analysis of genomic sequences.
FL	Stitched or stretched genomic sequences (see Example V).
INCY	Full length transcript and exon prediction from mapping of EST sequences to the genome. Genomic location and EST composition data are combined to predict the exons and resulting transcript.

- In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in Table 4 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA
- 25 identification numbers are not shown.

- Table 5 shows the representative cDNA libraries for those full length polynucleotides which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotides. The tissues and vectors which were used to
- 30 construct the cDNA libraries shown in Table 5 are described in Table 6.

Table 8 shows single nucleotide polymorphisms (SNPs) found in polynucleotide embodiments, along with allele frequencies in different human populations. Columns 1 and 2 show

the polynucleotide sequence identification number (SEQ ID NO:) and the corresponding Incyte project identification number (PID) for polynucleotides of the invention. Column 3 shows the Incyte identification number for the EST in which the SNP was detected (EST ID), and column 4 shows the identification number for the SNP (SNP ID). Column 5 shows the position within the EST sequence at which the SNP is located (EST SNP), and column 6 shows the position of the SNP within the full-length polynucleotide sequence (CB1 SNP). Column 7 shows the allele found in the EST sequence. Columns 8 and 9 show the two alleles found at the SNP site. Column 10 shows the amino acid encoded by the codon including the SNP site, based upon the allele found in the EST. Columns 11-14 show the frequency of allele 1 in four different human populations. An entry of n/d (not detected) indicates that the frequency of allele 1 in the population was too low to be detected, while n/a (not available) indicates that the allele frequency was not determined for the population.

The invention also encompasses PMOD variants. A preferred PMOD variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the PMOD amino acid sequence, and which contains at least one functional or structural characteristic of PMOD.

Various embodiments also encompass polynucleotides which encode PMOD. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:29-56, which encodes PMOD. The polynucleotide sequences of SEQ ID NO:29-56, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses variants of a polynucleotide encoding PMOD. In particular, such a variant polynucleotide will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a polynucleotide encoding PMOD. A particular aspect of the invention encompasses a variant of a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NO:29-56 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:29-56. Any one of the polynucleotide variants described above can encode a polypeptide which contains at least one functional or structural characteristic of PMOD.

In addition, or in the alternative, a polynucleotide variant of the invention is a splice variant of a polynucleotide encoding PMOD. A splice variant may have portions which have significant sequence identity to a polynucleotide encoding PMOD, but will generally have a greater or lesser number of polynucleotides due to additions or deletions of blocks of sequence arising from alternate splicing of exons during mRNA processing. A splice variant may have less than about 70%, or

alternatively less than about 60%, or alternatively less than about 50% polynucleotide sequence identity to a polynucleotide encoding PMOD over its entire length; however, portions of the splice variant will have at least about 70%, or alternatively at least about 85%, or alternatively at least about 95%, or alternatively 100% polynucleotide sequence identity to portions of the polynucleotide encoding PMOD. For example, a polynucleotide comprising a sequence of SEQ ID NO:31 is a splice variant of a polynucleotide comprising a sequence of SEQ ID NO:34, and a polynucleotide comprising a sequence of SEQ ID NO:44 is a splice variant of a polynucleotide comprising a sequence of SEQ ID NO:56. Any one of the splice variants described above can encode a polypeptide which contains at least one functional or structural characteristic of PMOD.

10 It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding PMOD, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring PMOD, and all such variations are to be considered as being specifically disclosed.

Although polynucleotides which encode PMOD and its variants are generally capable of hybridizing to polynucleotides encoding naturally occurring PMOD under appropriately selected conditions of stringency, it may be advantageous to produce polynucleotides encoding PMOD or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding PMOD and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of polynucleotides which encode PMOD and PMOD derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic polynucleotide may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a polynucleotide encoding PMOD or any fragment thereof.

Embodiments of the invention can also include polynucleotides that are capable of hybridizing to the claimed polynucleotides, and, in particular, to those having the sequences shown in SEQ ID NO:29-56 and fragments thereof, under various conditions of stringency. (See, e.g., Wahl,

G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Biosciences, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Invitrogen, Carlsbad CA). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Amersham Biosciences), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acids encoding PMOD may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National

Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotides or fragments thereof which encode PMOD may be cloned in recombinant DNA molecules that direct expression of PMOD, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other polynucleotides which encode substantially the same or a functionally equivalent polypeptides may be produced and used to express PMOD.

The polynucleotides of the invention can be engineered using methods generally known in the art in order to alter PMOD-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent No. 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of PMOD, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene

variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, polynucleotides encoding PMOD may be synthesized, in whole or in part, using one or more chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, PMOD itself or a fragment thereof may be synthesized using chemical methods known in the art. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of PMOD, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, *supra*, pp. 28-53.)

In order to express a biologically active PMOD, the polynucleotides encoding PMOD or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotides encoding PMOD. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of polynucleotides encoding PMOD. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where a polynucleotide sequence encoding PMOD and its initiation codon and upstream regulatory sequences

are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing polynucleotides encoding PMOD and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express polynucleotides encoding PMOD. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, *supra*; Ausubel, *supra*; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of polynucleotides to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotides encoding PMOD. For example, routine cloning, subcloning, and propagation of polynucleotides encoding PMOD can be achieved using a

multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPO11 plasmid (Invitrogen). Ligation of polynucleotides encoding PMOD into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for

5 *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of PMOD are needed, e.g. for the production of antibodies, vectors which direct high level expression of PMOD may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

10 Yeast expression systems may be used for production of PMOD. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign polynucleotide sequences into the host genome for stable propagation. (See;

15 e.g., Ausubel, 1995, *supra*; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of PMOD. Transcription of polynucleotides encoding PMOD may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J.

20 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology

25 (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, polynucleotides encoding PMOD may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to

30 obtain infective virus which expresses PMOD in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of

35 DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are

constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression
5 of PMOD in cell lines is preferred. For example, polynucleotides encoding PMOD can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is
10 to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine
15 phosphoribosyltransferase genes, for use in *tk* and *ap^r* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g.,
20 Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate
25 luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the
30 sequence encoding PMOD is inserted within a marker gene sequence, transformed cells containing polynucleotides encoding PMOD can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PMOD under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

35 In general, host cells that contain the polynucleotide encoding PMOD and that express

PMOD may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

5 Immunological methods for detecting and measuring the expression of PMOD using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PMOD is preferred, but a
10 competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

15 A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding PMOD include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, polynucleotides encoding PMOD, or any fragments thereof, may be cloned into a
20 vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Biosciences, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be
25 used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with polynucleotides encoding PMOD may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence
30 and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode PMOD may be designed to contain signal sequences which direct secretion of PMOD through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted polynucleotides or to process the expressed protein in the desired fashion. Such
35 modifications of the polypeptide include, but are not limited to, acetylation, carboxylation,

glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant polynucleotides encoding PMOD may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric PMOD protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of PMOD activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the PMOD encoding sequence and the heterologous protein sequence, so that PMOD may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In another embodiment, synthesis of radiolabeled PMOD may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

PMOD, fragments of PMOD, or variants of PMOD may be used to screen for compounds that specifically bind to PMOD. One or more test compounds may be screened for specific binding to PMOD. In various embodiments, 1, 2, 3, 4, 5, 10, 20, 50, 100, or 200 test compounds can be screened for specific binding to PMOD. Examples of test compounds can include antibodies, anticalins, oligonucleotides, proteins (e.g., ligands or receptors), or small molecules.

In related embodiments, variants of PMOD can be used to screen for binding of test

compounds, such as antibodies, to PMOD, a variant of PMOD, or a combination of PMOD and/or one or more variants PMOD. In an embodiment, a variant of PMOD can be used to screen for compounds that bind to a variant of PMOD, but not to PMOD having the exact sequence of a sequence of SEQ ID NO: 1-28. PMOD variants used to perform such screening can have a range of
5 about 50% to about 99% sequence identity to PMOD, with various embodiments having 60%, 70%, 75%, 80%, 85%, 90%, and 95% sequence identity.

In an embodiment, a compound identified in a screen for specific binding to PMOD can be closely related to the natural ligand of PMOD, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991)
10 Current Protocols in Immunology 1(2):Chapter 5.) In another embodiment, the compound thus identified can be a natural ligand of a receptor PMOD. (See, e.g., Howard, A.D. et al. (2001) Trends Pharmacol. Sci.22:132-140; Wise, A. et al. (2002) Drug Discovery Today 7:235-246.)

In other embodiments, a compound identified in a screen for specific binding to PMOD can be closely related to the natural receptor to which PMOD binds, at least a fragment of the receptor, or
15 a fragment of the receptor including all or a portion of the ligand binding site or binding pocket. For example, the compound may be a receptor for PMOD which is capable of propagating a signal, or a decoy receptor for PMOD which is not capable of propagating a signal (Ashkenazi, A. and V.M. Divit (1999) Curr. Opin. Cell Biol. 11:255-260; Mantovani, A. et al. (2001) Trends Immunol. 22:328-336). The compound can be rationally designed using known techniques. Examples of such
20 techniques include those used to construct the compound etanercept (ENBREL; Immunex Corp., Seattle WA), which is efficacious for treating rheumatoid arthritis in humans. Etanercept is an engineered p75 tumor necrosis factor (TNF) receptor dimer linked to the Fc portion of human IgG₁ (Taylor, P.C. et al. (2001) Curr. Opin. Immunol. 13:611-616).

In one embodiment, two or more antibodies having similar or, alternatively, different
25 specificities can be screened for specific binding to PMOD, fragments of PMOD, or variants of PMOD. The binding specificity of the antibodies thus screened can thereby be selected to identify particular fragments or variants of PMOD. In one embodiment, an antibody can be selected such that its binding specificity allows for preferential identification of specific fragments or variants of PMOD. In another embodiment, an antibody can be selected such that its binding specificity allows
30 for preferential diagnosis of a specific disease or condition having increased, decreased, or otherwise abnormal production of PMOD.

In an embodiment, anticalins can be screened for specific binding to PMOD, fragments of PMOD, or variants of PMOD. Anticalins are ligand-binding proteins that have been constructed based on a lipocalin scaffold (Weiss, G.A. and H.B. Lowman (2000) Chem. Biol. 7:R177-R184;
35 Skerra, A. (2001) J. Biotechnol. 74:257-275). The protein architecture of lipocalins can include a

beta-barrel having eight antiparallel beta-strands, which supports four loops at its open end. These loops form the natural ligand-binding site of the lipocalins, a site which can be re-engineered *in vitro* by amino acid substitutions to impart novel binding specificities. The amino acid substitutions can be made using methods known in the art or described herein, and can include conservative substitutions
5 (e.g., substitutions that do not alter binding specificity) or substitutions that modestly, moderately, or significantly alter binding specificity.

In one embodiment, screening for compounds which specifically bind to, stimulate, or inhibit PMOD involves producing appropriate cells which express PMOD, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells
10 expressing PMOD or cell membrane fractions which contain PMOD are then contacted with a test compound and binding, stimulation, or inhibition of activity of either PMOD or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example,
15 the assay may comprise the steps of combining at least one test compound with PMOD, either in solution or affixed to a solid support, and detecting the binding of PMOD to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a
20 solid support.

An assay can be used to assess the ability of a compound to bind to its natural ligand and/or to inhibit the binding of its natural ligand to its natural receptors. Examples of such assays include radio-labeling assays such as those described in U.S. Patent No. 5,914,236 and U.S. Patent No. 6,372,724. In a related embodiment, one or more amino acid substitutions can be introduced into a
25 polypeptide compound (such as a receptor) to improve or alter its ability to bind to its natural ligands. (See, e.g., Matthews, D.J. and J.A. Wells. (1994) Chem. Biol. 1:25-30.) In another related embodiment, one or more amino acid substitutions can be introduced into a polypeptide compound (such as a ligand) to improve or alter its ability to bind to its natural receptors. (See, e.g., Cunningham, B.C. and J.A. Wells (1991) Proc. Natl. Acad. Sci. USA 88:3407-3411; Lowman, H.B.
30 et al. (1991) J. Biol. Chem. 266:10982-10988.)

PMOD, fragments of PMOD, or variants of PMOD may be used to screen for compounds that modulate the activity of PMOD. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for PMOD activity, wherein PMOD is combined with at least one test compound, and the activity of
35 PMOD in the presence of a test compound is compared with the activity of PMOD in the absence of

the test compound. A change in the activity of PMOD in the presence of the test compound is indicative of a compound that modulates the activity of PMOD. Alternatively, a test compound is combined with an *in vitro* or cell-free system comprising PMOD under conditions suitable for PMOD activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of PMOD may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding PMOD or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (*neo*; Capecchi, M.R. (1989) *Science* 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) *Clin. Invest.* 97:1999-2002; Wagner, K.U. et al. (1997) *Nucleic Acids Res.* 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding PMOD may also be manipulated *in vitro* in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) *Science* 282:1145-1147).

Polynucleotides encoding PMOD can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding PMOD is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress PMOD, e.g., by secreting PMOD in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) *Biotechnol. Annu. Rev.* 4:55-74).

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of PMOD and protein modification and maintenance molecules. In addition, the expression of PMOD is closely associated with epithelial, brain, brain tumor, ileum, lymph node, liver, ovarian, placental, prostate, cerebellum, pituitary gland, small intestine, and testis tissues and promonocyte cells. Further examples of tissues expressing PMOD can be found in Table 6 and can also be found in Example XI. Therefore, PMOD appears to play a role in gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders. In the treatment of disorders associated with increased PMOD expression or activity, it is desirable to decrease the expression or activity of PMOD. In the treatment of disorders associated with decreased PMOD expression or activity, it is desirable to increase the expression or activity of PMOD.

Therefore, in one embodiment, PMOD or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMOD. Examples of such disorders include, but are not limited to, a gastrointestinal disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a cardiovascular disorder, such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular

heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease,

5 congenital heart disease, and complications of cardiac transplantation; an autoimmune/inflammatory disease, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, atherosclerotic plaque rupture, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact

10 dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, degradation of articular cartilage, osteoporosis,

15 pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis,

20 cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid,

25 penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, bone resorption, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary

30 keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, age-related macular degeneration, and sensorineural hearing loss; an epithelial disorder, such as dyshidrotic eczema, allergic contact dermatitis, keratosis pilaris, melasma, vitiligo, actinic keratosis, basal cell carcinoma,

35 squamous cell carcinoma, seborrheic keratosis, folliculitis, herpes simplex, herpes zoster, varicella,

candidiasis, dermatophytosis, scabies, insect bites, cherry angioma, keloid, dermatofibroma, acrochordons, urticaria, transient acantholytic dermatosis, xerosis, eczema, atopic dermatitis, contact dermatitis, hand eczema, nummular eczema, lichen simplex chronicus, asteatotic eczema, stasis dermatitis and stasis ulceration, seborrheic dermatitis, psoriasis, lichen planus, pityriasis rosea, impetigo, ecthyma, dermatophytosis, tinea versicolor, warts, acne vulgaris, acne rosacea, pemphigus vulgaris, pemphigus foliaceus, paraneoplastic pemphigus, bullous pemphigoid, herpes gestationis, dermatitis herpetiformis, linear IgA disease, epidermolysis bullosa acquisita, dermatomyositis, lupus erythematosus, scleroderma and morphea, erythroderma, alopecia, figurate skin lesions, telangiectasias, hypopigmentation, hyperpigmentation, vesicles/bullae, exanthems, cutaneous drug reactions, papulonodular skin lesions, chronic non-healing wounds, photosensitivity diseases, epidermolysis bullosa simplex, epidermolytic hyperkeratosis, epidermolytic and nonepidermolytic palmoplantar keratoderma, ichthyosis bullosa of Siemens, ichthyosis exfoliativa, keratosis palmaris et plantaris, keratosis palmoplantaris, palmoplantar keratoderma, keratosis punctata, Meesmann's corneal dystrophy, pachyonychia congenita, white sponge nevus, steatocystoma multiplex, epidermal nevi/epidermolytic hyperkeratosis type, monilethrix, trichothiodystrophy, chronic hepatitis/cryptogenic cirrhosis, and colorectal hyperplasia; a neurological disorder, such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a reproductive disorder, such as infertility, including tubal disease, ovulatory defects, and endometriosis, a disorder of prolactin production, a disruption of the estrous

cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia.

In another embodiment, a vector capable of expressing PMOD or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMOD including, but not limited to, those described above.

10 In a further embodiment, a composition comprising a substantially purified PMOD in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMOD including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of PMOD may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMOD including, but not limited to, those listed above.

In a further embodiment, an antagonist of PMOD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PMOD. Examples of such disorders include, but are not limited to, those gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders described above. In one aspect, an antibody which specifically binds PMOD may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express PMOD.

25 In an additional embodiment, a vector expressing the complement of the polynucleotide encoding PMOD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PMOD including, but not limited to, those described above.

In other embodiments, any protein, agonist, antagonist, antibody, complementary sequence, or vector embodiments may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

35 An antagonist of PMOD may be produced using methods which are generally known in the art. In particular, purified PMOD may be used to produce antibodies or to screen libraries of

pharmaceutical agents to identify those which specifically bind PMOD. Antibodies to PMOD may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use. Single chain antibodies (e.g., from camels or llamas) may be potent enzyme inhibitors and may have advantages in the design of peptide mimetics, and in the development of immuno-adsorbents and biosensors (Muyldermans, S. (2001) J. Biotechnol. 74:277-302).

For the production of antibodies, various hosts including goats, rabbits, rats, mice, camels, dromedaries, llamas, humans, and others may be immunized by injection with PMOD or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to PMOD have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of PMOD amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to PMOD may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce PMOD-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be

generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for PMOD may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between PMOD and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering PMOD epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for PMOD. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of PMOD-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple PMOD epitopes, represents the average affinity, or avidity, of the antibodies for PMOD. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular PMOD epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the PMOD-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of PMOD, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to

determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of PMOD-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and
5 guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, *supra*, and Coligan et al. *supra*.)

In another embodiment of the invention, polynucleotides encoding PMOD, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA,
10 RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding PMOD. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PMOD. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense
15 sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral
20 vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.*
25 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding PMOD may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined
30 immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal,
35 R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii)

express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as *Candida albicans* and *Paracoccidioides brasiliensis*; and protozoan parasites such as *Plasmodium falciparum* and *Trypanosoma cruzi*). In the case where a genetic deficiency in PMOD expression or regulation causes disease, the expression of PMOD from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

10 In a further embodiment of the invention, diseases or disorders caused by deficiencies in PMOD are treated by constructing mammalian expression vectors encoding PMOD and introducing these vectors by mechanical means into PMOD-deficient cells. Mechanical transfer technologies for use with cells *in vivo* or *ex vitro* include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene
15 transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of PMOD include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX, PCR2-TOPOTA vectors
20 (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). PMOD may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl.*
25 *Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, *supra*), or (iii) a tissue-specific promoter or the native promoter of
30 the endogenous gene encoding PMOD from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method
35 (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al.

(1982) EMBO J. 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to PMOD expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding PMOD under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent No. 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In an embodiment, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding PMOD to cells which have one or more genetic abnormalities with respect to the expression of PMOD. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent No. 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 389:239-242, both incorporated by reference herein.

In another embodiment, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding PMOD to target cells which have one or more genetic abnormalities with

respect to the expression of PMOD. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing PMOD to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has
5 been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent No. 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent No. 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the
10 control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) *J. Virol.* 73:519-532 and Xu, H. et al. (1994) *Dev. Biol.* 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids
15 containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another embodiment, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding PMOD to target cells. The biology of the prototypic alphavirus,
20 Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) *Curr. Opin. Biotechnol.* 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity
25 (e.g., protease and polymerase). Similarly, inserting the coding sequence for PMOD into the alphavirus genome in place of the capsid-coding region results in the production of a large number of PMOD-coding RNAs and the synthesis of high levels of PMOD in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN)
30 indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) *Virology* 228:74-83). The wide host range of alphaviruses will allow the introduction of PMOD into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA
35 transfections, and performing alphavirus infections, are well known to those with ordinary skill in the

art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of RNA molecules encoding PMOD.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA molecules encoding PMOD. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine,

cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding PMOD.

5 Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or
10 promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased PMOD expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding PMOD may be therapeutically useful, and in the treatment of disorders associated with decreased PMOD expression or activity, a compound which specifically promotes expression of the polynucleotide encoding PMOD may be therapeutically useful.

15 At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound
20 based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding PMOD is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding PMOD are assayed
25 by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding PMOD. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide
30 exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys.
35 Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a

combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruice, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruice, T.W. et al. (2000) U.S. Patent No. 6,022,691).

5 Many methods for introducing vectors into cells or tissues are available and equally suitable for use *in vivo*, *in vitro*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.

10 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition
15 which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of PMOD, antibodies to PMOD, and mimetics, agonists, antagonists, or inhibitors of PMOD.

20 The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form.
25 These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton,
30 J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

35 Specialized forms of compositions may be prepared for direct intracellular delivery of

macromolecules comprising PMOD or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, PMOD or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to
5 transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration
10 range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example PMOD or fragments thereof, antibodies of PMOD, and agonists, antagonists or inhibitors of PMOD, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined
15 by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are
20 used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the
25 subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week,
30 or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their
35 inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells,

conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind PMOD may be used for the diagnosis of disorders characterized by expression of PMOD, or in assays to monitor patients being
5 treated with PMOD or agonists, antagonists, or inhibitors of PMOD. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for PMOD include methods which utilize the antibody and a label to detect PMOD in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of
10 reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring PMOD, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PMOD expression. Normal or standard values for PMOD expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to PMOD
15 under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of PMOD expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, polynucleotides encoding PMOD may be used for
20 diagnostic purposes. The polynucleotides which may be used include oligonucleotides, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of PMOD may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of PMOD, and to monitor regulation of PMOD levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotides, including genomic sequences, encoding PMOD or closely related molecules may be used to identify
25 nucleic acid sequences which encode PMOD. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe
30 identifies only naturally occurring sequences encoding PMOD, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the PMOD encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:29-56 or from genomic sequences including promoters, enhancers, and introns of the PMOD gene.

35 Means for producing specific hybridization probes for polynucleotides encoding PMOD

include the cloning of polynucleotides encoding PMOD or PMOD derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotides encoding PMOD may be used for the diagnosis of disorders associated with expression of PMOD. Examples of such disorders include, but are not limited to, a gastrointestinal disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, α_1 -antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a cardiovascular disorder, such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; an autoimmune/inflammatory disease, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, atherosclerotic plaque rupture, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune

polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's

5 thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, degradation of articular cartilage, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and

10 extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including

15 adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, bone resorption, epilepsy, gonadal dysgenesis, WAGR

20 syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, age-related macular

25 degeneration, and sensorineural hearing loss; an epithelial disorder, such as dyshidrotic eczema, allergic contact dermatitis, keratosis pilaris, melasma, vitiligo, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, seborrheic keratosis, folliculitis, herpes simplex, herpes zoster, varicella, candidiasis, dermatophytosis, scabies, insect bites, cherry angioma, keloid, dermatofibroma, acrochordons, urticaria, transient acantholytic dermatosis, xerosis, eczema, atopic dermatitis, contact

30 dermatitis, hand eczema, nummular eczema, lichen simplex chronicus, asteatotic eczema, stasis dermatitis and stasis ulceration, seborrheic dermatitis, psoriasis, lichen planus, pityriasis rosea, impetigo, ecthyma, dermatophytosis, tinea versicolor, warts, acne vulgaris, acne rosacea, pemphigus vulgaris, pemphigus foliaceus, paraneoplastic pemphigus, bullous pemphigoid, herpes gestationis, dermatitis herpetiformis, linear IgA disease, epidermolysis bullosa acquisita, dermatomyositis, lupus

35 erythematosus, scleroderma and morphea, erythroderma, alopecia, figurate skin lesions,

telangiectasias, hypopigmentation, hyperpigmentation, vesicles/bullae, exanthems, cutaneous drug reactions, papulonodular skin lesions, chronic non-healing wounds, photosensitivity diseases, epidermolysis bullosa simplex, epidermolytic hyperkeratosis, epidermolytic and nonepidermolytic palmoplantar keratoderma, ichthyosis bullosa of Siemens, ichthyosis exfoliativa, keratosis palmaris et plantaris, keratosis palmoplantaris, palmoplantar keratoderma, keratosis punctata, Meesmann's

5 corneal dystrophy, pachyonychia congenita, white sponge nevus, steatocystoma multiplex, epidermal nevi/epidermolytic hyperkeratosis type, monilethrix, trichothiodystrophy, chronic hepatitis/cryptogenic cirrhosis, and colorectal hyperplasia; a neurological disorder, such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease,

10 Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases

15 including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve

20 disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia,

25 Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a reproductive disorder, such as infertility, including tubal disease, ovulatory defects, and endometriosis, a disorder of prolactin production, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic

30 pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia. Polynucleotides encoding PMOD may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick,

35 pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to

detect altered PMOD expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, polynucleotides encoding PMOD may be used in assays that detect the presence of associated disorders, particularly those mentioned above. Polynucleotides complementary to sequences encoding PMOD may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of polynucleotides encoding PMOD in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of PMOD, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PMOD, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier, thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding PMOD may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding PMOD, or a fragment of a polynucleotide complementary to the polynucleotide encoding

PMOD, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from polynucleotides encoding PMOD
5 may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from polynucleotides encoding PMOD are used to amplify DNA using the polymerase chain reaction
10 (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA
15 sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the
20 alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

SNPs may be used to study the genetic basis of human disease. For example, at least 16 common SNPs have been associated with non-insulin-dependent diabetes mellitus. SNPs are also useful for examining differences in disease outcomes in monogenic disorders, such as cystic fibrosis,
25 sickle cell anemia, or chronic granulomatous disease. For example, variants in the mannose-binding lectin, MBL2, have been shown to be correlated with deleterious pulmonary outcomes in cystic fibrosis. SNPs also have utility in pharmacogenomics, the identification of genetic variants that influence a patient's response to a drug, such as life-threatening toxicity. For example, a variation in N-acetyl transferase is associated with a high incidence of peripheral neuropathy in response to the
30 anti-tuberculosis drug isoniazid, while a variation in the core promoter of the ALOX5 gene results in diminished clinical response to treatment with an anti-asthma drug that targets the 5-lipoxygenase pathway. Analysis of the distribution of SNPs in different populations is useful for investigating genetic drift, mutation, recombination, and selection, as well as for tracing the origins of populations and their migrations. (Taylor, J.G. et al. (2001) Trends Mol. Med. 7:507-512; Kwok, P.-Y. and Z. Gu
35 (1999) Mol. Med. Today 5:538-543; Nowotny, P. et al. (2001) Curr. Opin. Neurobiol. 11:637-641.)

Methods which may also be used to quantify the expression of PMOD include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be
5 accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotides described herein may be used as elements on a microarray. The microarray can be
10 used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the
15 activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

20 In another embodiment, PMOD, fragments of PMOD, or antibodies specific for PMOD may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of
25 gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of
30 transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines,
35 biopsies, or other biological samples. The transcript image may thus reflect gene expression *in vivo*,

as in the case of a tissue or biopsy sample, or *in vitro*, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with *in vitro* model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In an embodiment, the toxicity of a test compound can be assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another embodiment relates to the use of the polypeptides disclosed herein to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric

focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of interest. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for PMOD to quantify the levels of PMOD expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendozze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the

polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London.

In another embodiment of the invention, nucleic acid sequences encoding PMOD may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent *in situ* hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, *supra*, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding PMOD on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, PMOD, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between PMOD and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with PMOD, or fragments thereof, and washed. Bound PMOD is then detected by methods well known in the art. Purified PMOD can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PMOD specifically compete with a test compound for binding PMOD. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PMOD.

In additional embodiments, the nucleotide sequences which encode PMOD may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder

of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, including U.S. Ser. No. 60/300,508, U.S. Ser. No. 60/303,445, U.S. Ser. No. 60/305,405, U.S. Ser. No. 60/311,442, U.S. Ser. No. 60/314,821, U.S. Ser. No. 60/315,992, and U.S. Ser. No. 60/378/205, are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA). Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Invitrogen), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Invitrogen), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Biosciences) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Invitrogen), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), PCR2-TOPOTA plasmid (Invitrogen), PCMV-ICIS plasmid (Stratagene), pIGEN (Incyte Genomics, Palo Alto CA), pRARE (Incyte Genomics), or pINCY (Incyte Genomics), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli*

cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Invitrogen.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Biosciences or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Amersham Biosciences); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public

databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM; PROTEOME databases with sequences from *Homo sapiens*, *Rattus norvegicus*, *Mus musculus*, *Caenorhabditis elegans*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Candida albicans* (Incyte Genomics, Palo Alto CA); hidden

5 Markov model (HMM)-based protein family databases such as PFAM, INCY, and TIGRFAM (Haft, D.H. et al. (2001) Nucleic Acids Res. 29:41-43); and HMM-based protein domain databases such as SMART (Schultz et al. (1998) Proc. Natl. Acad. Sci. USA 95:5857-5864; Letunic, I. et al. (2002) Nucleic Acids Res. 30:242-244). (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) Curr. Opin. Struct. Biol.

10 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and

15 Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein

20 databases (genpept), SwissProt, the PROTEOME databases, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, hidden Markov model (HMM)-based protein family databases such as PFAM, INCY, and TIGRFAM; and HMM-based protein domain databases such as SMART. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide

25 sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and

30 threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value,

35 the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:29-56. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 2.

5 **IV. Identification and Editing of Coding Sequences from Genomic DNA**

Putative protein modification and maintenance molecules were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, 10 and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode protein modification and maintenance molecules, 15 the encoded polypeptides were analyzed by querying against PFAM models for protein modification and maintenance molecules. Potential protein modification and maintenance molecules were also identified by homology to Incyte cDNA sequences that had been annotated as protein modification and maintenance molecules. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted 20 sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were 25 obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

30 **"Stitched" Sequences**

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm 35 based on graph theory and dynamic programming to integrate cDNA and genomic information,

generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpri public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

15 **"Stretched" Sequences**

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

25 **VI. Chromosomal Mapping of PMOD Encoding Polynucleotides**

The sequences which were used to assemble SEQ ID NO:29-56 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:29-56 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences

had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Génethon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the

entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

5 Alternatively, polynucleotides encoding PMOD are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive
10 system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following
15 disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding PMOD. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

20 **VIII. Extension of PMOD Encoding Polynucleotides**

Full length polynucleotides are produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06
25 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one
30 extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Biosciences), ELONGASE enzyme
35 (Invitrogen), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair

PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; 5 Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 µl PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 µl of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II 10 (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 µl to 10 µl aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, 15 digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Biosciences). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham 20 Biosciences), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase 25 (Amersham Biosciences) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% 30 dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Biosciences) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotides are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for 35 such extension, and an appropriate genomic library.

IX. Identification of Single Nucleotide Polymorphisms in PMOD Encoding Polynucleotides

Common DNA sequence variants known as single nucleotide polymorphisms (SNPs) were identified in SEQ ID NO:29-56 using the LIFESEQ database (Incyte Genomics). Sequences from the same gene were clustered together and assembled as described in Example III, allowing the
5 identification of all sequence variants in the gene. An algorithm consisting of a series of filters was used to distinguish SNPs from other sequence variants. Preliminary filters removed the majority of basecall errors by requiring a minimum Phred quality score of 15, and removed sequence alignment errors and errors resulting from improper trimming of vector sequences, chimeras, and splice variants. An automated procedure of advanced chromosome analysis analysed the original
10 chromatogram files in the vicinity of the putative SNP. Clone error filters used statistically generated algorithms to identify errors introduced during laboratory processing, such as those caused by reverse transcriptase, polymerase, or somatic mutation. Clustering error filters used statistically generated algorithms to identify errors resulting from clustering of close homologs or pseudogenes, or due to contamination by non-human sequences. A final set of filters removed duplicates and SNPs found in
15 immunoglobulins or T-cell receptors.

Certain SNPs were selected for further characterization by mass spectrometry using the high throughput MASSARRAY system (Sequenom, Inc.) to analyze allele frequencies at the SNP sites in four different human populations. The Caucasian population comprised 92 individuals (46 male, 46 female), including 83 from Utah, four French, three Venezuelan, and two Amish individuals. The
20 African population comprised 194 individuals (97 male, 97 female), all African Americans. The Hispanic population comprised 324 individuals (162 male, 162 female), all Mexican Hispanic. The Asian population comprised 126 individuals (64 male, 62 female) with a reported parental breakdown of 43% Chinese, 31% Japanese, 13% Korean, 5% Vietnamese, and 8% other Asian. Allele frequencies were first analyzed in the Caucasian population; in some cases those SNPs which showed
25 no allelic variance in this population were not further tested in the other three populations.

X. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:29-56 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide
30 fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Biosciences), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Biosciences). An aliquot containing 10^7
35 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of

human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16
5 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

XI. Microarrays

10 The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, *supra.*), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999), *supra*). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers.
15 Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Skena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645;
20 Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the
25 biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element
30 on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and
35 poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is

reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Biosciences). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with

5 GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by *in vitro* transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc.

10 (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

15 Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Biosciences).

20 Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a

25 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in U.S. Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

30 Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

35

Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC

computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping
5 emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

10 Array elements that exhibited at least about a two-fold change in expression, a signal-to-background ratio of at least 2.5, and an element spot size of at least 40% were identified as differentially expressed using the GEMTOOLS program (Incyte Genomics).

Expression

The human C3A cell line is a clonal derivative of HepG2/C3 (hepatoma cell line, isolated
15 from a 15-year-old male with liver tumor), which was selected for strong contact inhibition of growth. The use of a clonal population enhances the reproducibility of the cells. C3A cells have many characteristics of primary human hepatocytes in culture: i) expression of insulin receptor and insulin-like growth factor II receptor; ii) secretion of a high ratio of serum albumin compared with α -fetoprotein; iii) conversion of ammonia to urea and glutamine; iv) metabolism of aromatic amino
20 acids; and v) proliferation in glucose-free and insulin-free medium. The C3A cell line is now well established as an in vitro model of the mature human liver (Mickelson et al. (1995) Hepatology 22:866-875; Nagendra et al. (1997) Am. J. Physiol. 272:G408-G416). The expression of SEQ ID NO:29 was altered by a factor of 2 or more in cells treated with a variety of steroids including prednisone, dexamethasone, medroxyprogesterone, budesonide, and beclomthasone. In addition, the
25 expression of SEQ ID NO:29 was was altered by a factor of two or more in C3A cells. Therefore, SEQ ID NO:29 can be used in assays related to treatment for cell proliferative disorders.

For example, SEQ ID NO:31 and SEQ ID NO:34 showed differential expression in breast tumor cell lines versus normal breast epithelial cells as determined by microarray analysis. The expression of SEQ ID NO:31 and SEQ ID NO:34 was decreased by at least two fold in breast tumor
30 cell lines that were harvested from donors with both early and late stages of tumor progression and malignant transformation. Therefore, SEQ ID NO:31 and SEQ ID NO:34 can be used in diagnostic assays for breast cancer.

In another example, SEQ ID NO:33 showed differential expression in response to several compounds which produce known metabolic and toxicological responses. The expression of SEQ ID
35 NO:33 was reduced by at least two fold in the human C3A liver cell line incubated for varying

lengths of time with compounds including fenofibrate, clofibrate, dexamethasone, beclomethasone, medroxyprogesterone, budesonide, and betamethasone. Therefore, SEQ ID NO:33 can be used in toxicology testing.

5 SEQ ID NO:35 showed differential expression in human peripheral blood mononuclear cells (PBMCs) following exposure to 5 and 25 μ M prednisone for 24 hours. Prednisone is a corticosteroid that is metabolized in the liver to its active form, prednisolone. Prednisone is approximately four times more potent as a glucocorticoid than hydrocortisone. Glucocorticoids are naturally occurring hormones that prevent or suppress inflammation and immune responses when administered at pharmacologic doses. At the molecular level, unbound glucocorticoids readily cross cell membranes and bind with high affinity to specific cytoplasmic receptors. Subsequent to binding, transcription and, ultimately, protein synthesis are affected. The result can include inhibition of leukocyte infiltration at the site of inflammation, interference in the function of mediators of the inflammatory response, and suppression of humoral immune responses. PBMCs can be classified into discrete cellular populations representing the major cellular components of the immune system. PBMCs 15 contain about 52% lymphocytes (12% B lymphocytes, 40% T lymphocytes {25% CD4+ and 15% CD8+}), 20% NK cells, 25% monocytes, and 3% various cells that include dendritic cells and progenitor cells. These cells were pooled from the blood of 6 healthy volunteer donors. The expression of SEQ ID NO:35 was decreased by at least two-fold in prednisone-treated (5 and 25 μ M) cells as compared to untreated controls.

20 SEQ ID NO:44 showed differential expression in normal tissue versus tissue affected by prostate carcinoma by microarray analysis. Expression of SEQ ID NO:44 in a primary prostate epithelial cell line (PrEC) isolated from a normal donor was compared to expression of SEQ ID NO:44 in a prostate carcinoma cell line isolated (LNCaP) from a lymph node biopsy of a 50-year-old male with metastatic prostate carcinoma. Expression of SEQ ID NO:44 was decreased by at least 25 two-fold in the cell line affected by prostate carcinoma. In addition, expression of SEQ ID NO:44 in peripheral blood mononuclear cells isolated from a pool of healthy donors was decreased by at least two-fold by treatment with 1 ng/ml IL12 for 24 hours. The expression of SEQ ID NO:44 was also shown to be differentially expressed in a comparison of breast cells lines by microarray analysis. In four of the seven breast cancer cell lines tested, expression of SEQ ID NO:44 was shown to be 30 decreased by at least two-fold when compared to normal mammary epithelial cells (HMEC), indicating the use of SEQ ID NO:44 as a diagnostic marker, for disease staging, and as a therapeutic target for protease-associated diseases including prostate and breast cancer.

For example, SEQ ID NO:51 showed differential expression in toxicology studies as determined by microarray analysis. The expression of SEQ ID NO:51 was decreased by at least two 35 fold in a human C3A liver cell line treated with various drugs (e.g., steroids, steroid hormones)

relative to untreated C3A cells. The human C3A cell line is a clonal derivative of HepG2/C3 (hepatoma cell line, isolated from a 15-year-old male with liver tumor), which was selected for strong contact inhibition of growth. The C3A cell line is well established as an *in vitro* model of the mature human liver (Mickelson et al. (1995) Hepatology 22:866-875; Nagendra et al. (1997) Am J Physiol 5 272:G408-G416). Effects upon liver metabolism are important to understanding the pharmacodynamics of a drug. Therefore, SEQ ID NO:51 can be used for understanding the pharmacodynamics of a drug.

In another example, SEQ ID NO:55 showed differential expression in lung adenocarcinoma versus normal lung tissues as determined by microarray analysis. The expression of SEQ ID NO:55 10 was decreased by at least two fold in lung adenocarcinoma relative to grossly uninvolved normal lung tissue from the same donor. Therefore, SEQ ID NO:55 can be used as a diagnostic marker for disease staging or as a potential therapeutic target for lung cancer.

As another example, SEQ ID NO:56 is downregulated in breast cancer cell lines versus nonmalignant mammary epithelial cells, as determined by microarray analysis. In one experiment, 15 gene expression profiles of nonmalignant mammary epithelial cells were compared to gene expression profiles of various breast carcinoma lines at different stages of tumor progression. The cells were grown in defined serum-free H14 medium to 70-80% confluence prior to RNA harvest. Cell lines compared included: a) HMEC, a primary breast epithelial cell line isolated from a normal donor, b)MCF-10A, a breast mammary gland cell line isolated from a 36-year-old woman with 20 fibrocystic breast disease, c)MCF7, a nonmalignant breast adenocarcinoma cell line isolated from the pleural effusion of a 69- year-old female, d)T-47D, a breast carcinoma cell line isolated from a pleural effusion obtained from a 54-year-old female with an infiltrating ductal carcinoma of the breast, e)Sk-BR-3, a breast adenocarcinoma cell line isolated from a malignant pleural effusion of a 43-year-old female, f)BT-20, a breast carcinoma cell line derived *in vitro* from cells emigrating out of 25 thin slices of the tumor mass isolated from a 74-year-old female, g)MDA-mb-231, a breast tumor cell line isolated from the pleural effusion of a 51-year-old female, and h)MDA-mb-435S, a spindle-shaped strain that evolved from the parent line (435) isolated by R. Cailleau from pleural effusion of a 31-year-old female with metastatic, ductal adenocarcinoma of the breast. Therefore, SEQ ID NO:56 can be used in monitoring treatment of, and diagnostic assays for, breast cancer.

30 As another example, SEQ ID NO:56 is downregulated in prostate carcinomas versus primary prostate epithelial cells, as determined by microarray analysis. Primary prostate epithelial cells were compared with prostate carcinomas representative of the different stages of tumor progression. Cell lines compared included: a) PrEC, a primary prostate epithelial cell line isolated from a normal donor, b) DU 145, a prostate carcinoma cell line isolated from a metastatic site in the brain of 69-year old 35 male with widespread metastatic prostate carcinoma, c) LNCaP, a prostate carcinoma cell line

isolated from a lymph node biopsy of a 50-year-old male with metastatic prostate carcinoma, and d)PC-3, a prostate adenocarcinoma cell line isolated from a metastatic site in the bone of a 62-year-old male with grade IV prostate adenocarcinoma. In one experiment, cells were grown in basal media in the absence of growth factors and hormones. In a second experiment, cells were grown under optimal growth conditions, in the presence of growth factors and nutrients. Cells grown under restrictive conditions were compared to normal PrECs grown under restrictive conditions. Therefore, SEQ ID NO:56 can be used in monitoring treatment of, and diagnostic assays for, prostate cancer.

XII. Complementary Polynucleotides

Sequences complementary to the PMOD-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring PMOD. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of PMOD. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the PMOD-encoding transcript.

XIII. Expression of PMOD

Expression and purification of PMOD is achieved using bacterial or virus-based expression systems. For expression of PMOD in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express PMOD upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of PMOD in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding PMOD by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, PMOD is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Biosciences). Following purification, the GST moiety can be proteolytically cleaved from PMOD at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10 and 16). Purified PMOD obtained by these methods can be used directly in the assays shown in Examples XVII, XVIII, and XIX, where applicable.

XIV. Functional Assays

PMOD function is assessed by expressing the sequences encoding PMOD at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT plasmid (Invitrogen, Carlsbad CA) and PCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of PMOD on gene expression can be assessed using highly purified

populations of cells transfected with sequences encoding PMOD and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake
5 Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding PMOD and other genes of interest can be analyzed by northern analysis or microarray techniques.

XV. Production of PMOD Specific Antibodies

PMOD substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g.,
10 Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize animals (e.g., rabbits, mice, etc.) and to produce antibodies using standard protocols.

Alternatively, the PMOD amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for
15 selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-
Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to
20 increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-PMOD activity by, for example, binding the peptide or PMOD to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

25 XVI. Purification of Naturally Occurring PMOD Using Specific Antibodies

Naturally occurring or recombinant PMOD is substantially purified by immunoaffinity chromatography using antibodies specific for PMOD. An immunoaffinity column is constructed by covalently coupling anti-PMOD antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Biosciences). After the coupling, the resin is blocked and
30 washed according to the manufacturer's instructions.

Media containing PMOD are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PMOD (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/PMOD binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such
35 as urea or thiocyanate ion), and PMOD is collected.

XVII. Identification of Molecules Which Interact with PMOD

PMOD, or biologically active fragments thereof, are labeled with ^{125}I Bolton-Hunter reagent. (See, e.g., Bolton, A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled PMOD, washed, and any wells with labeled PMOD complex are assayed. Data obtained using different concentrations of PMOD are used to calculate values for the number, affinity, and association of PMOD with the candidate molecules.

Alternatively, molecules interacting with PMOD are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989) *Nature* 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

PMOD may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

XVII. Demonstration of PMOD Activity

PMOD activity can be demonstrated using a generic immunoblotting strategy or through a variety of specific activity assays, some of which are outlined below. As a general approach, cell lines or tissues transformed with a vector containing PMOD coding sequences can be assayed for PMOD activity by immunoblotting. Transformed cells are denatured in SDS in the presence of β -mercaptoethanol, nucleic acids are removed by ethanol precipitation, and proteins are purified by acetone precipitation. Pellets are resuspended in 20 mM Tris buffer at pH 7.5 and incubated with Protein G-Sepharose pre-coated with an antibody specific for PMOD. After washing, the Sepharose beads are boiled in electrophoresis sample buffer, and the eluted proteins subjected to SDS-PAGE. The SDS-PAGE is transferred to a membrane for immunoblotting, and the PMOD activity is assessed by visualizing and quantifying bands on the blot using the antibody specific for PMOD as the primary antibody and ^{125}I -labeled IgG specific for the primary antibody as the secondary antibody.

PMOD kinase activity is measured by quantifying the phosphorylation of a protein substrate by PMOD in the presence of gamma-labeled ^{32}P -ATP. PMOD is incubated with the protein substrate, ^{32}P -ATP, and an appropriate kinase buffer. The ^{32}P incorporated into the substrate is separated from free ^{32}P -ATP by electrophoresis and the incorporated ^{32}P is counted using a radioisotope counter. The amount of incorporated ^{32}P is proportional to the activity of PMOD. A determination of the specific amino acid residue phosphorylated is made by phosphoamino acid analysis of the hydrolyzed protein.

PMOD phosphatase activity is measured by the hydrolysis of P-nitrophenyl phosphate (PNPP). PMOD is incubated together with PNPP in HEPES buffer pH 7.5, in the presence of 0.1% β -mercaptoethanol at 37°C for 60 min. The reaction is stopped by the addition of 6 ml of 10 N NaOH

and the increase in light absorbance at 410 nm resulting from the hydrolysis of PNPP is measured using a spectrophotometer. The increase in light absorbance is proportional to the activity of PMOD in the assay (Diamond, R.H. et al. (1994) Mol. Cell. Biol. 14:3752-62).

5 The assay for SEQ ID NO:10 is carried out as described above for PMOD using a cysteine protease, such as papain, assayed in the absence and in the presence of various concentrations of SEQ ID NO:10. Inhibition of papain protease activity is proportional to the activity of SEQ ID NO:10 in the assay.

The assay for SEQ ID NO:11 is carried out as described above for PMOD using matrix metalloproteinase assayed in the absence and in the presence of various concentrations of SEQ ID
10 NO:11. Inhibition of matrix metalloproteinase activity is proportional to the activity of SEQ ID NO:11 in the assay.

In the alternative, PMOD phosphatase activity is determined by measuring the amount of phosphate removed from a phosphorylated protein substrate. Reactions are performed with 2 or 4 nM enzyme in a final volume of 30 μ l containing 60 mM Tris, pH 7.6, 1 mM EDTA, 1 mM EGTA, 0.1%
15 2-mercaptoethanol and 10 μ M substrate, 32 P-labeled on serine/threonine or tyrosine, as appropriate. Reactions are initiated with substrate and incubated at 30° C for 10-15 min. Reactions are quenched with 450 μ l of 4% (w/v) activated charcoal in 0.6 M HCl, 90 mM $\text{Na}_4\text{P}_2\text{O}_7$, and 2 mM NaH_2PO_4 , then centrifuged at 12,000 \times g for 5 min. Acid-soluble ^{32}Pi is quantified by liquid scintillation counting (Sinclair, C. et al. (1999) J. Biol. Chem. 274:23666-23672).

20 PMOD protease activity is measured by the hydrolysis of appropriate synthetic peptide substrates conjugated with various chromogenic molecules in which the degree of hydrolysis is quantified by spectrophotometric (or fluorometric) absorption of the released chromophore (Beynon, R.J. and J.S. Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University Press, New York, NY, pp.25-55). Peptide substrates are designed according to the category of protease activity
25 as endopeptidase (serine, cysteine, aspartic proteases, or metalloproteases), aminopeptidase (leucine aminopeptidase), or carboxypeptidase (carboxypeptidases A and B, procollagen C-proteinase). Commonly used chromogens are 2-naphthylamine, 4-nitroaniline, and furylacrylic acid. Assays are performed at ambient temperature and contain an aliquot of the enzyme and the appropriate substrate in a suitable buffer. Reactions are carried out in an optical cuvette, and the increase/decrease in
30 absorbance of the chromogen released during hydrolysis of the peptide substrate is measured. The change in absorbance is proportional to the enzyme activity in the assay.

In the alternative, an assay for PMOD protease activity takes advantage of fluorescence resonance energy transfer (FRET) that occurs when one donor and one acceptor fluorophore with an appropriate spectral overlap are in close proximity. A flexible peptide linker containing a cleavage
35 site specific for PMOD is fused between a red-shifted variant (RSGFP4) and a blue variant (BFP5) of

Green Fluorescent Protein. This fusion protein has spectral properties that suggest energy transfer is occurring from BFP5 to RSGFP4. When the fusion protein is incubated with PMOD, the substrate is cleaved, and the two fluorescent proteins dissociate. This is accompanied by a marked decrease in energy transfer which is quantified by comparing the emission spectra before and after the addition of PMOD (Mitra, R.D. et al (1996) *Gene* 173:13-17). This assay can also be performed in living cells. In this case the fluorescent substrate protein is expressed constitutively in cells and PMOD is introduced on an inducible vector so that FRET can be monitored in the presence and absence of PMOD (Sagot, I. et al (1999) *FEBS Letters* 447:53-57).

- An assay for ubiquitin hydrolase activity measures the hydrolysis of a ubiquitin precursor.
- 10 The assay is performed at ambient temperature and contains an aliquot of PMOD and the appropriate substrate in a suitable buffer. Chemically synthesized human ubiquitin-valine may be used as substrate. Cleavage of the C-terminal valine residue from the substrate is monitored by capillary electrophoresis (Franklin, K. et al. (1997) *Anal. Biochem.* 247:305-309).

- PMOD protease inhibitor activity for alpha 2-HS-glycoprotein (AHSG) can be measured as a decrease in osteogenic activity in dexamethasone-treated rat bone marrow cell cultures (dex-RBMC). Assays are carried out in 96-well culture plates containing minimal essential medium supplemented with 15% fetal bovine serum, ascorbic acid (50 µg/ml), antibiotics (100 µg/ml penicillin G, 50 µg/ml gentamicin, 0.3 µg/ml fungizone), 10 mM B-glycerophosphate, dexamethasone (10^{-8} M) and various concentrations of PMOD for 12-14 days. Mineralized tissue formation in the cultures is quantified by measuring the absorbance at 525 nm using a 96-well plate reader (Binkert, C. et al. *supra*).
- 20

- PMOD protease inhibitor activity for inter-alpha-trypsin inhibitor (ITI) can be measured by a continuous spectrophotometric rate determination of trypsin activity. The assay is performed at ambient temperature in a quartz cuvette in pH 7.6 assay buffer containing 63 mM sodium phosphate, 0.23 mM N α -benzoyl-L-arginine ethyl ester, 0.06 mM hydrochloric acid, 100 units trypsin, and various concentrations of PMOD. Immediately after mixing by inversion, the increase in $A_{253\text{ nm}}$ is recorded for approximately 5 minutes and the enzyme activity is calculated (Bergmeyer, H.U. et al. (1974) *Meth. Enzym. Anal.* 1:515-516)
- 25

- PMOD isomerase activity such as peptidyl prolyl *cis/trans* isomerase activity can be assayed by an enzyme assay described by Rahfeld, J.U., et al. (1994) (*FEBS Lett.* 352: 180-184). The assay is performed at 10 °C in 35 mM HEPES buffer, pH 7.8, containing chymotrypsin (0.5 mg/ml) and PMOD at a variety of concentrations. Under these assay conditions, the substrate, Suc-Ala-Xaa-Pro-Phe-4-NA, is in equilibrium with respect to the prolyl bond, with 80-95% in *trans* and 5-20% in *cis* conformation. An aliquot (2 µl) of the substrate dissolved in dimethyl sulfoxide (10 mg/ml) is added to the reaction mixture described above. Only the *cis* isomer of the substrate is a substrate for cleavage by chymotrypsin. Thus, as the substrate is isomerized by PMOD, the product is cleaved by
- 30
- 35

chymotrypsin to produce 4-nitroanilide, which is detected by its absorbance at 390 nm. 4-Nitroanilide appears in a time-dependent and a PMOD concentration-dependent manner.

PMOD galactosyltransferase activity can be determined by measuring the transfer of radiolabeled galactose from UDP-galactose to a GlcNAc-terminated oligosaccharide chain (Kolbinger, F. et al. (1998) J. Biol. Chem. 273:58-65). The sample is incubated with 14 μ l of assay stock solution (180 mM sodium cacodylate, pH 6.5, 1 mg/ml bovine serum albumin, 0.26 mM UDP-galactose, 2 μ l of UDP-[3 H]galactose), 1 μ l of MnCl_2 (500 mM), and 2.5 μ l of $\text{GlcNAc}\beta\text{O}-(\text{CH}_2)_8\text{-CO}_2\text{Me}$ (37 mg/ml in dimethyl sulfoxide) for 60 minutes at 37 °C. The reaction is quenched by the addition of 1 ml of water and loaded on a C18 Sep-Pak cartridge (Waters), and the column is washed twice with 5 ml of water to remove unreacted UDP-[3 H]galactose. The [3 H]galactosylated $\text{GlcNAc}\beta\text{O}-(\text{CH}_2)_8\text{-CO}_2\text{Me}$ remains bound to the column during the water washes and is eluted with 5 ml of methanol. Radioactivity in the eluted material is measured by liquid scintillation counting and is proportional to galactosyltransferase activity in the starting sample.

PMOD induction by heat or toxins may be demonstrated using primary cultures of human fibroblasts or human cell lines such as CCL-13, HEK293, or HEP G2 (ATCC). To heat induce PMOD expression, aliquots of cells are incubated at 42 °C for 15, 30, or 60 minutes. Control aliquots are incubated at 37 °C for the same time periods. To induce PMOD expression by toxins, aliquots of cells are treated with 100 μ M arsenite or 20 mM azetidine-2-carboxylic acid for 0, 3, 6, or 12 hours. After exposure to heat, arsenite, or the amino acid analogue, samples of the treated cells are harvested and cell lysates prepared for analysis by western blot. Cells are lysed in lysis buffer containing 1% Nonidet P-40, 0.15 M NaCl, 50 mM Tris-HCl, 5 mM EDTA, 2 mM N-ethylmaleimide, 2 mM phenylmethylsulfonyl fluoride, 1 mg/ml leupeptin, and 1 mg/ml pepstatin. Twenty micrograms of the cell lysate is separated on an 8% SDS-PAGE gel and transferred to a membrane. After blocking with 5% nonfat dry milk/phosphate-buffered saline for 1 h, the membrane is incubated overnight at 4°C or at room temperature for 2-4 hours with an appropriate dilution of anti-PMOD serum in 2% nonfat dry milk/phosphate-buffered saline. The membrane is then washed and incubated with a 1:1000 dilution of horseradish peroxidase-conjugated goat anti-rabbit IgG in 2% dry milk/phosphate-buffered saline. After washing with 0.1% Tween 20 in phosphate-buffered saline, the PMOD protein is detected and compared to controls using chemiluminescence.

PMOD lysyl hydroxylase activity is determined by measuring the production of hydroxy[14 C]lysine from [14 C]lysine. Radiolabeled procollagen is incubated with PMOD in buffer containing ascorbic acid, iron sulfate, dithiothreitol, bovine serum albumin, and catalase. Following a 30 minute incubation, the reaction is stopped by the addition of acetone, and centrifuged. The sedimented material is dried, and the hydroxy[14 C]lysine is converted to [14 C]formaldehyde by oxidation with periodate, and then extracted into toluene. The amount of 14 C extracted into toluene is

quantified by scintillation counting, and is proportional to the activity of PMOD in the sample (Kivirikko, K., and Myllyla, R. (1982) *Methods Enzymol.* 82:245-304).

XVIII. Identification of PMOD Substrates

Phage display libraries can be used to identify optimal substrate sequences for PMOD. A random hexamer followed by a linker and a known antibody epitope is cloned as an N-terminal extension of gene III in a filamentous phage library. Gene III codes for a coat protein, and the epitope will be displayed on the surface of each phage particle. The library is incubated with PMOD under proteolytic conditions so that the epitope will be removed if the hexamer codes for a PMOD cleavage site. An antibody that recognizes the epitope is added along with immobilized protein A. Uncleaved phage, which still bear the epitope, are removed by centrifugation. Phage in the supernatant are then amplified and undergo several more rounds of screening. Individual phage clones are then isolated and sequenced. Reaction kinetics for these peptide substrates can be studied using an assay in Example XVII, and an optimal cleavage sequence can be derived (Ke, S.H. et al. (1997) *J. Biol. Chem.* 272:16603-16609).

To screen for in vivo PMOD substrates, this method can be expanded to screen a cDNA expression library displayed on the surface of phage particles (T7SELECT™10-3 Phage display vector, Novagen, Madison, WI) or yeast cells (pYD1 yeast display vector kit, Invitrogen, Carlsbad, CA). In this case, entire cDNAs are fused between Gene III and the appropriate epitope.

XIX. Identification of PMOD Inhibitors

Compounds to be tested are arrayed in the wells of a multi-well plate in varying concentrations along with an appropriate buffer and substrate, as described in the assays in Example XVII. PMOD activity is measured for each well and the ability of each compound to inhibit PMOD activity can be determined, as well as the dose-response kinetics. This assay could also be used to identify molecules which enhance PMOD activity.

In the alternative, phage display libraries can be used to screen for peptide PMOD inhibitors. Candidates are found among peptides which bind tightly to a protease. In this case, multi-well plate wells are coated with PMOD and incubated with a random peptide phage display library or a cyclic peptide library (Koivunen, E. et al. (1999) *Nature Biotech* 17:768-774). Unbound phage are washed away and selected phage amplified and rescreened for several more rounds. Candidates are tested for PMOD inhibitory activity using an assay described in Example XVII.

Various modifications and variations of the described compositions, methods, and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. It will be appreciated that the invention provides novel and useful proteins, and their encoding polynucleotides, which can be used in the drug discovery process, as well as methods for

using these compositions for the detection, diagnosis, and treatment of diseases and conditions.

Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments.

Nor should the description of such embodiments be considered exhaustive or limit the invention to
5 the precise forms disclosed. Furthermore, elements from one embodiment can be readily recombined
with elements from one or more other embodiments. Such combinations can form a number of
embodiments within the scope of the invention. It is intended that the scope of the invention be
defined by the following claims and their equivalents.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Incyte Full Length Clones
7994355	1	7994355CD1	29	7994355CB1	
7475875	2	7475875CD1	30	7475875CB1	
71231882	3	71231882CD1	31	71231882CB1	
2875922	4	2875922CD1	32	2875922CB1	90129891CA2
8158136	5	8158136CD1	33	8158136CB1	90038744CA2
5969491	6	5969491CD1	34	5969491CB1	
7497367	7	7497367CD1	35	7497367CB1	90092215CA2
7632424	8	7632424CD1	36	7632424CB1	
1804436	9	1804436CD1	37	1804436CB1	
7486358	10	7486358CD1	38	7486358CB1	
7472344	11	7472344CD1	39	7472344CB1	
7192959	12	7192959CD1	40	7192959CB1	
6169565	13	6169565CD1	41	6169565CB1	6169565CA2
7494717	14	7494717CD1	42	7494717CB1	
7497510	15	7497510CD1	43	7497510CB1	90166158CA2, 90166182CA2, 90166190CA2, 90166282CA2, 90166290CA2, 90189432CA2, 90189440CA2, 90189464CA2, 90189480CA2, 90189516CA2, 90189532CA2, 90189548CA2, 90189596CA2, 90189833CA2, 90189849CA2, 90189857CA2, 90189865CA2, 90189881CA2, 90189933CA2, 90189941CA2, 90189949CA2, 90189957CA2, 90189981CA2, 90189989CA2, 90190189CA2
7498882	16	7498882CD1	44	7498882CB1	
5524205	17	5524205CD1	45	5524205CB1	
7102342	18	7102342CD1	46	7102342CB1	
4169939	19	4169939CD1	47	4169939CB1	

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Incyte Full Length Clones
6539977	20	6539977CD1	48	6539977CB1	90188738CA2, 90188893CA2, 95003737CA2, 95003761CA2, 95003853CA2, 95003869CA2, 95003905CA2, 95003913CA2, 95003969CA2, 95004005CA2, 95004061CA2, 95004069CA2
7675588	21	7675588CD1	49	7675588CB1	4213559CA2, 90166613CA2
6244077	22	6244077CD1	50	6244077CB1	90110106CA2, 90110114CA2, 90110154CA2, 90110162CA2, 90110170CA2, 90110186CA2, 90110194CA2, 90110270CA2, 90110278CA2, 90110286CA2, 90110294CA2, 90110470CA2
7498404	23	7498404CD1	51	7498404CB1	
7391748	24	7391748CD1	52	7391748CB1	
7499780	25	7499780CD1	53	7499780CB1	
7499881	26	7499881CD1	54	7499881CB1	
7488579	27	7488579CD1	55	7488579CB1	
7510521	28	7510521CD1	56	7510521CB1	

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
1	7994355CD1	g7108521	1.3E-85	[Arabidopsis thaliana] Ubiquitin-protein ligase 2 Bates, P.W. and Vierstra, R.D. (1999) Plant J. 20 (2), 183-195 UPL1 and 2, two 405 kDa ubiquitin-protein ligases from Arabidopsis thaliana related to the HECT-domain protein family
2	7475875CD1	g4262617	1.8E-42	[Caenorhabditis elegans] contains similarity to dual specificity phosphatase, catalytic domain (Pfam:PF00782)
3	71231882CD1	g1545952	1.7E-36	[Homo sapiens] herpesvirus associated ubiquitin-specific protease (HAUSP) Everett, R.D., et al. (1997) A novel ubiquitin-specific protease is dynamically associated with the PML nuclear domain and binds to a herpesvirus regulatory protein. EMBO J. 16, 566-577
4	2875922CD1	g14582773	0.0	[Homo sapiens] sumo/sentrin-specific protease
5	8158136CD1	g11993492	1.1E-79	[Arabidopsis thaliana] ubiquitin-specific protease 26 Yan, N., et al. (2000) The Ubiquitin-Specific Protease Family from Arabidopsis. AtUBP1 and 2 Are Required for the Resistance to the Amino Acid Analog Canavanine. Plant Physiol. 124, 1828-1843
6	5969491CD1	g1545952	1.6E-36	[Arabidopsis thaliana] ubiquitin carboxyl-terminal hydrolase-like protein Everett et al., supra
7	7497367CD1	g179936	1.7E-235	Homo sapiens] carboxypeptidase N (EC 3.4.17.3) Tan, F. et al. (1990) The deduced protein sequence of the human carboxypeptidase N high molecular weight subunit reveals the presence of leucine-rich tandem repeats J. Biol. Chem. 265:13-19
8	7632424CD1	g11071727	5.1E-216	[Homo sapiens] putative sialoglycoprotease type 2
10	7486358CD1	g3928491	1.4E-27	[Mus musculus] testatin Tohonen, V. et al. (1998) Proc. Natl. Acad. Sci. U.S.A. 95 (24), 14208-14213
11	7472344CD1	g15559064	0.0	[Mus musculus] SNAG1
12	7192959CD1	g6456116	9.5E-70	[Mus musculus] F-box protein FBX17
13	6169565CD1	g804764	1.3E-10	[Homo sapiens] neutral protease large subunit Johansen, T. et al. (1989) Members of the RTVL-H family of human endogenous retrovirus-like elements are expressed in placenta Gene 79:259-267

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
14	7494717CD1	g8489831	4.8E-31	[Homo sapiens] ubiquitin-conjugating BIR-domain enzyme APOLLON Chen, Z. et al. (1999) A human IAP-family gene, apollon, expressed in human brain cancer cells Biochem. Biophys. Res. Commun. 264:847-854
15	7497510CD1	g450529	6.3E-34	[Homo sapiens] Kunitz-type protease inhibitor, HKIB9
16	7498882CD1	g17933077	0.0	[Homo sapiens] cathepsin C
17	5524205CD1	g3170887	2.0E-30	[Mus musculus] ubiquitin-protein ligase E3-alpha Kwon, Y.T. et al. (1998) The mouse and human genes encoding the recognition component of the N-end rule pathway Proc. Natl. Acad. Sci. U.S.A. 95:7898-7903
18	7102342CD1	g20070980	0.0	[Homo sapiens] Similar to a disintegrin and metalloproteinase domain 18
18	7102342CD1	g4585653	9.1E-134	[Homo sapiens] tMDC III protein
19	4169939CD1	g13937848	8.5E-114	[Homo sapiens] (BC007028) Similar to elastase 3, pancreatic (protease E)
20	6539977CD1	g13897995	2.1E-136	[Homo sapiens] kallikrein 14 Hooper, J.D. et al. (2001) Identification and characterization of klk14, a novel kallikrein serine protease gene located on human chromosome 19q13.4 and expressed in prostate and skeletal muscle Genomics 73:117-122
21	7675588CD1	g13591753	9.7E-22	[Oryctolagus cuniculus] eppin Richardson, R.T. et al. (2001) Cloning and sequencing of human Eppin: a novel family of protease inhibitors expressed in the epididymis and testis. Gene 270:93-102
22	6244077CD1	g6648623	3.3E-76	[Homo sapiens] DNAj homolog
23	7498404CD1	g3062806	3.4E-121	[Homo sapiens] dolichol-phosphate-mannose synthase Tomita, S. et al. (1998) A homologue of Saccharomyces cerevisiae Dpm1p is not sufficient for synthesis of dolichol-phosphate-mannose in mammalian cells. J. Biol. Chem. 273:9249-9254
24	7391748CD1	g181182	1.8E-75	[Homo sapiens] preprocathepsin G Salvesen, G. et al. (1987) Molecular cloning of human cathepsin G: structural similarity to mast cell and cytotoxic T lymphocyte proteinases. Biochemistry 26:2289-2293
25	7499780CD1	g7546824	0.0	[Homo sapiens] lysyl hydroxylase 3 Rautavuoma, K. et al. (2000) Complete exon-intron organization of the gene for human lysyl hydroxylase 3 (LH3). Matrix Biol. 19:73-79

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
26	7499881CD1	g5257133	0.0	[Homo sapiens] protein O-mannosyl-transferase 1
27	7488579CD1	g4336577	1.8E-130	[Homo sapiens] putative mast cell mMCP-7-like II typtase Pallaoro, M. et al. (1999) Characterization of genes encoding known and novel human mast cell tryptases on chromosome 16p13.3. J. Biol. Chem. 274:3355-3362
28	7510521CD1	g1006657	4.5E-17	[Homo sapiens] cathepsin C Paris, A. et al. (1995) Molecular cloning and sequence analysis of human preprocathepsin C FEBS Lett. 369, 326-330
28	7510521CD1	334886 CTSC	3.9E-18	[Homo sapiens][Hydrolase; Protease (other than proteasomal)][Lysosome/vacuole; Cytoplasmic] Cathepsin C (dipeptidyl peptidase D), a lysosomal aminopeptidase and member of the papain family of cysteine proteinases, involved in activation of serine proteases in immune and inflammatory cells, mutations are associated with Papillon-Lefevre syndrome Hart, T. C. et al.
				Mutations of the cathepsin C gene are responsible for Papillon-Lefevre syndrome. J Med Genet 36, 881-7 (1999).

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	7994355CD1	774	HECT-domain (ubiquitin-transferase): D467-A774	HMMER_PFAM
			Ank repeat: V26-I58, S59-K92, N93-I125	HMMER_PFAM
			Ank repeat proteins: PF00023: L31-L46, G60-R69	BLIMPS_PFAM
			HECT-domain (ubiquitin-transferase): PF00632: V586-G592, W680-P707, L735-C766	BLIMPS_PFAM
			PROTEIN LIGASE UBIQUITIN CONJUGATION REPEAT UBIQUITIN D20PROTEIN DNA BINDING PROBABLE ONCOGENIC PD002225: P492-Y771	BLAST_PRODROM
			HECT DOMAIN DM01690; P39940 513-808: N475-L764; P51593 9-306: N475-Y771; A38919 785-1082: N475-A774; P53119 615-909: N475-S768	BLAST_DOMO
			Potential Phosphorylation Sites: S7 S15 S159 S176 S231 S250 S291 S338 S430 S669 S703 T96 T128 T216 T689 Y397	MOTIFS
			Potential Glycosylation Sites: N283 N722	MOTIFS
2	7475875CD1	703	Transmembrane domain: V113-A129, A141-V169 N-terminus is non-cytosolic	TMAP
			Tyrosine specific protein phosphatase BL00383: I137-G147, R170-C185	BLIMPS_BLOCKS
			Tyrosine specific protein phosphatases signature and profiles: L116-P174	PROFILESCAN
			Tyrosine specific protein phosphatases active site: I137-L149	MOTIFS
			Leucine zipper pattern: L333-L354	MOTIFS
			Potential Phosphorylation Sites: D18S21 S201 S341 S343 S448 S486 S496 S503 S594 T118 T272 T493 T605 T679 T687 T695	MOTIFS
			Potential Glycosylation Sites: N550	MOTIFS
3	71231882CD1	1256	Ubiquitin carboxyl-terminal hydrolases family 1: S41-L72	HMMER_PFAM
			Ubiquitin carboxyl-terminal hydrolase family 2: L285-K482	HMMER_PFAM
			Ubiquitin carboxyl-terminal hydrolases family 2 proteins BL00972: G42-L59, M129-N138, I158-C172, I288-D312, C446-Q467	BLIMPS_BLOCKS
			PROTEASE UBIQUITIN HYDROLASE UBIQUITINSPECIFIC ENZYME DEUBIQUITINATING CARBOXYLTERMINAL THIOLESTERASE PROCESSING CONJUGATION	BLAST_PRODROM
			UBIQUITIN CARBOXYL-TERMINAL HYDROLASES FAMILY 2; DM00659 P50101 209-458: N45-G299	BLAST_DOMO

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	7994355CD1	774	HECT-domain (ubiquitin-transferase): D467-A774 Ank repeat: V26-I58, S59-K92, N93-I125 Ank repeat proteins; PF00023; L31-L46, G60-R69 HECT-domain (ubiquitin-transferase): PF00632; V586-G592, W680-P707, L735-C766 PROTEIN LIGASE UBIQUITIN CONJUGATION REPEAT UBIQUITIN D20PROTEIN DNA BINDING PROBABLE ONCOGENIC PD002225: P492-Y771 HECT DOMAIN DM01690; P39940 513-808; N475-L764; P51593 9-306; N475-Y771; A38919 785-1082; N475-A774; P53119 615-909; N475-S768 Potential Phosphorylation Sites: S7 S15 S159 S176 S231 S250 S291 S338 S430 S669 S703 T96 T128 T216 T689 Y397 Potential Glycosylation Sites: N283 N722	HMMER_PFAM HMMER_PFAM BLIMPS_PFAM BLIMPS_PFAM BLAST_PRODROM BLAST_DOMO MOTIFS MOTIFS
2	7475875CD1	703	Transmembrane domain: V113-A129, A141-V169 N-terminus is non-cytosolic Tyrosine specific protein phosphatase BL00383: I137-G147, R170-C185 Tyrosine specific protein phosphatases signature and profiles: L116-P174 Tyrosine specific protein phosphatases active site: I137-L149 Leucine zipper pattern: L333-L354 Potential Phosphorylation Sites: D18S21 S201 S341 S343 S448 S486 S496 S503 S594 T118 T272 T493 T605 T679 T687 T695 Potential Glycosylation Sites: N550	TMAP BLIMPS_BLOCKS PROFILES SCAN MOTIFS MOTIFS MOTIFS MOTIFS
3	71231882CD1	1256	Ubiquitin carboxyl-terminal hydrolases family 1: S41-L72 Ubiquitin carboxyl-terminal hydrolase family 2: L285-K482 Ubiquitin carboxyl-terminal hydrolases family 2 proteins BL00972: G42-L59, M129-N138, I158-C172, I288-D312, C446-Q467 PROTEASE UBIQUITIN HYDROLASE UBIQUITINSPECIFIC ENZYME DEUBIQUITINATING CARBOXYL-TERMINAL THIOLESTERASE PROCESSING CONJUGATION UBIQUITIN CARBOXYL-TERMINAL HYDROLASES FAMILY 2; DM00659 P50101 209-458; N45-G299	HMMER_PFAM HMMER_PFAM BLIMPS_BLOCKS BLAST_PRODROM BLAST_DOMO

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			Ubiquitin carboxyl-terminal hydrolases family 2 signature 2: Y289-Y306	MOTIFS
			Potential Phosphorylation Sites: S113 S125 S143 S153 S179 S196 S232 S280 S326 S337 S402 S550 S554 S653 S729 S745 S868 S907 S971 S995 S1122 S1220 S1233 S1254 T124 T407 T516 T565 T608 T635 T651 T712 T737 T818 T857 T928 T1017 T1058 T1161 T1200 T1205 Y265 Y896 Y1104	MOTIFS
			Potential Glycosylation Sites: N39 N177 N194 N413 N441 N453 N525 N720 N977	MOTIFS
4	2875922CD1	755	PROTEIN CHROMOSOME C17A5.07C1 SMT4 SIMILARITY A SMALL REGION OF; PD009801: K602-L748	BLAST_PRODROM
			Potential Phosphorylation Sites: S18 S93 S107 S122 S126 S176 S300 S339 S363 S388 S405 S416 S423 S430 S447 S465 S488 S492 S521 S538 S655 S660 S733 T278 T390 T626 Y130	MOTIFS
			Potential Glycosylation Sites: N88 N349 N401 N709	MOTIFS
5	8158136CD1	1034	Ubiquitin carboxyl-terminal hydrolases family 1: V89-C120	HMMER_PFAM
			Ubiquitin carboxyl-terminal hydrolase family 2: G332-L420	HMMER_PFAM
			Ubiquitin carboxyl-terminal hydrolases family 2 proteins BL00972: G90-W107, G177-S186, Y336-D360, S363-L384	BLIMPS_BLOCKS
			PROTEASE UBIQUITIN HYDROLASE UBIQUITINSPECIFIC ENZYME	BLAST_PRODROM
			DEUBIQUITINATING CARBOXYL-TERMINAL THIOLESTERASE PROCESSING	
			F30A10.10 PROTEIN; PD185574: E641-A879	BLAST_PRODROM
			UBIQUITIN CARBOXYL-TERMINAL HYDROLASES FAMILY 2; DM00659P40818782-1103: Q200-I374, L94-N202	BLAST_DOMO
			Ubiquitin carboxyl-terminal hydrolases family 2 signature 2: Y336-Y354	MOTIFS
			Potential Phosphorylation Sites: S87 S158 S191 S445 S461 S531 S597 S600 S633 S659 S686 S878 S885 S887 S934 T17 T123 T143 T280 T398 T549 T760 T788 T819 T889 T913 T953 Y268 Y687	MOTIFS
			Potential Glycosylation Sites: N278 N427 N624 N883 N907 N951 N973	MOTIFS
6	5969491CD1	1236	Ubiquitin carboxyl-terminal hydrolases family 1: S41-L72	HMMER_PFAM
			Ubiquitin carboxyl-terminal hydrolase family 2: L285-K482	HMMER_PFAM
			Ubiquitin carboxyl-terminal hydrolases family 2 proteins BL00972: G42-L59, M129-N138, I158-C172, L288-D312, C446-Q467	BLIMPS_BLOCKS

Table 3

SEQ ID NO:	Incye Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			PROTEASE UBIQUITIN HYDROLASE UBIQUITINSPECIFIC ENZYME DEUBIQUITINATING CARBOXYLTERMINAL THIOLESTERASE PROCESSING CONJUGATION; PD017412: R181-E286; (P-value = 1.1e-10)	BLAST_PRODUM
			UBIQUITIN CARBOXYL-TERMINAL HYDROLASES FAMILY 2; DM00659 P50101 209-458: N45-G299	BLAST_DOMO
			Ubiquitin carboxyl-terminal hydrolases family 2 signature 2: Y289-Y306	MOTIFS
			Potential Phosphorylation Sites: S113 S125 S143 S153 S179 S196 S232 S280 S326 S337 S402 S550 S554 S653 S729 S745 S868 S951 S975 S1102 S1200 S1213 S1234 T124 T407 T516 T565 T608 T635 T651 T712 T737 T818 T857 T908 T997 T1038 T1141 T1180 T1185 Y265 Y896 Y1084	MOTIFS
			Potential Glycosylation Sites: N39 N177 N194 N413 N441 N453 N525 N720 N957	MOTIFS
7	7497367CD1	545	signal_cleavage: M1-P21	SPSCAN
			Signal Peptide: M1-P21	HMMER
			Leucine Rich Repeat: H170-T193, N266-P289, S122-A145, K362-Y385, C290-S313, R98-T121, N74-P97, N314-E337, S194-G217, S218-F241, A146-T169, E338-S361, C242-G265	HMMER_PFAM
			Leucine rich repeat C-terminal domain: N395-P446	HMMER_PFAM
			Leucine rich repeat N-terminal domain: P21-P48	HMMER_PFAM
			Leucine zipper pattern: L126-L147, L150-L171, L174-L195, L270-L291, L342-L363	MOTIFS
			Potential Phosphorylation Sites: S36 S218 S361 S484 S507 S531 T62 T197 T303 T375 T414 T457	MOTIFS
			Potential Glycosylation Sites: N74 N111 N119 N228 N266 N348 N359 N518	MOTIFS
8	7632424CD1	414	Glycoprotease family: K37-R372	HMMER_PFAM
			Glycoprotease family proteins BL01016: G325-R334, L355-A364, V39-V53, G76-H87, D108-A152, L166-A178, L189-C216, C249-T260	BLIMPS_BLOCKS
			O-sialoglycoprotein endopeptidase (M22) metallo-protease family signature PR00789: L40-V53, I112-L132, Q133-A152, L166-A178, L189-L210, L321-N330	BLIMPS_PRINTS
			PROTEIN HYDROLASE METALLOPROTEASE ZINC ENDOPEPTIDASE O-SIALOGLYCOPROTEIN PUTATIVE GLYCOPROTEASE INTERGENIC REGION PD002367: I38-I368	BLAST_PRODUM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			GLYCOPROTEASE FAMILY DM02236; [P05852]1-276: V39-I305; [P43764]1-281: V39-I305; [P47292]5-277: V39-I305; [P43122]33-333: V39-R334	BLAST_DOMO
			Potential Phosphorylation Sites: S15 S45 S67 S105 S209 S281 S404 T44 T115 T154 T158 T260 T301	MOTIFS
			Potential Glycosylation Sites: N342	MOTIFS
9	1804436CD1	611	Serine proteases, V8 family, histidine proteins; BL00672: T531-D547	BLIMPS_BLOCKS
			V8 serine protease family signature; PR00839: A370-I387, I530-F546, D547-G559	BLIMPS_PRINTS
			Potential Phosphorylation Sites: S2 S8 S12 S45 S62 S233 S284 S302 S307 S343 S503 S570 S576 S607 T81 T97 T220 T248 T289 T325 T382 T516 Y359 Y424 Y511	MOTIFS
			Potential Glycosylation Sites: N35 N75 N101 N246 N340 N566	MOTIFS
10	7486358CD1	147	signal_cleavage: M1-A28	SPSCAN
			Signal Peptide: M1-V25, M1-A28	HMMER
			Cystatin domain: P49-C142	HMMER_PFAM
			CYSTEINE PROTEASES INHIBITORS; DM00182[P01035]1-110: L51-C142; DM00182[P01034]34-143: L51-C142; DM00182[P28325]29-139: L51-C142; DM00182[P01038]29-138: L47-C142	BLAST_DOMO
			Potential Phosphorylation Sites: S76 T133 Y68	MOTIFS
			Potential Glycosylation Sites: N117 N139	MOTIFS
11	7472344CD1	624	PX domain: P273-P382	HMMER_PFAM
			SH3 domain: L3-I59	HMMER_PFAM
			Potential Phosphorylation Sites: S20 S31 S44 S96 S150 S196 S238 S297 S350 S383 S384 S426 S470 S549 T163 T197 T279 T341 T435 T448 T577 T611 Y320	MOTIFS
			Potential Glycosylation Sites: N446	MOTIFS
12	7192959CD1	283	signal_cleavage: M1-L47	SPSCAN
			F-box domain: L24-A72	HMMER_PFAM
			Potential Phosphorylation Sites: S181 T78 T214	MOTIFS
			Potential Glycosylation Sites: N141 N212 N273	MOTIFS
13	6169565CD1	142	Signal_cleavage: M1-G38	SPSCAN
			Signal Peptide: M1-G35	HMMER

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
14	7494717CD1	354	Potential Phosphorylation Sites: S52 S106 S135	MOTIFS
			Signal_cleavage: M1-G57	SPSCAN
			Ubiquitin-conjugating enzyme: G93-R250	HMMER_PFAM
			Ubiquitin-conjugating enzymes active site: F152-E219	PROFILERSCAN
			UBIQUITIN LIGASE ENZYME CONJUGATING CARRIER; MULTIGENE FAMILY PD000461: T86-P221	BLAST_PRODROM
			UBIQUITIN-CONJUGATING ENZYMES DM00225; P35133 1-146: R103-V253; P35128 1-148: R103-P221; P25865 2-149: R103-R245; P23566 2-149: R103-R245	BLAST_DOMO
			Potential Phosphorylation Sites: S4 S81 S89 S274 S342 S351 T141 T174 T248 T296 T344 Y240	MOTIFS
			Potential Glycosylation Sites: N172	MOTIFS
15	7497510CD1	89	Signal_cleavage: M1-S21	SPSCAN
			Signal Peptide: M1-E18, M1-S21, M1-L23, M1-A24	HMMER
			Kunitz/Bovine pancreatic trypsin inhibitor: C36-C86	HMMER_PFAM
			Pancreatic trypsin inhibitor (Kunitz) family signature: P44-K87	PROFILERSCAN
			Basic protease (Kunitz-type) inhibitor family signature PR00759: P33-T47, C61-G71, G71-C86	BLIMPS_PRINTS
			INHIBITOR PROTEASE SERINE GLYCOPROTEIN RECURSOR; SIGNAL FACTOR REPEAT TISSUE PD000222: C36-C86	BLAST_PRODROM
			ANIMAL KUNITZ-TYPE PROTEINASE INHIBITOR DM00114; S41399 5-61: D30-K87; P10646 48-110: L31-C86; P10646 112-175: L23-K87; P10280 1-55: C36-C86	BLAST_DOMO
			Pancreatic trypsin inhibitor (Kunitz) family signature: F64-C82	MOTIFS
			Potential Phosphorylation Sites: T27	MOTIFS
16	7498882CD1	419	Signal_cleavage: M1-G20	SPSCAN
			Signal Peptide: M1-C24, M1-G20, M1-A28, M1-R23	HMMER
			Papain family cysteine protease: G253-T414, L231-Q252	HMMER_PFAM
			Eukaryotic thiol (cysteine) proteases cysteine proteins BL00139: N251-P259, N360-G369, Y378-Y394	BLIMPS_BLOCKS
			Eukaryotic thiol (cysteine) proteases active sites: D334-F395	PROFILERSCAN
			Papain cysteine protease (C1) family signature PR00705: H361-D371, Y378-S384	BLIMPS_PRINTS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			PRECURSOR C HYDROLASE SIGNAL CATHEPSIN THIOL PROTEASE ZYMOGEN DIPEPTIDYL PEPTIDASE I; PD013797: G20-H230	BLAST_PRODOM
			PROTEASE PRECURSOR SIGNAL CYSTEINE PROTEINASE HYDROLASE THIOL ZYMOGEN CATHEPSIN GLYCOPROTEIN; PD000158: N251-R399	BLAST_PRODOM
			EUKARYOTIC THIOL (CYSTEINE) PROTEASES CYSTEINE DM00081P80067 124-457: Q252-P415, G124-G264; P53634 124-458: Q252-P415, G124-Y368; P09668 25-332: N251-P417, H155-G256; S52212 34-335: G253-P415	BLAST_DOMO
			Eukaryotic thiol (cysteine) proteases asparagine active site: Y378-I397	MOTIFS
			Eukaryotic thiol (cysteine) proteases histidine active site: T359-G369	MOTIFS
			Potential Phosphorylation Sites: S48 S164 S209 S374 T31 T73 T138 T144 T189 T197 T233 T281 T401 Y296	MOTIFS
			Potential Glycosylation Sites: N29 N53 N119	MOTIFS
17	5524205CD1	951	LIGASE UBIQUITIN E3 COMPONENT PROTEIN NENDRECOGNIZING N RECOGNIN CONJUGATION; PD007850: I776-H935	BLAST_PRODOM
			Potential Phosphorylation Sites: S25 S30 S78 S116 S129 S164 S190 S266 S308 S341 S400 S404 S440 S451 S454 S639 S643 S659 S734 S765 S786 S860 T70 T82 T196 T224 T233 T270 T286 T493 T520 T574 T660 T788	MOTIFS
			Potential Glycosylation Sites: N5 N114 N452 N477 N491 N882	MOTIFS
18	7102342CD1	668	Signal cleavage: M1-A16	SPSCAN
			Signal Peptide: M1-A16, M1-P19, M1-G20, M1-Q22	HMMER
			Non-cytosolic domain: M1-E574; Transmembrane domain: N575-A597; Cytosolic domain: R598-	TMHMMER
			Reprolysin family propeptide: H63-S176	HMMER_PFAM
			Reprolysin (M12B) family zinc metalloprotease: L186-P346	HMMER_PFAM
			Disintegrin: A301-C365	HMMER_PFAM
			Disintegrins signature: G287-D368	PROFILES CAN
			Disintegrin signature PR00289: C326-R345, E356-D368	BLIMPS PRINTS
			EGF-like domain signature 2: C536-C547	MOTIFS
			Potential Phosphorylation Sites: S42 S135 S160 S173 S234 S311 S340 S361 S376 S417 S419 S494 S564 S570 S616 S629 S648 T197 T210 T247 T452 T483 T642 T652 Y275	MOTIFS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			Potential Glycosylation Sites: N39 N125 N359 N492	MOTIFS
19	4169939CD1	206	Signal_cleavage: M1-G17	SPSCAN
			Signal Peptide: M2-G17, M2-G19, M1-P21, M1-S23, M1-S25, M1-G17, M1-G19, M2-S25	HMMER
			Trypsin: I36-I199	HMMER_PFAM
			Serine proteases, trypsin family, histidine proteins BL00134: S147-G170, P186-I199	BLIMPS_BLOCKS
			Apple domain proteins BL00495: V171-W198, S147-T187, A69-T103, V139-S173, G177-S205	BLIMPS_BLOCKS
			Type I fibronectin domain proteins BL01253: R67-T103, N104-G142, R146-C159, V168-T202	BLIMPS_BLOCKS
			Serine proteases, trypsin family, active sites: V134-C180	PROFILESAN
			Chymotrypsin serine protease family (S1) signature PR00722: A55-A69, R146-N158	BLIMPS_PRINTS
			PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZYMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR; D000046: S68-I199, G19-G165	BLAST_PRODOR
			TRYPSIN DM00018; P05805 1-236: V29-I203; P08217 28-265: A35-I203; P55091 29-265: E33-I203, R28-S40; P05208 31-267: S23-I203	BLAST_DOMO
			Serine proteases, trypsin family, serine active site: S147-N158	MOTIFS
			Potential Phosphorylation Sites: S26 T133	MOTIFS
			Potential Glycosylation Sites: N50	MOTIFS
20	6539977CD1	267	Signal_cleavage: M1-A30	SPSCAN
			Signal Peptide: P15-A30, M17-S34	HMMER
			Trypsin: I41-I260	HMMER_PFAM
			Kringle domain proteins; BL00021: C68-G85, R144-G165, D219-I260	BLIMPS_BLOCKS
			Serine proteases, trypsin family, histidine proteins BL00134: C68-C84, D214-G237, P247-I260	BLIMPS_BLOCKS
			Type I fibronectin domain proteins BL01253: C68-A81, S169-G207, K213-C226, Q229-T263	BLIMPS_BLOCKS
			Serine proteases, trypsin family, active sites: A60-T104, I199-A242	PROFILESAN
			Chymotrypsin serine protease family (S1) signature PR00722: G69-C84, T123-A137, K213-V225	BLIMPS_PRINTS
			REPEAT PRECURSOR GLYCOPROTEIN PD00120: G69-A81, D214-G222	BLIMPS_PRODOR
			PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZYMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR; PD000046: P155-I260, I41-P136	BLAST_PRODOR
			TRYPSIN DM00018; P32821 24-242: K40-M264; P19799 21-239: I41-M264; P06872 24-242: I41-M264; P06871 23-242: K40-M264	BLAST_DOMO

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			Serine proteases, trypsin family, serine active site: D214-V225	MOTIFS
			Serine proteases, trypsin family, histidine active site: I79-C84	MOTIFS
			Potential Phosphorylation Sites: S2 S34 S158 S186 S258 T263	MOTIFS
21	7675588CD1	86	Signal_cleavage: M1-A25	SPSCAN
			Signal Peptide: M1-G27	HMMER
			Non-cytosolic domain: M1-E86	MOTIFS
			WAP-type (Whey Acidic Protein) 'four-disulfide core:K31-F72	HMMER_PFAM
			WAP-type 'four-disulfide core' domain signature: P32-D71	PROFILES
			4-disulphide core signature PR00003C: C54-F63,Q47-C54, S64-F42	BLIMPS_PRINTS
			Potential Phosphorylation Sites: S76	MOTIFS
22	6244077CD1	232	DnaJ domain: N3-G69	HMMER_PFAM
			Nt-dnaJ domain proteins BL00636: D18-K34, F46-D66	BLIMPS_BLOCKS
			dnaJ domains signatures and profile: R24-P86	PROFILES
			DnaJ protein family signature PR00625: A14-D33, F46-D66, S184-I199	BLIMPS_PRINTS
			DNAJ PROTEIN HOMOLOG HSJ1 HSJ1 CHAPERONE NEURONE ALTERNATIVE	BLAST_PROD
			SPLICING TESTIS PD013370: Y91-K232	
			PROTEIN CHAPERONE DNAJ HEAT SHOCK DNA REPLICATION REPEAT ANTIGEN T	BLAST_PROD
			PD000231: N3-D66	
			NT-DNAJ DOMAIN; DM00098[S23509][1-108: M1-G105; DM00098[P25686][1-108: M1-G105; DM00098[P30725][1-102: M1-G106; DM00098[S34632][1-99: Y4-G106	BLAST_DOMO
			Binding-protein-dependent transport systems inner membrane comp. sign: F114-P142	MOTIFS
			Nt-dnaJ domain signature: F46-Y65	MOTIFS
			Potential Phosphorylation Sites: S15 S59 S63 S72 S85 S121 S184 T92 T195 T196	MOTIFS
23	7498404CD1	237	signal_cleavage: M1-S52	SPSCAN
			Glycosyl transferase: S28-R195	HMMER_PFAM
			Glycosyl transferases: PF00535: I61-T71, A109-D118; P < 3e-3	BLIMPS_PFAM
			SYNTHASE DOLICHOL-PHOSPHATE MANNOSE MANNOSE MANNOSE TRANSFERASE	BLAST_PROD
			DOLICHYLPHOSPHATE BETADMANNOSE TRANSFERASE GLYCOSYL-TRANSFERASE TRANSMEMBRANE PD151196: L175-A235	

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			GLYCOSYL-TRANSFERASE BIOSYNTHESIS GLYCOSYL SYNTHASE TRANSMEMBRANE N-ACETYL-GALACTOSAMINYLTRANSFERASE MEMBRANE PD000196: S28-R147	BLAST_PRODUM
			Potential Phosphorylation Sites: S9 S13 S21 S50 S84 S207 T33 T200	MOTIFS
			Potential Glycosylation Sites: N198	MOTIFS
24	7391748CD1	146	Trypsin: M1-I129	HMIMER_PFAM
			Serine proteases, trypsin family, histidine proteins BL00134: A86-G109, P116-I129	BLIMPS_BLOCKS
			Type I fibronectin domain proteins BL01253: K85-C98, C98-T132, R7-M43	BLIMPS_BLOCKS
			Serine proteases, trypsin family, active sites: L27-V114	PROFILESAN
			PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZYMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR PD000046: N14-G113	BLAST_PRODUM
			TRYPSIN; DM00018 P08311 21-241: M1-M133; DM00018 P28293 21-241: M1-M133; DM00018 P80219 1-221: M1-M133; DM00018 P20718 21-242: M1-M133	BLAST_DOMO
			ATP/GTP-binding site motif A (P-loop): G104-S111	MOTIFS
			Serine proteases, trypsin family, serine active site: A86-L97	MOTIFS
			Potential Phosphorylation Sites: S6 S42 S135 T47 T49 T132	MOTIFS
25	7499780CD1	696	signal_cleavage: M1-A24	SPSCAN
			Signal Peptide: M1-S23, M1-A24, M1-R27	HMIMER
			Lysyl hydrolase: Q293-P696, E37-G292	HMIMER_PFAM
			PROCOLLAGEN-LYSINE 2-OXOGLUTARATE 5-DIOXYGENASE PRECURSOR LYSYL HYDROXYLASE OXIDOREDUCTASE DIOXYGENASE SIGNAL IRON ; PD011980: L13-D285; PD011578: E500-P696; PD009947: E294-K499	BLAST_PRODUM
			LYSYL HYDROXYLASE CHAIN; DM07920 P24802 1-729: H297-P696, L12-G291; DM07920 Q02809 1-726: E294-P696, P15-E294	BLAST_DOMO
			Potential Phosphorylation Sites: S23 S25 S121 S325 S391 S441 S443 S454 S477 S621 S660 S692 T65 T159 T360 T429 T445 T476 Y64 Y346	MOTIFS
			Potential Glycosylation Sites: N63 N506	MOTIFS
26	7499881CD1	630	MIR domain: D339-H396, P275-V332, P201-P264	HMIMER_PFAM
			Dolichyl-phosphate-mannose-protein mannosyltransferase: M1-F172	HMIMER_PFAM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
			Intracellular domains: M1-E8, N58-S63, D111-R148, R503-D514, Q565-H630; Transmembrane domains: L9-T31, L35-F57, P64-I86, V91-G110, A149-V171, I480-L502, A515-F537, Y547-L564; Extracellular domains: Q32-R34, K87-G90, F172-N479, F538-L546	TMHMMER
			DETHIOBIOTIN SYNTHETASE PD02561: N479-R506, V332-T343	BLIMPS_PRODOME
			DOLICHYLPHOSPHATEMANNOSE PROTEIN MANNOSYLTRANSFERASE	BLAST_PRODOME
			GLYCOSYLTRANSFERASE GLYCOPROTEIN TRANSMEMBRANE ENDOPLASMIC RETICULUM MULTIGENE FAMILY; PD009956: H144-H288; PD005044: M1-L108;	
			ROTATED ABDOMEN PROTEIN PD116484: Q402-G523	BLAST_PRODOME
			LUMENAL DOMAIN; DM02305[P31382]36-704: H144-L560, M1-S115; DM02305[P47190]23-698: H144-L560, M1-S115; DM02305[P46971]28-693: H144-S574, Y3-L109; DM02305[P42934]30-714: H144-L564, F11-V117	BLAST_DOMO
			Leucine zipper pattern: L539-L560	MOTIFS
			Potential Phosphorylation Sites: S226 S336 S451 S610 T211 T291 T598 T603	MOTIFS
			Potential Glycosylation Sites: N318 N354 N422	MOTIFS
27	7488579CD1	242	signal_cleavage: M1-A25	SPSCAN
			Signal Peptides: M8-A23, M1-A23, M8-A25, M1-A25	HMMER
			Trypsin: I38-T242	HMMER_PFAM
			Type I fibronectin domain proteins BL01253: C66-A79, E136-N172, H224-C237	BLIMPS_BLOCKS
			Serine proteases, trypsin family, active sites: H64-Q107, V212-T242	PROFESCAN
			Chymotrypsin serine protease family (S1) signature PR00722: G67-C82, Q124-V138, H224-REPEAT PRECURSOR GLYCOPROTEIN PD00120: G67-A79, D128-L132, D225-G233	BLIMPS_PRINTS
			PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZYMOGEN GLYCOPROTEIN	BLIMPS_PRODOME
			FAMILY MULTIGENE FACTOR PD000046: D106-K238, I38-G220	BLAST_PRODOME
			TRYPSIN; DM00018[P15157]31-270: I38-T242; DM00018[P02844]29-268: I38-T242; DM00018[P21845]31-271: G37-T242; DM00018[P15944]31-270: I38-G241	BLAST_DOMO
			Serine proteases, trypsin family, histidine active site: L77-C82	MOTIFS
			Serine proteases, trypsin family, serine active site: D225-V236	MOTIFS
			Potential Phosphorylation Sites: S54 T125	MOTIFS
			Potential Glycosylation Sites: N139	MOTIFS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
28	7510521CD1	48	Signal_cleavage: M1-G20	SPSCAN
			Signal Peptide: M1-G20	HMMER
			Signal Peptide: M1-C24	HMMER
			Signal Peptide: M1-A28	HMMER
			Signal Peptide: M1-R23	HMMER
			Plant lipid transfer proteins signature: M1-K48	PROFILESAN
			Potential Phosphorylation Sites: T31	MOTIFS
			Potential Glycosylation Sites: N29	MOTIFS

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
29/7994355CB1/4384	1-677, 1-779, 432-679, 432-950, 432-1041, 432-4364, 863-1620, 930-1257, 1087-1751, 1104-1357, 1338-1984, 1386-1900, 1446-1630, 1659-1969, 1659-2021, 1661-2064, 1812-2430, 1839-2144, 1843-2450, 1847-2151, 1872-2135, 1872-2348, 1976-2242, 1976-2562, 2002-2245, 2182-2636, 2212-2466, 2277-2797, 2277-3069, 2298-2770, 2376-2812, 2381-2672, 2408-2450, 2446-2937, 2476-3046, 2499-2718, 2499-2973, 2577-2886, 2666-3114, 2701-3285, 2766-3485, 2963-3504, 3039-3266, 3039-3284, 3068-3538, 3083-3701, 3088-3331, 3095-3607, 3110-3779, 3120-3528, 3122-3690, 3128-3267, 3128-3382, 3128-3396, 3130-3556, 3130-3605, 3178-3451, 3180-3446, 3220-3700, 3274-3514, 3274-3551, 3274-3560, 3308-3725, 3309-3491, 3311-3627, 3314-3957, 3357-3530, 3357-3540, 3383-3632, 3383-3653, 3389-3546, 3409-3709, 3419-3528, 3430-3527, 3440-3585, 3491-3721, 3491-3725, 3491-4023, 3494-3646, 3494-3738, 3500-3990, 3572-3826, 3572-3962, 3579-4002, 3597-4006, 3625-3885, 3660-3922, 3665-4247, 3691-4287, 3701-4353, 3704-3936, 3704-4198, 3709-3925, 3733-4376, 3768-4307, 3774-4031, 3782-4324, 3799-3983, 3802-4000, 3834-4333, 3857-4127, 3859-4009, 3888-4383, 3900-4382, 3933-4181, 3933-4352, 3933-4360, 3942-4200, 3944-4382, 4006-4265, 4025-4379, 4036-4333, 4041-4265, 4049-4275, 4049-4359, 4058-4384, 4070-4361, 4094-4343, 4094-4351, 4150-4352, 4150-4363, 4150-4366, 4150-4380, 4171-4384, 4193-4384, 4196-4360, 4246-4367, 4249-4374, 4257-4384, 4311-4384
30/7475875CB1/4007	1-180, 25-740, 433-1078, 809-1509, 961-1177, 961-1575, 974-1232, 1327-1981, 1373-1926, 1420-1875, 1611-1902, 1620-1905, 1705-1850, 1705-1951, 1705-2022, 1705-2052, 1705-2067, 1705-2068, 1705-2249, 1705-2279, 1713-2264, 1727-2068, 1727-2157, 1755-2385, 1774-2290, 1800-2021, 1800-2258, 1800-2266, 1800-2346, 1802-2269, 1885-2113, 1885-2409, 1885-2410, 1913-2316, 1920-2309, 1959-2047, 2011-2329, 2078-2312, 2159-2247, 2219-2770, 2303-2955, 2381-2812, 2428-2642, 2430-2875, 2453-3077, 2456-2994, 2467-3100, 2565-3092, 2568-3216, 2572-3061, 2577-3035, 2603-3036, 2609-3045, 2611-3030, 2614-3212, 2616-3180, 2630-2899, 2632-3221, 2635-3211, 2638-3159, 2641-3134, 2655-3148, 2656-2943, 2660-3272, 2663-3361, 2672-3180, 2672-3332, 2672-3337, 2674-3162, 2678-3188, 2681-2925, 2681-3192, 2684-3262, 2690-2976, 2703-3216, 2705-3376, 2711-3279, 2712-3216, 2715-3238, 2715-3410, 2715-3461, 2722-3304, 2724-3273, 2730-2969, 2732-2844, 2732-2995, 2734-3290, 2736-3093, 2754-3452, 2754-3454, 2767-3422, 2781-3285, 2787-3338, 2790-3359, 2792-3313, 2794-3005,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
	2794-3066, 2794-3250, 2794-3276, 2797-3412, 2806-2970, 2811-3067, 2822-3373, 2833-3451, 2836-3326, 2836-3714, 2854-3355, 2876-3419, 2879-3447, 2885-3153, 2885-3461, 2888-3497, 2922-3451, 2929-3488, 2944-3547, 3010-3447, 3015-3521, 3021-3564, 3036-3663, 3059-3650, 3063-3475, 3070-3213, 3074-3442, 3094-3667, 3094-3668, 3100-3663, 3107-3556, 3116-3679, 3129-3432, 3136-3845, 3149-3721, 3173-3668, 3184-3679, 3186-3448, 3188-3703, 3231-3754, 3258-3783, 3266-3790, 3287-3948, 3319-3844, 3320-3811, 3328-3795, 3330-4004, 3331-3960, 3341-3594, 3346-3555, 3346-3839, 3359-3914, 3388-3602, 3388-3627, 3388-3638, 3388-3664, 3390-4001, 3409-3983, 3420-4007, 3424-3937, 3425-4000, 3426-3790, 3438-3981, 3448-4005, 3463-4003, 3468-3896, 3473-3711, 3493-3941, 3552-4007, 3554-4007, 3555-4007, 3560-4007, 3573-4007, 3581-4007, 3582-4007, 3591-4007, 3594-4007, 3604-4007, 3615-3992, 3620-4007, 3659-4007, 3679-4007, 3790-4007, 3796-4007, 3805-4007, 3810-3951, 3907-4007
31/71231882CB1/4524	1-372, 6-295, 8-714, 20-270, 21-590, 21-620, 21-651, 37-651, 87-4524, 102-651, 114-677, 129-850, 129-851, 131-726, 131-761, 131-776, 131-798, 132-779, 231-488, 306-677, 437-677, 566-1211, 757-798, 872-1236, 1045-1219, 1045-1332, 1045-1522, 1045-1558, 1045-1576, 1048-1274, 1053-1576, 1189-1839, 1431-1912, 1469-1772, 1485-2017, 1744-2336, 1774-2519, 1904-2425, 1930-2569, 1955-2713, 2018-2497, 2072-2716, 2153-2694, 2172-2352, 2172-2386, 2172-2615, 2172-2655, 2172-2694, 2172-2724, 2172-2730, 2172-2742, 2255-2463, 2255-2911, 2271-2488, 2370-2772, 2400-2771, 2491-2724, 2491-2805, 2498-2686, 2501-2687, 2637-3165, 2696-3192, 2718-3058, 2821-3239, 2959-3204, 3105-4037, 3168-3564, 3204-3835, 3208-3588, 3213-3594, 3253-3596, 3258-4032, 3267-4032, 3286-3655, 3294-4032, 3295-4032, 3296-3753, 3297-4032, 3302-3930, 3344-4032, 3362-3591, 3375-3824, 3413-4032, 3488-3680, 3488-3812, 3488-4018, 3488-4079, 3510-4032, 3530-4117, 3555-4144, 3566-4503, 3593-3840, 3593-3872, 3593-4200, 3594-3907, 3600-4508, 3606-4236, 3626-4347, 3626-4358, 3634-4141, 3648-4428, 3653-4364, 3656-4102, 3659-4323, 3661-4368, 3702-4524, 3709-3983, 3733-3919, 3758-3972, 3758-3997, 3874-4220, 3876-4206, 3900-4441, 3911-4524

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
32/2875922CB1/3250	1-565, 9-482, 112-594, 190-821, 190-860, 256-469, 318-1062, 394-861, 473-668, 473-977, 638-1128, 682-943, 682-1241, 1108-1735, 1141-1573, 1414-1719, 1414-1930, 1414-1933, 1414-1936, 1414-1937, 1414-1951, 1414-1985, 1414-2050, 1453-2185, 1458-1983, 1616-1894, 1616-2120, 1626-2140, 1663-2321, 1672-1977, 1689-2110, 1708-2045, 1708-2071, 1708-2083, 1708-2088, 1728-2171, 1731-2368, 1753-2415, 1787-2376, 1813-2117, 1820-2568, 1841-2324, 1850-2403, 1858-2330, 1870-2511, 1881-2123, 1895-2246, 1903-2180, 1923-2059, 1924-2631, 1925-2466, 1977-2534, 1992-2521, 2035-2658, 2049-2687, 2059-2197, 2060-2197, 2063-2658, 2067-2618, 2087-2673, 2147-2742, 2157-2742, 2161-2673, 2176-2622, 2194-2584, 2214-2519, 2217-2673, 2226-2742, 2232-2523, 2232-2716, 2233-2481, 2266-2631, 2296-2700, 2323-2750, 2333-2750, 2339-2749, 2365-2742, 2377-2879, 2444-2708, 2683-3152, 2688-3110, 2715-3250, 2737-3203, 2750-3196, 2783-3016, 2798-3075, 2800-3128, 2866-3196, 2963-3195, 3039-3182, 3113-3250
33/8158136CB1/3834	1-523, 1-633, 1-638, 1-3802, 19-869, 26-302, 26-340, 26-603, 34-300, 34-528, 43-595, 61-947, 132-833, 207-447, 221-784, 419-916, 433-1161, 509-1021, 646-1308, 672-1213, 679-1111, 719-1106, 735-1120, 766-1264, 775-1408, 792-1292, 804-1351, 810-1309, 823-1510, 839-1163, 855-1430, 895-1147, 895-1399, 897-1201, 919-1168, 970-1202, 983-1237, 990-1391, 1013-1655, 1041-1537, 1041-1541, 1065-1524, 1094-1593, 1100-1369, 1116-1368, 1116-1654, 1116-1691, 1117-1642, 1127-1389, 1132-1771, 1184-1433, 1190-1406, 1217-1629, 1220-1448, 1237-1711, 1246-1491, 1252-1810, 1329-1738, 1368-2054, 1387-1933, 1391-1668, 1392-1956, 1437-1597, 1460-1688, 1461-1609, 1490-2065, 1501-1799, 1523-1806, 1529-1784, 1556-1820, 1586-1841, 1686-2203, 1717-2475, 1722-1981, 1795-2397, 1798-2537, 1848-2107, 1855-2601, 1900-2578, 1908-2625, 1910-2273, 1911-2623, 1995-2606, 2002-2705, 2019-2628, 2040-2775, 2061-2763, 2061-2804, 2061-2845, 2066-2694, 2083-2769, 2089-2678, 2091-2797, 2097-2827, 2102-2550, 2123-2527, 2133-2421, 2178-2619, 2179-2473, 2191-2804, 2207-2706, 2210-2921,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
34/596949/ICB1/4493	<p>2215-2901, 2215-2938, 2247-2496, 2256-3023, 2261-2477, 2266-2865, 2267-2543, 2299-2604, 2325-2786, 2329-2838, 2329-2968, 2336-3076, 2336-3162, 2336-3209, 2361-2624, 2366-2938, 2372-2875, 2405-3011, 2435-3071, 2449-2718, 2581-3450, 2636-3304, 2666-3445, 2741-3405, 2812-3481, 2816-3480, 2824-3480, 2828-3480, 2832-3441, 2860-3478, 2863-3436, 2866-3483, 2888-3588, 2896-3450, 2907-3482, 2916-3490, 2922-3464, 2923-3467, 2926-3376, 2941-3482, 2954-3482, 2961-3479, 2966-3475, 2982-3490, 2997-3448, 3016-3804, 3034-3503, 3042-3476, 3043-3487, 3047-3483, 3062-3488, 3064-3482, 3067-3483, 3087-3494, 3093-3483, 3099-3375, 3112-3487, 3118-3488, 3124-3323, 3137-3390, 3137-3558, 3144-3357, 3159-3489, 3176-3490, 3177-3444, 3178-3381, 3186-3391, 3186-3416, 3187-3803, 3205-3470, 3222-3487, 3230-3804, 3232-3804, 3237-3495, 3257-3698, 3257-3798, 3268-3483, 3270-3488, 3281-3465, 3311-3490, 3320-3519, 3320-3616, 3331-3482, 3334-3481, 3334-3487, 3334-3494, 3343-3805, 3357-3808, 3398-3483, 3400-3804, 3403-3833, 3412-3834, 3426-3804, 3430-3802</p> <p>1-568, 1-598, 1-666, 65-4493, 80-621, 92-316, 92-655, 107-828, 107-829, 109-704, 109-739, 109-754, 109-776, 110-757, 209-466, 284-655, 415-655, 544-1189, 735-776, 850-1211, 1023-1197, 1023-1310, 1023-1500, 1023-1536, 1023-1554, 1026-1252, 1032-1554, 1049-1498, 1136-1420, 1167-1407, 1167-1817, 1235-1472, 1263-1554, 1301-1517, 1409-1890, 1447-1750, 1463-1995, 1722-2314, 1752-2497, 1882-2403, 1908-2547, 1933-2691, 1995-2475, 2051-2692, 2133-2672, 2150-2330, 2150-2364, 2150-2593, 2150-2633, 2150-2672, 2150-2701, 2150-2708, 2150-2720, 2153-2888, 2233-2441, 2249-2466, 2298-2901, 2348-2750, 2378-2749, 2378-2803, 2392-2842, 2469-2702, 2469-2783, 2476-2664, 2479-2665, 2498-2803, 2538-2845, 2568-2827, 2581-3326, 2583-3077, 2587-3122, 2592-2994, 2632-2952, 2636-3358, 2648-2910, 2650-2901, 2666-2940, 2677-2950, 2698-2914, 2716-3143, 2756-3243, 2793-3022, 2793-3195, 2793-3368, 2793-3457, 2793-3473, 2793-3475, 2793-3502, 2948-3511, 2959-3266, 3043-3955, 3060-3516, 3122, 2883-3411, 2907-3499, 2919-3490, 2943-3259, 2945-3502, 2948-3511, 2959-3266, 3043-3955, 3060-3516, 3086-3367, 3086-3482, 3094-3526, 3118-3425, 3122-3716, 3131-3512, 3139-3428, 3161-3520, 3171-3514, 3176-3950, 3185-3950, 3204-3511, 3208-3573, 3211-3526, 3212-3950, 3213-3950, 3214-3671, 3215-3950, 3220-3848, 3262-3950, 3280-3509, 3296-3742, 3331-3950, 3346-3512, 3406-3597, 3406-3730, 3406-3936, 3406-3996, 3428-3950, 3448-4035, 3473-4021, 3495-3685, 3504-4243, 3511-3758, 3511-3790, 3511-4118, 3512-3825, 3518-4427, 3525-4153, 3544-4276, 3545-4265, 3554-4059, 3566-4347, 3571-4282, 3574-3937, 3577-4158, 3579-4286, 3620-4472, 3627-3901, 3651-3837, 3676-3890, 3676-3915, 3792-4138, 3799-4124, 3818-4360, 3830-4442</p>

Table 4

Polynucleotide SEQ ID NO: / Incyte ID/ Sequence Length	Sequence Fragments
35/7497367CB1/2921	1-798, 33-671, 37-635, 48-194, 48-197, 50-191, 51-197, 59-194, 59-197, 95-197, 95-1824, 303-663, 332-434, 356-396, 356-410, 526-1210, 552-1231, 1199-1347, 1199-1799, 1199-1864, 1199-1871, 1199-1968, 1199-1992, 1199-2061, 1199-2064, 1199-2123, 1200-1968, 1452-1563, 1871-2287, 1871-2601, 1871-2762, 1874-2428, 1874-2467, 1874-2496, 2047-2707, 2051-2712, 2064-2585, 2109-2552, 2156-2687, 2193-2646, 2196-2752, 2291-2921, 2423-2907, 2435-2921, 2756-2858, 2756-2862, 2759-2861, 2780-2820, 2780-2834, 2867-2921
36/7632424CB1/2572	1-714, 24-727, 33-612, 49-776, 49-811, 50-586, 60-720, 71-700, 85-885, 87-591, 87-618, 98-895, 122-664, 122-754, 123-606, 210-707, 251-928, 370-413, 370-565, 370-633, 370-737, 420-924, 420-1020, 453-704, 453-959, 468-937, 515-1213, 575-1277, 580-826, 580-1180, 581-1188, 599-1257, 780-1292, 788-1404, 809-961, 862-1416, 902-1548, 906-1144, 936-1683, 945-1613, 956-1108, 962-1108, 963-1108, 969-1108, 969-1167, 985-1108, 989-1534, 1021-1736, 1025-1680, 1031-1736, 1063-1564, 1063-1719, 1084-1880, 1090-1714, 1106-1153, 1137-1768, 1150-1797, 1151-1790, 1195-1798, 1197-1797, 1201-1810, 1219-1850, 1228-1804, 1240-1594, 1241-1664, 1338-1624, 1387-1999, 1417-2183, 1456-2180, 1497-1756, 1545-2118, 1642-2083, 1660-2052, 1665-2086, 1667-1717, 1670-2204, 1679-1967, 1693-1967, 1695-1972, 1695-2078, 1695-2083, 1700-2078, 1749-2063, 1754-2183, 1755-2063, 1774-2018, 1774-2079, 1774-2088, 1810-2041, 1817-2074, 1826-2086, 1870-2075, 1897-2204, 1941-2183, 2101-2556, 2101-2572, 2108-2359, 2117-2572, 2264-2570, 2383-2572
37/1804436CB1/3758	1-426, 23-459, 66-703, 87-455, 159-805, 162-643, 162-677, 170-811, 455-1225, 497-1051, 540-1201, 588-878, 626-1361, 637-1122, 670-1491, 731-1134, 774-1469, 782-1261, 792-1211, 819-1398, 863-1409, 893-1531, 911-1409, 991-1577, 998-1236, 1003-1138, 1012-1394, 1015-1630, 1079-1546, 1086-1673, 1087-1597, 1087-1599, 1088-1602, 1153-1435, 1187-1522, 1199-1467, 1332-1948, 1358-3567, 1374-1951, 1397-1738, 1399-1988, 1411-1844, 1424-1723, 1438-1723, 1492-2157, 1572-2168, 1601-2175, 1608-1884, 1632-1902, 1635-1900, 1643-2079, 1693-2337, 1695-2296, 1695-2318, 1695-2379, 1695-2406, 1796-2046, 1807-2082, 1807-2276, 1807-2280, 1807-2311, 1823-2331, 1846-2389, 1860-2393, 1941-2206, 2002-2224, 2024-2243, 2050-2227, 2284-2773, 2329-2906, 2392-2881, 2445-2749, 2449-2792, 2449-2929, 2449-2935, 2449-2948, 2449-2969, 2449-3034, 2449-3043, 2449-3052, 2449-3061, 2449-3082, 2457-2746, 2503-2782, 2524-2776, 2670-3308, 2681-3222, 2685-2875, 2685-3230, 2697-3334, 2704-2943, 2704-2946, 2704-2947, 2704-2949, 2713-3322, 2718-3329, 2736-3331, 2737-3282, 2749-3369,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
	2750-3407, 2753-3035, 2759-3156, 2763-3306, 2767-3369, 2771-3405, 2775-3027, 2775-3400, 2783-3287, 2785-3034, 2803-3199, 2803-3456, 2809-3350, 2815-3158, 2820-3402, 2826-3078, 2826-3453, 2840-3357, 2854-3317, 2855-3369, 2857-3314, 2857-3318, 2858-3098, 2865-3102, 2865-3481, 2866-3149, 2878-3539, 2881-3551, 2901-3439, 2906-3553, 2908-3347, 2918-3514, 2920-3555, 2923-3502, 2937-3094, 2944-3201, 2944-3213, 2947-3567, 2960-3562, 2975-3223, 2975-3248, 2975-3553, 2994-3207, 2999-3257, 3001-3270, 3004-3290, 3008-3233, 3011-3464, 3014-3276, 3015-3252, 3026-3280, 3027-3281, 3035-3567, 3046-3274, 3046-3306, 3046-3314, 3046-3315, 3052-3567, 3057-3324, 3061-3567, 3063-3326, 3069-3330, 3072-3567, 3074-3567, 3076-3378, 3077-3563, 3090-3360, 3092-3567, 3100-3559, 3102-3560, 3105-3564, 3108-3556, 3110-3489, 3113-3562, 3113-3567, 3117-3567, 3125-3560, 3130-3387, 3135-3559, 3138-3436, 3140-3394, 3141-3367, 3141-3559, 3144-3567, 3151-3559, 3155-3559, 3157-3562, 3157-3563, 3162-3554, 3169-3546, 3171-3560, 3171-3562, 3175-3559, 3176-3428, 3180-3235, 3180-3559, 3181-3564, 3181-3693, 3181-3758, 3187-3410, 3197-3563, 3199-3566, 3205-3559, 3210-3559, 3218-3442, 3218-3445, 3218-3455, 3220-3559, 3240-3565, 3243-3510, 3246-3559, 3265-3559, 3266-3559, 3269-3559, 3286-3507, 3286-3552, 3286-3559, 3286-3562, 3289-3560, 3290-3544, 3291-3559, 3294-3529, 3297-3566, 3335-3561, 3352-3567, 3359-3566, 3388-3567, 3420-3557
38/7486358CB1/1036	1-698, 1-832, 143-831, 148-831, 588-1030, 588-1036, 592-831
39/7472344CB1/3651	1-654, 179-734, 214-871, 515-980, 542-1000, 812-1067, 852-1392, 958-1474, 1002-1222, 1031-1598, 1037-1799, 1039-1738, 1062-1440, 1200-1824, 1241-1860, 1281-1824, 1675-1960, 1675-1986, 1675-2039, 1675-2167, 1675-2184, 1675-2207, 1675-2299, 1675-2324, 1709-2324, 1748-2324, 2014-2637, 2204-2328, 2235-2328, 2550-2826, 2660-2889, 2689-2938, 2689-2952, 2692-2889, 2693-2923, 2693-2984, 2693-3189, 2694-3222, 2695-3173, 2697-2990, 2697-3183, 2699-3248, 2701-3002, 2703-2896, 2763-3021, 2787-3021, 2834-3218, 2834-3238, 2837-3388, 2845-3232, 3005-3181, 3005-3317, 3011-3414, 3015-3409, 3074-3314, 3081-3321, 3081-3651, 3084-3345, 3084-3601
40/7192959CB1/1989	1-517, 170-815, 170-870, 170-882, 172-564, 172-775, 172-810, 174-871, 197-769, 222-826, 430-889, 539-1122, 559-802, 559-1127, 573-875, 621-1361, 640-1063, 666-904, 720-991, 963-1617, 991-1282, 991-1497, 1012-1622, 1069-1618, 1145-1539, 1275-1507, 1275-1630, 1370-1589, 1370-1622, 1378-1619, 1424-1743, 1465-1716, 1645-1989, 1730-1989, 1751-1935, 1751-1989, 1766-1989, 1773-1989, 1780-1989, 1784-1989, 1785-1989, 1829-1989, 1840-1989, 1845-1989, 1853-1989, 1880-1989, 1886-1989

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
41/6169565CB1/1629	1-747, 275-554, 275-957, 275-992, 275-1064, 275-1082, 282-1085, 643-1230, 690-1438, 694-1407, 733-1590, 842-1629, 863-1629, 868-1628, 871-1587, 935-1629
42/7494717CB1/3166	1-237, 90-702, 100-922, 100-955, 104-842, 127-307, 137-222, 137-776, 137-784, 141-405, 144-307, 144-608, 150-279, 150-613, 166-600, 171-756, 193-977, 391-696, 394-885, 424-887, 459-649, 496-1311, 506-916, 541-1014, 566-864, 566-1031, 566-1077, 574-989, 629-1193, 637-1266, 638-1116, 639-1235, 648-1283, 673-866, 684-1028, 689-946, 701-1248, 735-977, 748-1311, 759-1002, 819-1297, 824-1450, 842-1479, 853-1495, 853-1535, 857-1079, 878-1118, 883-1567, 893-1393, 912-1320, 924-1531, 944-1549, 954-1235, 1018-1486, 1020-1264, 1032-1331, 1036-1515, 1037-1467, 1053-1179, 1072-1560, 1151-1760, 1196-1484, 1217-1543, 1268-1486, 1280-1579, 1286-1750, 1295-1560, 1320-1641, 1379-1625, 1390-1645, 1398-1526, 1437-1756, 1444-1689, 1476-1736, 1501-1733, 1719-1950, 1720-2098, 1798-2096, 1827-2175, 1879-2107, 1889-2144, 1937-2197, 1954-2196, 1958-2248, 1958-2490, 1974-2216, 2001-2257, 2003-2646, 2012-2208, 2045-2280, 2046-2248, 2113-2353, 2114-2309, 2144-2409, 2204-2454, 2304-2654, 2338-2537, 2340-2552, 2346-2583, 2357-2702, 2379-2596, 2425-2703, 2477-2731, 2482-2697, 2495-2740, 2503-2697, 2522-2782, 2630-3132, 2631-2844, 2634-3116, 2649-3126, 2665-2904, 2737-3020, 2737-3143, 2737-3166, 2814-3027, 2818-3007, 2828-3074, 2861-2999, 2965-3109
43/7497510CB1/399	1-144, 69-338, 156-335, 192-399, 212-399
44/7498882CB1/1811	1-237, 15-181, 21-247, 22-466, 22-519, 23-197, 27-253, 27-289, 28-302, 28-366, 29-280, 34-315, 35-161, 35-278, 35-281, 37-248, 37-291, 48-200, 48-309, 48-421, 58-310, 59-359, 60-294, 60-871, 61-184, 61-273, 61-299, 61-304, 61-315, 61-328, 61-333, 61-334, 61-391, 61-657, 61-703, 62-251, 62-299, 62-315, 62-321, 62-344, 62-460, 63-319, 63-442, 63-592, 64-302, 65-274, 65-307, 65-308, 65-326, 65-328, 65-329, 65-342, 65-630, 66-283, 66-301, 66-305, 66-315, 66-322, 66-325, 66-326, 66-331, 66-343, 66-440, 66-634, 67-196, 67-248, 67-275, 67-292, 67-294, 67-302, 67-316, 67-381, 67-490, 68-300, 68-306, 68-645, 69-308, 69-460, 70-300, 70-321, 70-328, 70-331, 70-339, 70-354, 70-356, 70-423, 71-171, 71-673, 72-287, 73-351, 73-386, 76-364, 79-346, 80-334, 82-346, 83-346, 83-348, 85-329, 85-338, 85-360, 85-361, 85-364, 85-374, 85-526, 85-564, 86-284, 86-310, 86-322, 86-329, 86-352, 86-355, 86-361, 86-362, 86-365, 86-373, 86-381, 86-383, 86-386, 86-387, 86-432, 86-450, 86-549, 86-555, 86-574, 86-619, 86-664, 86-668, 86-669, 86-675, 86-680, 86-714, 87-446, 92-619, 96-380, 102-472, 103-308, 103-329, 103-684, 104-351,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
	104-1374, 107-290, 117-359, 117-401, 118-365, 129-381, 129-387, 135-388, 141-403, 156-431, 162-822, 165-632, 166-448, 172-576, 172-769, 173-386, 183-415, 183-601, 197-534, 210-621, 211-518, 224-403, 225-427, 243-528, 261-522, 267-553, 393-851, 412-841, 443-683, 447-699, 449-715, 449-742, 451-698, 453-721, 454-601, 459-778, 474-762, 487-744, 487-750, 490-801, 507-705, 507-764, 509-759, 522-827, 527-752, 527-785, 553-867, 565-838, 565-869, 582-869, 584-871, 595-843, 597-821, 597-856, 602-845, 602-871, 603-826, 608-706, 632-868, 656-853, 663-766, 674-829, 693-1238, 694-983, 868-1351, 868-1509, 870-962, 871-1143, 873-1154, 873-1292, 880-1109, 909-1168, 909-1457, 913-1175, 915-1156, 915-1379, 916-1116, 916-1143, 930-1176, 933-1232, 934-1194, 935-1191, 935-1368, 936-1181, 938-1204, 939-1177, 941-1265, 974-1195, 990-1264, 991-1257, 1005-1254, 1008-1252, 1020-1253, 1020-1363, 1024-1314, 1027-1315, 1027-1322, 1028-1270, 1039-1285, 1039-1329, 1043-1292, 1044-1265, 1044-1351, 1046-1289, 1046-1303, 1048-1331, 1051-1714, 1055-1267, 1055-1361, 1079-1358, 1079-1584, 1086-1381, 1087-1276, 1087-1351, 1092-1598, 1099-1383, 1101-1391, 1107-1530, 1109-1370, 1112-1430, 1115-1377, 1115-1576, 1117-1624, 1123-1317, 1125-1406, 1130-1438, 1131-1315, 1131-1318, 1132-1440, 1135-1403, 1136-1425, 1141-1391, 1143-1427, 1145-1775, 1146-1414, 1149-1407, 1149-1592, 1152-1393, 1152-1406, 1154-1380, 1154-1402, 1154-1410, 1158-1772, 1161-1260, 1161-1413, 1161-1430, 1165-1390, 1166-1773, 1167-1740, 1175-1437, 1175-1463, 1176-1419, 1176-1449, 1176-1739, 1184-1729, 1188-1438, 1188-1456, 1195-1676, 1201-1772, 1207-1470, 1210-1456, 1219-1414, 1219-1497, 1246-1773, 1249-1403, 1250-1552, 1255-1523, 1256-1775, 1265-1555, 1275-1711, 1289-1811, 1300-1787, 1304-1547, 1306-1775, 1312-1775, 1315-1787, 1319-1747, 1324-1775, 1325-1431, 1327-1775, 1331-1787, 1332-1787, 1333-1781, 1336-1582, 1347-1593, 1352-1781, 1354-1782, 1355-
	1783, 1357-1644, 1357-1787, 1358-1790, 1359-1781, 1360-1787, 1360-1790, 1361-1775, 1361-1794, 1364-1777, 1364-1783, 1365-1775, 1365-1781, 1367-1734, 1367-1774, 1368-1781, 1370-1749, 1371-1791, 1373-1782, 1374-1781

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
45/5524205CB1/4407	1-981, 25-703, 144-868, 183-482, 286-913, 531-1263, 614-868, 615-1263, 617-915, 617-1000, 769-1034, 871-1014, 871-1186, 924-1568, 926-1167, 926-1243, 926-1372, 926-1568, 931-1514, 1004-1385, 1004-1595, 1008-1595, 1049-1294, 1122-1654, 1155-1399, 1155-4378, 1160-1457, 1160-1501, 1400-1600, 1400-1737, 1409-1600, 1409-1737, 1601-1737, 1601-1823, 1627-1931, 1627-2195, 1627-2218, 1687-2376, 1772-2386, 1917-2373, 1939-2661, 2008-2267, 2049-2394, 2094-2371, 2137-2635, 2138-2408, 2186-2427, 2285-2698, 2311-2499, 2317-2548, 2367-3019, 2396-3011, 2398-2699, 2398-3213, 2403-3014, 2460-3046, 2546-3044, 2555-2808, 2555-2817, 2566-3177, 2568-3049, 2570-2825, 2665-2881, 2679-3099, 2834-3025, 2834-3282, 2924-3178, 2947-3553, 2996-3550, 3067-3325, 3098-3667, 3200-3704, 3285-3544, 3290-3605, 3291-3454, 3417-3750, 3429-3672, 3541-3830, 3541-3831, 3571-3822, 3571-4122, 3572-3817, 3587-4067, 3671-3938, 3671-3946, 3683-3971, 3740-4007, 3765-4057, 3825-3892, 3836-4044, 3874-4060, 3883-4197, 3906-4150, 3915-4249, 3984-4224, 4008-4291, 4011-4247, 4012-4262, 4030-4321, 4147-4407, 4161-4374
46/7102342CB1/2177	1-745, 6-581, 13-543, 15-510, 21-759, 24-769, 44-772, 194-371, 194-531, 194-578, 194-580, 195-580, 198-268, 198-579, 198-580, 199-580, 204-580, 209-580, 220-580, 255-578, 278-580, 279-580, 650-1213, 650-1252, 650-1310, 650-1328, 650-1355, 902-1145, 902-1453, 902-1508, 1185-1584, 1206-1722, 1219-1616, 1220-1735, 1227-1907, 1229-1657, 1272-1444, 1322-2121, 1342-2177, 1370-1666, 1384-2176, 1404-2177, 1414-2177, 1514-2161, 1542-2177, 1598-2171, 1894-1961, 1909-1961, 1948-2000, 2051-2171, 2055-2171, 2112-2171
47/4169939CB1/703	1-603, 5-300, 109-703, 112-703, 129-417, 132-427
48/6539977CB1/1295	1-448, 1-450, 1-1295, 3-322, 150-455, 156-455, 449-933, 459-778, 459-842, 459-865, 475-721, 475-1050, 865-1295, 1061-1112, 1061-1149, 1061-1186, 1061-1222, 1061-1259, 1097-1295, 1134-1295, 1171-1295, 1207-1295, 1244-1295
49/7675588CB1/575	1-273, 1-492, 1-566, 65-323, 79-566, 89-319, 196-328, 243-575, 262-575, 420-566
50/6244077CB1/1062	1-652, 50-1060, 490-970, 490-986, 490-1018, 490-1019, 490-1062, 493-1050, 514-1021, 624-1049, 647-735, 656-1055, 667-1062, 696-1053, 744-1055, 770-852
51/7498404CB1/1029	1-599, 2-1029
52/7391748CB1/905	1-528, 30-256, 30-309, 246-475, 250-492, 261-697, 265-842, 317-626, 321-583, 325-780, 335-887, 336-618, 360-558, 360-836, 374-606, 378-837, 386-553, 387-618, 388-865, 399-645, 400-658, 418-905, 422-905, 428-888, 429-884, 430-701, 433-659, 433-660, 444-836, 445-889, 446-888, 449-713, 454-579, 456-752, 459-625, 459-884, 464-888, 472-888, 480-710, 482-732

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
53/7499780CB1/2667	1-585, 66-747, 173-512, 173-650, 173-745, 203-450, 215-494, 215-498, 215-742, 215-803, 234-663, 266-841, 288-573, 297-799, 373-1007, 389-924, 445-850, 449-950, 510-791, 512-837, 513-1071, 516-782, 518-652, 520-772, 520-1123, 521-1054, 528-1044, 530-807, 530-1022, 530-1081, 532-703, 535-1022, 536-797, 539-655, 550-1138, 558-1155, 570-860, 572-704, 572-814, 572-829, 589-1087, 591-909, 613-892, 618-822, 622-926, 635-1121, 666-927, 675-782, 784-1015, 784-1028, 801-992, 844-1144, 849-957, 849-1536, 872-1121, 921-1155, 933-1155, 964-1155, 1026-1130, 1155-1278, 1159-1758, 1159-1797, 1162-1698, 1165-1606, 1186-1686, 1196-1372, 1198-1677, 1210-1516, 1210-1731, 1211-1775, 1215-1682, 1215-1793, 1232-1487, 1232-1509, 1260-1924, 1263-1519, 1267-1916, 1273-1502, 1277-1758, 1278-1943, 1282-1573, 1286-1555, 1286-1938, 1291-1925, 1296-1925, 1298-1576, 1300-1553, 1322-1620, 1327-1570, 1329-1847, 1343-1602, 1349-1633, 1352-2038, 1362-1485, 1365-1962, 1371-1929, 1379-2064, 1381-1513, 1382-1641, 1382-1669, 1403-2073, 1405-2004, 1414-2054, 1423-1916, 1427-1937, 1435-1869, 1441-1695, 1453-1908, 1473-1704, 1475-1731, 1476-1978, 1476-2024, 1476-2036, 1476-2056, 1480-2036, 1497-1719, 1497-1782, 1507-1708, 1515-2186, 1517-2215, 1518-2011, 1520-2072, 1534-1809, 1536-2234, 1537-1774, 1543-2102, 1547-2009, 1550-2012, 1551-2147, 1551-2168, 1552-2248, 1563-2005, 1565-1910, 1568-2089, 1574-1832, 1574-2181, 1576-2289, 1578-2282, 1580-2218, 1583-1852, 1587-2126, 1596-2278, 1606-2264, 1608-2108, 1609-1912, 1611-2126, 1614-1767, 1620-2110, 1622-2173, 1624-1872, 1624-1917, 1624-2237, 1630-2179, 1638-2211, 1641-1869, 1642-2291, 1649-2105, 1650-2277, 1653-1787, 1658-1927, 1659-1926, 1667-2182, 1667-2275, 1668-1907, 1668-1943, 1670-1938, 1675-2288, 1679-1952, 1679-2192, 1681-1974, 1682-2363, 1682-2383, 1685-1955, 1685-2379, 1687-1955, 1719-2343, 1719-2363, 1721-2200, 1723-1983, 1723-2005, 1725-2312, 1726-2032, 1726-2279, 1727-2433, 1742-1980, 1764-1981, 1765-1833, 1765-2240, 1765-2355, 1766-2053, 1769-2347, 1773-1834, 1774-2278, 1775-2267, 1776-2078, 1778-2196, 1793-2390, 1799-2388, 1809-2081, 1809-2358, 1821-2012, 1822-2088, 1822-2096, 1823-2416, 1824-2102, 1829-2113, 1835-2355, 1844-2510, 1850-2138, 1850-2170, 1861-2071, 1861-2174, 1870-2152, 1876-2514, 1880-2281, 1880-2297, 1881-2145, 1882-2151, 1884-2489, 1885-2141, 1885-2468, 1888-2121, 1888-2451, 1893-2180, 1895-2416, 1896-2563, 1896-2584, 1898-2618, 1902-2121, 1902-2523, 1904-2113, 1909-2484, 1920-2128, 1920-2559, 1921-2578, 1923-2141, 1923-2550, 1929-2576, 1933-2172, 1953-2244, 1959-2405, 1962-2226, 1963-2326, 1963-2646, 1975-2667, 1981-2635, 1982-2653, 1989-2258, 1989-2583, 1992-2234, 1992-2239, 1992-2248, 2003-2274, 2004-2583, 2005-2651, 2005-2653, 2006-2240, 2007-2266, 2008-2298, 2009-2248, 2015-2347, 2021-2503, 2021-2667, 2022-2360, 2030-2480, 2040-2666, 2040-2667, 2041-2302, 2044-2309, 2044-2410, 2048-2355, 2050-2667, 2051-2172, 2051-2306, 2051-2320,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
	2051-2345, 2051-2355, 2056-2603, 2059-2221, 2060-2621, 2064-2311, 2064-2334, 2064-2365, 2070-2324, 2074-2298, 2088-2661, 2089-2650, 2091-2372, 2091-2667, 2094-2399, 2098-2630, 2100-2347, 2104-2667, 2105-2667, 2108-2667, 2110-2656, 2111-2666, 2122-2556, 2132-2667, 2135-2666, 2141-2653, 2143-2667, 2156-2667, 2160-2664, 2162-2424, 2172-2409, 2172-2663, 2172-2664, 2181-2667, 2187-2440, 2191-2667, 2194-2379, 2195-2667, 2196-2662, 2196-2667, 2197-2457, 2202-2661, 2204-2666, 2207-2662, 2208-2663, 2211-2457, 2211-2660, 2211-2664, 2212-2545, 2212-2667, 2216-2459, 2218-2510, 2218-2653, 2218-2667, 2221-2506, 2221-2663, 2223-2667, 2225-2663, 2227-2666, 2230-2665, 2231-2663, 2231-2667, 2243-2663, 2251-2639, 2251-2662, 2253-2664, 2254-2664, 2260-2666, 2261-2663, 2263-2662, 2269-2662, 2271-2664, 2274-2664, 2277-2666, 2278-2664, 2286-2662, 2288-2664, 2290-2662, 2298-2547, 2305-2665, 2313-2663, 2317-2660, 2336-2573, 2336-2626, 2358-2505
54/7499881CB1/2977	1-258, 1-310, 33-767, 42-258, 42-610, 306-729, 372-941, 407-687, 440-794, 545-713, 795-1350, 1031-1646, 1062-1309, 1088-1337, 1108-1603, 1137-1345, 1161-1432, 1200-1581, 1230-1793, 1239-1499, 1245-1793, 1288-1738, 1348-1793, 1368-1517, 1375-1656, 1380-2242, 1383-1629, 1393-1715, 1415-1631, 1433-1571, 1439-1744, 1461-1671, 1469-1693, 1469-2019, 1478-1665, 1495-1707, 1517-1732, 1520-2009, 1528-1763, 1589-1862, 1606-1910, 1640-1928, 1679-1914, 1681-1929, 1682-2087, 1692-1862, 1727-2227, 1746-2533, 1790-2063, 1795-2120, 1806-2059, 1809-2083, 1827-2533, 1836-2067, 1845-2098, 1870-2160, 1876-2169, 1904-2179, 1911-2533, 1923-2260, 1925-2288, 1940-2200, 1941-2214, 1945-2209, 1947-2533, 1956-2554, 1957-2215, 1957-2535, 1966-2088, 1972-2533, 1975-2312, 1977-2217, 1977-2246, 1983-2254, 1997-2533, 1999-2291, 2033-2320, 2046-2499, 2062-2563, 2073-2687, 2074-2646, 2077-2499, 2079-2341, 2083-2329, 2090-2355, 2093-2420, 2103-2336, 2115-2386, 2118-2533, 2120-2396, 2136-2636, 2179-2910, 2185-2506, 2192-2848, 2199-2895, 2200-2790, 2201-2714, 2214-2488, 2533, 2120-2396, 2136-2636, 2179-2910, 2185-2506, 2192-2848, 2199-2895, 2200-2790, 2201-2714, 2214-2488,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
	2215-2458, 2228-2927, 2236-2732, 2262-2579, 2264-2520, 2285-2552, 2288-2541, 2292-2916, 2304-2964, 2311-2952, 2338-2848, 2358-2598, 2362-2665, 2368-2977, 2372-2927, 2374-2956, 2376-2848, 2397-2655, 2402-2921, 2408-2952, 2415-2977, 2417-2905, 2422-2702, 2422-2952, 2424-2650, 2436-2617, 2443-2947, 2463-2759, 2469-2690, 2474-2977, 2484-2727, 2491-2965, 2494-2748, 2494-2961, 2494-2965, 2496-2970, 2502-2971, 2509-2966, 2509-2972, 2510-2965, 2512-2971, 2513-2778, 2513-2951, 2513-2975, 2514-2965, 2515-2977, 2517-2971, 2518-2764, 2518-2964, 2519-2822, 2519-2969, 2525-2805, 2526-2965, 2527-2780, 2527-2965, 2529-2965, 2533-2959, 2535-2965, 2544-2974, 2546-2809, 2548-2974, 2551-2803, 2551-2969, 2554-2965, 2555-2969, 2558-2971, 2559-2965, 2560-2977, 2561-2962, 2561-2965, 2563-2965, 2564-2965, 2566-2965, 2567-2969, 2568-2965, 2573-2972, 2579-2977, 2598-2965, 2600-2958, 2604-2964, 2604-2965, 2604-2970, 2608-2965, 2609-2965, 2617-2969, 2618-2965, 2640-2959, 2653-2917, 2653-2921, 2657-2965, 2659-2965, 2693-2888, 2699-2965, 2702-2965, 2702-2970, 2703-2965, 2773-2965, 2811-2959, 2858-2977
55/7488579CB1/729	1-729, 23-520, 27-331, 27-357, 27-361, 34-327, 50-337, 83-364, 117-503, 127-405, 130-410, 173-438, 192-481, 218-503, 232-503
56/7510521CB1/1879	1-237, 15-181, 21-241, 21-1869, 23-197, 27-241, 30-241, 35-161, 37-241, 48-200, 61-96, 61-241, 62-241, 66-241, 67-196, 67-240, 70-241, 71-171, 80-231, 112-378, 118-375, 118-386, 268-364, 269-525, 270-808, 271-515, 294-524, 300-565, 305-592, 312-804, 317-894, 317-925, 322-575, 328-911, 330-578, 335-649, 341-597, 341-888, 347-688, 353-933, 356-604, 366-622, 366-795, 367-525, 367-769, 368-614, 377-687, 379-943, 397-637, 400-675, 401-653, 403-669, 403-696, 403-987, 404-715, 405-652, 407-675, 413-732, 428-716, 428-1015, 430-772, 430-998, 433-1043, 436-723, 436-1072, 439-1150, 441-698, 441-704, 444-755, 445-1009, 458-991, 459-918, 461-659, 461-718, 463-713, 465-995, 469-758, 469-1013, 472-1053, 476-781, 481-706, 481-739, 494-904, 496-655, 496-768, 499-764, 500-962, 504-852, 507-821, 519-743, 519-792, 519-823, 527-753, 528-816, 535-817, 536-823, 538-825, 545-1094, 549-797, 551-775, 551-810, 554-839, 556-799, 556-835, 557-780, 560-660, 562-807, 565-783, 567-919, 567-943, 567-970, 586-822, 588-1129, 594-963, 595-856, 595-1316, 596-877, 598-1196, 602-844, 610-807, 612-865,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
	612-873, 617-720, 617-1093, 619-923, 625-867, 625-875, 625-900, 626-890, 627-875, 628-783, 628-903, 628-917, 631-879, 647-886, 660-921, 660-1127, 660-1231, 661-930, 669-920, 669-943, 669-1208, 670-905, 671-907, 672-1285, 673-944, 676-918, 679-925, 685-890, 685-897, 685-936, 688-947, 688-1270, 689-943, 689-960, 689-1257, 696-1013, 699-1333, 703-1168, 705-999, 705-1316, 706-957, 711-999, 711-1263, 720-1072, 724-950, 724-981, 724-1295, 732-1009, 741-1026, 745-987, 745-996, 745-998, 755-990, 769-1041, 769-1042, 778-979, 778-1174, 779-1007, 779-1017, 781-1036, 782-1351, 788-1329, 788-1467, 791-1026, 791-1035, 797-1028, 799-1291, 803-1005, 803-1350, 814-1071, 815-1507, 821-1083, 822-1101, 824-1082, 824-1507, 829-1012, 834-1108, 836-1138, 839-1013, 839-1087, 839-1172, 845-1113, 851-1062, 852-1288, 867-1121, 867-1133, 867-1153, 867-1166, 867-1170, 869-1122, 872-1110, 872-1135, 873-1096, 874-1712, 876-1142, 876-1146, 876-1147, 876-1150, 877-1119, 886-1448, 888-1242, 888-1514, 896-1352.
	898-1161, 901-1110, 902-1286, 904-1149, 905-1155, 908-1177, 908-1334, 909-1258, 909-1355, 910-1134, 912-1712, 914-1103, 914-1436, 924-1165, 925-1357, 926-1204, 927-1165, 928-1034, 935-1337, 937-1081, 937-1140, 937-1150, 937-1154, 941-1136, 944-1140, 946-1595, 954-1286, 957-1225, 957-1229, 958-1712, 959-1151, 959-1240, 959-1378, 966-1195, 966-1201, 966-1235, 967-1712, 970-1269, 977-1131, 981-1537, 985-1401, 993-1632, 995-1254, 995-1543, 999-1261, 1000-1684, 1001-1242, 1001-1465, 1002-1202, 1002-1229, 1002-1248, 1003-1712, 1014-1545, 1016-1262, 1016-1306, 1019-1318, 1020-1280, 1020-1616, 1021-1277, 1022-1267, 1022-1273, 1024-1290, 1025-1263, 1027-1351, 1060-1281, 1068-1356, 1076-1350, 1077-1343, 1077-1732, 1077-1784, 1078-1367, 1084-1352, 1086-1691, 1091-1271, 1091-1340, 1094-1338, 1106-1339, 1106-1449, 1110-1400, 1113-1401, 1113-1408, 1114-1356, 1114-1385, 1125-1371, 1125-1415, 1125-1675, 1129-1378, 1130-1351, 1130-1437, 1132-1375, 1132-1389, 1133-

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
	1635, 1134-1417, 1134-1717, 1137-1800, 1141-1286, 1141-1353, 1141-1412, 1141-1429, 1141-1431, 1147-1627, 1165-1444, 1165-1668, 1172-1467, 1173-1362, 1173-1437, 1173-1580, 1178-1684, 1185-1469, 1187-1477, 1193-1616, 1195-1456, 1198-1487, 1198-1516, 1201-1463, 1201-1662, 1203-1710, 1209-1376, 1209-1398, 1209-1403, 1210-1356, 1210-1840, 1211-1489, 1211-1492, 1216-1524, 1217-1401, 1217-1404, 1217-1470, 1218-1526, 1221-1489, 1222-1511, 1227-1477, 1229-1462, 1229-1513, 1230-1497, 1231-1861, 1232-1500, 1234-1869, 1235-1414, 1235-1493, 1235-1678, 1238-1479, 1238-1492, 1240-1466, 1240-1488, 1240-1496, 1247-1346, 1247-1499, 1247-1516, 1251-1476, 1255-1527, 1259-1869, 1261-1523, 1261-1549, 1262-1505, 1262-1535, 1270-1823, 1274-1524, 1274-1542, 1275-1826, 1281-1762, 1283-1802, 1292-1869, 1293-1556, 1293-1825, 1296-1542, 1296-1570, 1301-1639, 1301-1824, 1305-1500, 1305-1583, 1313-1540, 1313-1578, 1314-1829, 1318-1773, 1332-1828, 1335-1489, 1336-1638, 1340-1865, 1340-1879, 1341-1609, 1341-1828, 1342-1861, 1351-1641, 1352-1869, 1358-1879, 1361-1797, 1362-1633, 1371-1689, 1375-1869, 1377-1869, 1383-1835, 1384-1630, 1386-1869, 1392-1861, 1398-1861, 1401-1869, 1404-1868, 1405-1836, 1409-1869, 1410-1861, 1411-1517, 1411-1869, 1412-1869, 1413-1861, 1417-1869, 1418-1869, 1419-1867, 1420-1867, 1422-1668, 1424-1869, 1424-1879, 1428-1696, 1433-1679, 1433-1879, 1438-1632, 1438-1867, 1440-1868, 1440-1869, 1441-1869, 1443-1596, 1443-1730, 1443-1869, 1444-1869, 1445-1867, 1446-1869, 1447-1861, 1447-1869, 1448-1869, 1450-1711, 1450-1863, 1450-1869, 1451-1861, 1451-1867, 1451-1868, 1451-1869, 1452-1861, 1452-1879, 1453-1820, 1453-1869, 1454-1867, 1456-1835, 1456-1869, 1457-1759, 1457-1869, 1458-1709, 1459-1868, 1460-1867, 1462-1861, 1462-1867, 1463-1868, 1463-1869, 1464-1867, 1465-1689, 1465-1869, 1466-1601, 1466-1712, 1466-1729, 1466-1850, 1466-1861, 1467-1869, 1471-1861, 1471-1867, 1475-1861, 1476-1854, 1477-1868, 1478-1763, 1480-1869, 1481-1834, 1481-1867, 1481-1868, 1483-1861, 1483-1867, 1484-1741, 1486-1737, 1491-1857, 1491-1869, 1492-1879, 1493-1869, 1495-1678, 1496-1756, 1498-1869, 1500-1861, 1501-1869, 1502-1861, 1504-1869, 1505-1759, 1506-1731, 1506-1748, 1507-1869, 1513-1784, 1524-1861, 1524-1879, 1527-1869, 1530-1755, 1531-1862, 1533-1809, 1535-1815, 1541-1710, 1543-1784, 1544-1795, 1552-1826, 1554-1869, 1555-1867, 1556-1861, 1556-1868, 1563-1861, 1564-1854, 1575-1850, 1577-1879, 1586-1869, 1588-1869, 1589-1869, 1590-1868, 1592-1868, 1592-1869, 1593-1861, 1596-1868, 1596-1869, 1598-1861, 1599-1867, 1600-1851, 1600-1869, 1601-1869, 1602-1861, 1604-1879, 1605-1868, 1611-1869, 1612-1861, 1613-1835, 1623-1868, 1623-1869, 1631-1869, 1641-1861, 1642-1869, 1643-1864, 1644-1868, 1645-1861, 1658-1868, 1665-1861, 1666-1861, 1669-1869, 1672-1866, 1674-1869, 1675-1879, 1686-1869, 1716-1879, 1735-1824, 1739-1869, 1744-1879, 1745-1869, 1773-1861, 1795-1861

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID:	Representative Library
29	7994355CB1	ESOGTUT02
30	7475875CB1	ENDCNOT03
31	71231882CB1	BRAZNOT01
32	2875922CB1	PLACFER06
33	8158136CB1	LNODNON02
34	5969491CB1	BEPINON01
35	7497367CB1	LIVRTMR01
36	7632424CB1	PROSNOT11
37	1804436CB1	SINTBST01
39	7472344CB1	OVARDIR01
40	7192959CB1	BRAITUT08
41	6169565CB1	UTRSTDT01
42	7494717CB1	BRSTNOT02
44	7498882CB1	DENDNOT01
45	5524205CB1	BRAUTDR02
46	7102342CB1	BRSTTUT02
47	4169939CB1	PANCNOT19
48	6539977CB1	BRAIFEN08
49	7675588CB1	BRONDI01
50	6244077CB1	TESTNOT17
52	7391748CB1	THPINOT01
53	7499780CB1	PITUNON01
54	7499881CB1	CRBLNOT01
55	7488579CB1	SINTTMR02
56	7510521CB1	DENDNOT01

Table 6

Library	Vector	Library Description
BEPINON01	PSPORT	Normalized library was constructed from 5.12 million independent clones from a bronchial epithelium library. RNA was made from a bronchial epithelium primary cell line derived from a 54-year-old Caucasian male. The normalization and hybridization conditions were adapted from Soares et al., PNAS (1994) 91:9228, using a longer (24-hour) reannealing hybridization period.
BRAIFEN08	pINCY	This normalized fetal brain tissue library was constructed from 400 thousand independent clones from a fetal brain tissue library. Starting RNA was made from brain tissue removed from a Caucasian male fetus who was stillborn with a hypoplastic left heart at 23 weeks' gestation. The library was normalized in 2 rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228 and Bonaldo et al., Genome Research (1996) 6:791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
BRAITUT08	pINCY	Library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 47-year-old Caucasian male during excision of cerebral meningeal tissue. Pathology indicated grade 4 fibrillary astrocytoma with focal tumoral radionecrosis. Patient history included cerebrovascular disease, deficiency anemia, hyperlipidemia, epilepsy, and tobacco use. Family history included cerebrovascular disease and a malignant prostate neoplasm.
BRAUTDR02	PCDNA2.1	This random primed library was constructed using RNA isolated from pooled amygdala and entorhinal cortex tissue removed from a 55-year-old Caucasian female who died from cholangiocarcinoma. Pathology indicated mild meningeal fibrosis predominately over the convexities, scattered axonal spheroids in the white matter of the cingulate cortex and the thalamus, and a few scattered neurofibrillary tangles in the entorhinal cortex and the periaqueductal gray region. Pathology for the associated tumor tissue indicated well-differentiated cholangiocarcinoma of the liver with residual or relapsed tumor. Patient history included cholangiocarcinoma, post-operative Budd-Chiari syndrome, biliary ascites, hydrothorax, dehydration, malnutrition, oliguria and acute renal failure. Previous surgeries included cholecystectomy and resection of 85% of the liver.
BRAZNOT01	pINCY	Library was constructed using RNA isolated from striatum, globus pallidus and posterior putamen tissue removed from a 45-year-old Caucasian female who died from a dissecting aortic aneurysm and ischemic bowel disease. Pathology indicated mild arteriosclerosis involving the cerebral cortical white matter and basal ganglia. Grossly, there was mild meningeal fibrosis and mild focal atherosclerotic plaque in the middle cerebral artery, as well as vertebral arteries bilaterally. Microscopically, the cerebral hemispheres, brain stem and cerebellum reveal focal areas in the white matter that contain blood vessels that were barrel-shaped, hyalinized, with hemosiderin-laden macrophages in the Virchow-Robin space. In addition, there were scattered neurofibrillary tangles within the basolateral nuclei of the amygdala. Patient history included mild atheromatosis of aorta and coronary arteries, bowel and liver infarct due to aneurysm, physiologic fatty liver associated with obesity, mild diffuse emphysema, thrombosis of mesenteric and portal veins, cardiomegaly due to

Table 6

Library	Vector	Library Description
		hypertrophy of left ventricle, arterial hypertension, acute pulmonary edema, splenomegaly, obesity (300 lb.), leiomyoma of uterus, sleep apnea, and iron deficiency anemia.
BRONDIT01	pINCY	Library was constructed using RNA isolated from right lower lobe bronchial tissue removed from a pool of 3 asthmatic Caucasian male and female donors, 22- to 51-years-old during bronchial pinch biopsies. Patient history included atopy as determined by positive skin tests to common aero-allergens.
BRSTNOT02	PSPORT1	Library was constructed using RNA isolated from diseased breast tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated proliferative fibrocystic changes characterized by apocrine metaplasia, sclerosing adenosis, cyst formation, and ductal hyperplasia without atypia. Pathology for the associated tumor tissue indicated an invasive grade 4 mammary adenocarcinoma. Patient history included atrial tachycardia and a benign neoplasm. Family history included cardiovascular and cerebrovascular disease.
BRSTTUT02	PSPORT1	Library was constructed using RNA isolated from breast tumor tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy with reconstruction. Pathology indicated residual invasive grade 3 mammary ductal adenocarcinoma. The remaining breast parenchyma exhibited proliferative fibrocystic changes without atypia. One of 10 axillary lymph nodes had metastatic tumor as a microscopic intranodal focus. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia, and a malignant colon neoplasm.
CRBLNOT01	PSPORT1	Library was constructed using RNA isolated from the cerebellum tissue of a 69-year-old Caucasian male who died from chronic obstructive pulmonary disease. Patient history included myocardial infarction, hypertension, and osteoarthritis.
DENDNOT01	pINCY	Library was constructed using RNA isolated from untreated dendritic cells from peripheral blood.
ENDCNOT03	pINCY	Library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male.
ESOGTUT02	pINCY	Library was constructed using RNA isolated from esophageal tumor tissue obtained from a 61-year-old Caucasian male during a partial esophagectomy, proximal gastrectomy, pyloromyotomy, and regional lymph node excision. Pathology indicated an invasive grade 3 adenocarcinoma in the esophagus. Family history included atherosclerotic coronary artery disease, type II diabetes, chronic liver disease, primary cardiomyopathy, benign hypertension, and cerebrovascular disease.

Table 6

Library	Vector	Library Description
LIVRTMR01	PCDNA2.1	This random primed library was constructed using RNA isolated from liver tissue removed from a 62-year-old Caucasian female during partial hepatectomy and exploratory laparotomy. Pathology for the matched tumor tissue indicated metastatic intermediate grade neuroendocrine carcinoma, consistent with islet cell tumor, forming nodules ranging in size, in the lateral and medial left liver lobe. The pancreas showed fibrosis, chronic inflammation and fat necrosis consistent with pseudocyst. The gallbladder showed mild chronic cholecystitis. Patient history included malignant neoplasm of the pancreas tail, pulmonary embolism, hyperlipidemia, thrombophlebitis, joint pain in multiple joints, type II diabetes, benign hypertension, cerebrovascular disease, and normal delivery. Previous surgeries included distal pancreatectomy, total splenectomy, and partial hepatectomy. Family history included pancreas cancer with secondary liver cancer, benign hypertension, and hyperlipidemia.
LNODNON02	pINCY	This normalized lymph node tissue library was constructed from .56 million independent clones from a lymph node tissue library. Starting RNA was made from lymph node tissue removed from a 16-month-old Caucasian male who died from head trauma. Serologies were negative. Patient history included bronchitis. Patient medications included Dopamine, Dobutamine, Vancomycin, Vasopressin, Proventil, and Atarax. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9932 and Bonaldo et al., Genome Research 6 (1996):791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
OVARDIR01	PCDNA2.1	This random primed library was constructed using RNA isolated from right ovary tissue removed from a 45-year-old Caucasian female during total abdominal hysterectomy, bilateral salpingo-oophorectomy, vaginal suspension and fixation, and incidental appendectomy. Pathology indicated stromal hyperthecosis of the right and left ovaries. Pathology for the matched tumor tissue indicated a dermoid cyst (benign cystic teratoma) in the left ovary. Multiple (3) intramural leiomyomata were identified. The cervix showed squamous metaplasia. Patient history included metrorrhagia, female stress incontinence, alopecia, depressive disorder, pneumonia, normal delivery, and deficiency anemia. Family history included benign hypertension, atherosclerotic coronary artery disease, hyperlipidemia, and primary tuberculous complex.
PANCN0T19	pINCY	Library was constructed using RNA isolated from pancreatic tissue removed from an 8-year-old Black male who died from anoxia.

Table 6

Library	Vector	Library Description
PITUNON01	pINCY	This normalized pituitary gland tissue library was constructed from 6.92 million independent clones from a pituitary gland tissue library. Starting RNA was made from pituitary gland tissue removed from a 55-year-old male who died from chronic obstructive pulmonary disease. Neuropathology indicated there were no gross abnormalities, other than mild ventricular enlargement. There was no apparent microscopic abnormality in any of the neocortical areas examined, except for a number of silver positive neurons with apical dendrite staining, particularly in the frontal lobe. The significance of this was undetermined. The only other microscopic abnormality was that there was prominent silver staining with some swollen axons in the CA3 region of the anterior and posterior hippocampus. Microscopic sections of the cerebellum revealed mild Bergmann's gliosis in the Purkinje cell-layer. Patient history included schizophrenia. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research
PLACFER06	pINCY	This random primed library was constructed using RNA isolated from placental tissue removed from a Caucasian fetus who died after 16 weeks' gestation from fetal demise and hydrocephalus. Patient history included umbilical cord wrapped around the head (3 times) and the shoulders (1 time). Serology was positive for anti-CMV. Family history included multiple pregnancies and live births, and an abortion.
PROSNOT11	pINCY	Library was constructed using RNA isolated from the prostate tissue of a 28-year-old Caucasian male, who died from a self-inflicted gunshot wound.
SINTBST01	pINCY	Library was constructed using RNA isolated from ileum tissue obtained from an 18-year-old Caucasian female during bowel anastomosis. Pathology indicated Crohn's disease of the ileum, involving 15 cm of the small bowel. Family history included cerebrovascular disease and atherosclerotic coronary artery disease.
SINTTMR02	PCDNA2.1	This random primed library was constructed using RNA isolated from small intestine tissue removed from a 59-year-old male. Pathology for the matched tumor tissue indicated multiple (9) carcinoid tumors, grade 1, in the small bowel. The largest tumor was associated with a large mesenteric mass. Multiple convoluted segments of bowel were adhered to the tumor. A single (1 of 13) regional lymph node was positive for malignancy. The peritoneal biopsy indicated focal fat necrosis.
TESTNOT17	pINCY	Library was constructed from testis tissue removed from a 26-year-old Caucasian male who died from head trauma due to a motor vehicle accident. Serologies were negative. Patient history included a hernia at birth, tobacco use (1 1/2 ppd), marijuana use, and daily alcohol use (beer and hard liquor).
THPINOT01	pINCY	Library was constructed using RNA isolated from untreated THP-1 cells. THP-1 is a human promonocyte line derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (ref: Int. J. Cancer (1980) 26:171).
UTRSTDT01	pINCY	Library was constructed using RNA isolated from uterus tissue removed from a 46-year-old Caucasian female who died from cardiopulmonary arrest. Patient history included liver and breast cancer.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value = 1.0E-8 or less; Full Length sequences: Probability value = 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value = 1.0E-6; Assembled ESTs: fasta Identity = 95% or greater and Match length = 200 bases or greater; fastx E value = 1.0E-8 or less; Full Length sequences: fastx score = 100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-426.	Probability value = 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM, INCY, SMART and TIGRFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-10.	PFAM, INCY, SMART or TIGRFAM hits: Probability value = 1.0E-3 or less; Signal peptide hits: Score = 0 or greater
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score \geq GCG specified "HIGH" value for that particular Prosite motif. Generally, score = 1.4-2.1.

Table 7

Program	Description	Reference	Parameter Threshold
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score = 120 or greater; Match length = 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score = 3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. On Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence (AAAI) Press, Menlo Park, CA, and MIT Press, Cambridge, MA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
56	7510521	064728H1	SNP00006579	124	526	T	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	073306H1	SNP00105444	119	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	073566H1	SNP00105444	51	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	073573H1	SNP00105444	51	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	074430H1	SNP00105444	119	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	075234H1	SNP00105444	119	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	075593H1	SNP00105444	43	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	076131H1	SNP00105444	119	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	076800H1	SNP00105444	119	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	076808H1	SNP00105444	119	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	078964H1	SNP00105444	119	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	123967H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	136239H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	139719H1	SNP00067426	37	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	142453H1	SNP00105444	235	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	145280H1	SNP00105444	166	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	145328H1	SNP00067426	28	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	145480H1	SNP00067426	28	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	151495H1	SNP00067426	28	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	156942R1	SNP00105444	457	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	1636617H1	SNP00105444	43	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	163725H1	SNP00105444	43	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	1806536H1	SNP00105444	243	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	1806750H1	SNP00006580	47	1241	G	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	1880982H1	SNP00105444	131	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	1900090H1	SNP00006579	197	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	1974452H1	SNP00105444	148	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	1979551H1	SNP00105444	95	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	1979551R6	SNP00105444	95	1262	C	C	T	noncoding	n/a	n/a	n/a	n/a

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
56	7510521	2107073H1	SNP00067426	8	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2116360H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2135075H1	SNP00067426	70	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2137253H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2183517H1	SNP00067426	13	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2211782H1	SNP00067426	28	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2212924H1	SNP00067426	28	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2303437H1	SNP00105444	25	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2311978H1	SNP00105444	20	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2369247H1	SNP00006579	8	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	2397207H1	SNP00067426	28	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2398368H1	SNP00006579	161	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	2436959H1	SNP00006579	130	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	2437672H1	SNP00067426	35	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2438148H1	SNP00067426	32	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2454291H1	SNP00006579	46	526	T	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	2582571H1	SNP00067426	35	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2591249H2	SNP00105444	13	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2591549H1	SNP00105444	13	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2620592H1	SNP00067426	29	97	C	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2621494H1	SNP00105444	169	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2622226H1	SNP00006580	226	1241	T	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	2675269H1	SNP00067426	27	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2686334H1	SNP00105444	31	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2722514H1	SNP00067426	50	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2741742H1	SNP00105444	238	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2745948H1	SNP00067426	15	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2767239H1	SNP00067426	26	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2799215H1	SNP00105444	13	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
56	7510521	2848443H1	SNP00067426	30	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2862235H1	SNP00105444	39	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	2884679H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2910110H1	SNP00067426	16	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2913667H1	SNP00067426	33	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2969030H1	SNP00067426	10	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	2994484H1	SNP00006579	20	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3045590H1	SNP00006579	28	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3095345H1	SNP00006579	120	526	T	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3111703H1	SNP00067426	26	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3138158H1	SNP00105444	144	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3220920H1	SNP00105444	42	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3236768H1	SNP00105444	19	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3240887H1	SNP00105444	143	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3283818H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3343887H1	SNP00006579	159	526	T	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3345625H1	SNP00105444	236	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3377028H1	SNP00105444	33	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3402507H1	SNP00067426	61	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3403476H1	SNP00006579	121	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3405513H1	SNP00067426	30	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3411758H1	SNP00067426	33	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3422192H1	SNP00067426	25	97	C	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3450266H1	SNP00006579	205	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3455078H1	SNP00006579	36	526	T	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3473335H1	SNP00105444	126	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3503020H1	SNP00067426	61	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3510182H1	SNP00067426	33	97	C	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3556979H1	SNP00105444	144	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
56	7510521	3595275H1	SNP00067426	10	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3600404H1	SNP00105444	87	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3617096H1	SNP00067426	11	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3628252H1	SNP00105444	132	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3814147H1	SNP00006580	172	1241	T	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	3814616H1	SNP00006579	81	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3833454H1	SNP00105444	60	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3836524H1	SNP00006580	221	1241	T	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	3837242H1	SNP00006579	124	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3841051H1	SNP00105444	62	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3842154H1	SNP00105444	72	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3843327H1	SNP00006579	126	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3843721H1	SNP00105444	74	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3844448H1	SNP00006579	58	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3844674H1	SNP00006579	99	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	3845294H1	SNP00105444	127	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	3845517H1	SNP00067426	31	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	3877879H1	SNP00006579	122	526	T	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	4044544H1	SNP00105444	154	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	4054975H1	SNP00006580	162	1241	G	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	4083278H1	SNP00006580	107	1241	G	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	415116R6	SNP00006579	97	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	4209957H1	SNP00006579	123	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	4335846H1	SNP00105444	48	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	4335847H1	SNP00105444	46	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	4415058H1	SNP00067426	30	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	4424212H1	SNP00006579	171	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	4543723H1	SNP00105444	57	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	4551410H1	SNP00105444	23	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
56	7510521	4557539H1	SNP00105444	18	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	4559026H1	SNP00105444	18	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	4610804H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	4637077H1	SNP00067426	27	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	4671661H1	SNP00067426	30	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	4705376H1	SNP00067426	29	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	4781251H1	SNP00067426	11	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	4794967H1	SNP00006580	244	1241	G	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	4894666H1	SNP00006579	29	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	4993290H1	SNP00006579	257	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	5038495H1	SNP00006579	227	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	5075161H1	SNP00067426	32	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	5111563H1	SNP00006580	156	1241	T	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	5115634H1	SNP00105444	5	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	5187513H1	SNP00105444	170	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	5274442H1	SNP00105444	128	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	5313705H1	SNP00105444	43	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	548863H1	SNP00067426	30	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	5551463H1	SNP00105444	30	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	5610627H1	SNP00105444	184	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	5636114H1	SNP00006579	8	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	5679473H1	SNP00067426	31	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	5685751H1	SNP00067426	44	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	5733518H1	SNP00105444	38	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	5955250H1	SNP00006579	21	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	6028821H1	SNP00105444	22	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	6096755H1	SNP00105444	148	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	6109371H1	SNP00006579	150	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	6116936H1	SNP00067426	32	97	T	T	C	noncoding	0.95	0.95	n/d	0.96

Table 8

SEQ ID NO.	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
56	7510521	6325544H1	SNP00006579	171	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	6325576H1	SNP00006579	177	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	641917H1	SNP00006579	67	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	6458184H1	SNP00006579	91	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	652597H1	SNP00067426	28	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	6529484H1	SNP00006579	255	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	6532850H1	SNP00105444	415	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	6869191H1	SNP00105444	123	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	6916821H1	SNP00006579	170	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	6966242H1	SNP00006579	338	526	C	C	T	noncoding	0.82	n/a	n/a	n/a
56	7510521	7185105H1	SNP00105444	117	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	7218669H1	SNP00105444	314	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	7332779H1	SNP00105444	351	1224	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	7365079H1	SNP00067426	36	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	7370713H1	SNP00006580	557	1241	T	T	G	noncoding	n/a	n/a	n/a	n/a
56	7510521	7628538J1	SNP00105444	459	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	7632152H1	SNP00105444	561	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	766430H1	SNP00105444	135	1259	C	C	T	noncoding	n/a	n/a	n/a	n/a
56	7510521	777571H1	SNP00067426	12	97	T	T	C	noncoding	0.95	0.95	n/d	0.96
56	7510521	7936627H1	SNP00006579	71	526	C	C	T	noncoding	0.82	n/a	n/a	n/a

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28,
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3-7, SEQ ID NO:9-19, SEQ ID NO:21-26, and SEQ ID NO:28,
 - c) a polypeptide comprising a naturally occurring amino acid sequence at least 94% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:20,
 - d) a polypeptide comprising a naturally occurring amino acid sequence at least 96% identical to an amino acid sequence of SEQ ID NO:8,
 - e) a polypeptide comprising a naturally occurring amino acid sequence at least 97% identical to an amino acid sequence of SEQ ID NO:27,
 - f) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, and
 - g) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-28.
2. An isolated polypeptide of claim 1 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28.
3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
5. An isolated polynucleotide of claim 4 comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
9. A method of producing a polypeptide of claim 1, the method comprising:
- 5 a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
- b) recovering the polypeptide so expressed.
- 10 10. A method of claim 9, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-28.
11. An isolated antibody which specifically binds to a polypeptide of claim 1.
- 15 12. An isolated polynucleotide selected from the group consisting of:
- a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:29-56,
- b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of
- 20 SEQ ID NO:29-53 and SEQ ID NO:55-56,
- c) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 91% identical to a polynucleotide sequence of SEQ ID NO:54,
- d) a polynucleotide complementary to a polynucleotide of a),
- e) a polynucleotide complementary to a polynucleotide of b),
- 25 f) a polynucleotide complementary to a polynucleotide of c), and
- g) an RNA equivalent of a)-f).
13. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 12.
- 30 14. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:
- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample,
- 35 and which probe specifically hybridizes to said target polynucleotide, under

conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

5

15. A method of claim 14, wherein the probe comprises at least 60 contiguous nucleotides.

16. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:

- 10 a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

15 17. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

18. A composition of claim 17, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-28.

20

19. A method for treating a disease or condition associated with decreased expression of functional PMOD, comprising administering to a patient in need of such treatment the composition of claim 17.

25 20. A method of screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

30 21. A composition comprising an agonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with decreased expression of functional PMOD, comprising administering to a patient in need of such treatment a composition of claim 21.

35

23. A method of screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

5

24. A composition comprising an antagonist compound identified by a method of claim 23 and a pharmaceutically acceptable excipient.

25. A method for treating a disease or condition associated with overexpression of functional PMOD, comprising administering to a patient in need of such treatment a composition of claim 24.

10

26. A method of screening for a compound that specifically binds to the polypeptide of claim 1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

15

27. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, the method comprising:

20

- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

25

30

28. A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,

35

- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

5 29. A method of assessing toxicity of a test compound, the method comprising:

- a) treating a biological sample containing nucleic acids with the test compound,
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 12 under conditions whereby a specific hybridization complex is formed between said probe and a target
10 polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 12 or fragment thereof,
- c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a
15 difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

30. A diagnostic test for a condition or disease associated with the expression of PMOD in a biological sample, the method comprising:

- 20 a) combining the biological sample with an antibody of claim 11, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex, and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

25

31. The antibody of claim 11, wherein the antibody is:

- a) a chimeric antibody,
- b) a single chain antibody,
- c) a Fab fragment,
- 30 d) a F(ab')₂ fragment, or
- e) a humanized antibody.

35

32. A composition comprising an antibody of claim 11 and an acceptable excipient.

33. A method of diagnosing a condition or disease associated with the expression of PMOD

in a subject, comprising administering to said subject an effective amount of the composition of claim 32.

34. A composition of claim 32, wherein the antibody is labeled.

35. A method of diagnosing a condition or disease associated with the expression of PMOD in a subject, comprising administering to said subject an effective amount of the composition of claim 34.

36. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 11, the method comprising:

- a) immunizing an animal with a polypeptide consisting of an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- b) isolating antibodies from said animal, and
- c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28.

37. A polyclonal antibody produced by a method of claim 36.

38. A composition comprising the polyclonal antibody of claim 37 and a suitable carrier.

39. A method of making a monoclonal antibody with the specificity of the antibody of claim 11, the method comprising:

- a) immunizing an animal with a polypeptide consisting of an amino acid sequence selected from the group consisting of SEQ ID NO:1-28, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- b) isolating antibody producing cells from the animal,
- c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells,
- d) culturing the hybridoma cells, and
- e) isolating from the culture monoclonal antibody which specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28.

40. A monoclonal antibody produced by a method of claim 39.

41. A composition comprising the monoclonal antibody of claim 40 and a suitable carrier.

5 42. The antibody of claim 11, wherein the antibody is produced by screening a Fab expression library.

43. The antibody of claim 11, wherein the antibody is produced by screening a recombinant immunoglobulin library.

10

44. A method of detecting a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28 in a sample, the method comprising:

- a) incubating the antibody of claim 11 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- 15 b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28 in the sample.

45. A method of purifying a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-28 from a sample, the method comprising:

- a) incubating the antibody of claim 11 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) separating the antibody from the sample and obtaining the purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID
- 25 NO:1-28.

46. A microarray wherein at least one element of the microarray is a polynucleotide of claim 13.

30 47. A method of generating an expression profile of a sample which contains polynucleotides, the method comprising:

- a) labeling the polynucleotides of the sample,
- b) contacting the elements of the microarray of claim 46 with the labeled polynucleotides of the sample under conditions suitable for the formation of a
- 35 hybridization complex, and

- c) quantifying the expression of the polynucleotides in the sample.

48. An array comprising different nucleotide molecules affixed in distinct physical locations on a solid substrate, wherein at least one of said nucleotide molecules comprises a first
5 oligonucleotide or polynucleotide sequence specifically hybridizable with at least 30 contiguous nucleotides of a target polynucleotide, and wherein said target polynucleotide is a polynucleotide of claim 12.

49. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is
10 completely complementary to at least 30 contiguous nucleotides of said target polynucleotide.

50. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 60 contiguous nucleotides of said target polynucleotide.

51. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is
15 completely complementary to said target polynucleotide.

52. An array of claim 48, which is a microarray.

53. An array of claim 48, further comprising said target polynucleotide hybridized to a
20 nucleotide molecule comprising said first oligonucleotide or polynucleotide sequence.

54. An array of claim 48, wherein a linker joins at least one of said nucleotide molecules to said solid substrate.

55. An array of claim 48, wherein each distinct physical location on the substrate contains multiple nucleotide molecules, and the multiple nucleotide molecules at any single distinct physical location have the same sequence, and each distinct physical location on the substrate contains nucleotide molecules having a sequence which differs from the sequence of nucleotide molecules at
30 another distinct physical location on the substrate.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

58. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.
59. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.
- 5 60. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.
61. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.
62. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.
- 10 63. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.
64. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.
- 15 65. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.
66. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:11.
67. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:12.
- 20 68. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:13.
69. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:14.
- 25 70. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:15.
71. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:16.
72. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:17.
- 30 73. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:18.
74. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:19.
- 35 75. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:20.

76. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:21.
77. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:22.
- 5 78. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:23.
79. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:24.
80. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:25.
- 10 81. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:26.
82. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:27.
- 15 83. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:28.
84. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:29.
- 20 85. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:30.
86. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:31.
- 25 87. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:32.
88. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:33.
- 30 89. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:34.
- 35 90. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:35.

91. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:36.

5

92. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:37.

93. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

10 NO:38.

94. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:39.

95. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

15

NO:40.

96. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:41.

20

97. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:42.

98. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

25

NO:43.

99. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:44.

100. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

30

NO:45.

101. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID

NO:46.

35

102. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:47.

103. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
5 NO:48.

104. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:49.

105. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
10 NO:50.

106. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:51.

15 107. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:52.

108. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
20 NO:53.

109. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:54.

25 110. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:55.

111. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:56.

30

<110> INCYTE GENOMICS, INC.
GANDHI, Aameena R.
KABLE, Amy E.
SWARNAKAR, Anita
HAFALIA, April J.A.
TRAN, Bao
DUGGAN, Brendan M.
WARREN, Bridget A.
ISON, Craig H.
HONCHELL, Cynthia D.
NGUYEN, Danniel B.
LU, Dyung Aina M.
LEE, Ernestine A.
YUE, Henry
FORSYTHE, Ian J.
BARROSO, Ines
RAMKUMAR, Jayalaxmi
GRIFFIN, Jennifer A.
LI, Joana X.
YANG, Junming
THANGAVELU, Kavitha
GIETZEN, Kimberly J.
DING, Li
BAUGHN, Mariah R.
BOROWSKY, Mark L.
YAO, Monique G.
WALIA, Narinder K.
MASON, Patricia M.
GURURAJAN, Rajagopal
LEE, Sally
BECHA, Shanya D.
LEE, Soo Yeun
TRAN, Uyen K.
ELLIOTT, Vicki S.
LUO, Wen
SPRAGUE, William
TANG, Y. Tom
LU, Yan
ZEBARJADIAN, Yeganeh

<120> PROTEIN MODIFICATION AND MAINTENANCE MOLECULES

<130> PF-1040 PCT

<140> To Be Assigned

<141> Herewith

<150> US 60/300,508

<151> 2001-06-22

<150> US 60/303,445

<151> 2001-07-06

<150> US 60/305,405

<151> 2001-07-13

<150> US 60/311,442

<151> 2001-08-09

<150> US 60/314,821

<151> 2001-08-24

<150> US 60/315,992

<151> 2001-08-29

<150> US 60/378,205

<151> 2002-05-03

<160> 56

<170> PERL Program

<210> 1

<211> 774

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7994355CD1

<400> 1

```

Met Ala Asp Gln His Arg Ser Val Ser Glu Leu Leu Ser Asn Ser
 1          5          10          15
Lys Phe Asp Val Asn Tyr Ala Phe Gly Arg Val Lys Arg Ser Leu
          20          25          30
Leu His Ile Ala Ala Asn Cys Gly Ser Val Glu Cys Leu Val Leu
          35          40          45
Leu Leu Lys Lys Gly Ala Asn Pro Asn Tyr Gln Asp Ile Ser Gly
          50          55          60
Cys Thr Pro Leu His Leu Ala Ala Arg Asn Gly His Gly Gln Arg
          65          70          75
Asp Thr Ala Gln Ile Leu Leu Leu Arg Gly Ala Lys Tyr Leu Pro
          80          85          90
Asp Lys Asn Gly Val Thr Pro Leu Asp Leu Cys Val Gln Gly Gly
          95          100          105
Tyr Gly Glu Thr Cys Glu Val Leu Ile Gln Tyr His Pro Arg Leu
          110          115          120
Phe Gln Thr Ile Ile Gln Met Thr Gln Asn Glu Asp Leu Arg Glu
          125          130          135
Asn Met Leu Arg Gln Val Leu Glu His Leu Ser Gln Gln Ser Glu
          140          145          150
Ser Gln Tyr Leu Lys Ile Leu Thr Ser Leu Ala Glu Val Ala Thr
          155          160          165
Thr Asn Gly His Lys Leu Leu Ser Leu Ser Ser Asn Tyr Asp Ala
          170          175          180
Gln Met Lys Ser Leu Leu Arg Ile Val Arg Met Phe Cys His Val
          185          190          195
Phe Arg Ile Gly Pro Ser Ser Pro Ser Asn Gly Ile Asp Met Gly
          200          205          210
Tyr Asn Gly Asn Lys Thr Pro Arg Ser Gln Val Phe Lys Pro Leu
          215          220          225
Glu Leu Leu Trp His Ser Leu Asp Glu Trp Leu Val Leu Ile Ala
          230          235          240
Thr Glu Leu Met Lys Asn Lys Arg Asp Ser Thr Glu Ile Thr Ser
          245          250          255
Ile Leu Leu Lys Gln Lys Gly Gln Asp Gln Asp Ala Ala Ser Ile
          260          265          270
Pro Pro Phe Glu Pro Pro Gly Pro Gly Ser Tyr Glu Asn Leu Ser
          275          280          285
Thr Gly Thr Arg Glu Ser Lys Pro Asp Ala Leu Ala Gly Arg Gln
          290          295          300
Glu Ala Ser Ala Asp Cys Gln Asp Val Ile Ser Met Thr Ala Asn
          305          310          315
Arg Leu Ser Ala Val Ile Gln Ala Phe Tyr Met Cys Cys Ser Cys

```

	320		325		330
Gln Met Pro Pro	Gly Met Thr Ser Pro	Arg Phe Ile Glu Phe	Val		
	335		340		345
Cys Lys His Asp	Glu Val Leu Lys Cys	Phe Val Asn Arg Asn	Pro		
	350		355		360
Lys Ile Ile Phe	Asp His Phe His Phe	Leu Leu Glu Cys Pro	Glu		
	365		370		375
Leu Met Ser Arg	Phe Met His Ile Ile	Lys Ala Gln Pro Phe	Lys		
	380		385		390
Asp Arg Cys Glu	Trp Phe Tyr Glu His	Leu His Ser Gly Gln	Pro		
	395		400		405
Asp Ser Asp Met	Val His Arg Pro Val	Asn Glu Asn Asp Ile	Leu		
	410		415		420
Leu Val His Arg	Asp Ser Ile Phe Arg	Ser Ser Cys Glu Val	Val		
	425		430		435
Ser Lys Ala Asn	Cys Ala Lys Leu Lys	Gln Gly Ile Ala Val	Arg		
	440		445		450
Phe His Gly Glu	Glu Gly Met Gly Gln	Gly Val Val Arg Glu	Trp		
	455		460		465
Phe Asp Ile Leu	Ser Asn Glu Ile Val	Asn Pro Asp Tyr Ala	Leu		
	470		475		480
Phe Thr Gln Ser	Ala Asp Gly Thr Thr	Phe Gln Pro Asn Ser	Asn		
	485		490		495
Ser Tyr Val Asn	Pro Asp His Leu Asn	Tyr Phe Arg Phe Ala	Gly		
	500		505		510
Gln Ile Leu Gly	Leu Ala Leu Asn His	Arg Gln Leu Val Asn	Ile		
	515		520		525
Tyr Phe Thr Arg	Ser Phe Tyr Lys His	Ile Leu Gly Ile Pro	Val		
	530		535		540
Asn Tyr Gln Asp	Val Ala Ser Ile Asp	Pro Glu Tyr Ala Lys	Asn		
	545		550		555
Leu Gln Trp Ile	Leu Asp Asn Asp Ile	Ser Asp Leu Gly Leu	Glu		
	560		565		570
Leu Thr Phe Ser	Val Glu Thr Asp Val	Phe Gly Ala Met Glu	Glu		
	575		580		585
Val Pro Leu Lys	Pro Gly Gly Gly Ser	Ile Leu Val Thr Gln	Asn		
	590		595		600
Asn Lys Ala Glu	Tyr Val Gln Leu Val	Thr Glu Leu Arg Met	Thr		
	605		610		615
Arg Ala Ile Gln	Pro Gln Ile Asn Ala	Phe Leu Gln Gly Phe	His		
	620		625		630
Met Phe Ile Pro	Pro Ser Leu Ile Gln	Leu Phe Asp Glu Tyr	Glu		
	635		640		645
Leu Glu Leu Leu	Leu Ser Gly Met Pro	Glu Ile Asp Val Ser	Asp		
	650		655		660
Trp Ile Lys Asn	Thr Glu Tyr Thr Ser	Gly Tyr Glu Arg Glu	Asp		
	665		670		675
Pro Val Ile Gln	Trp Phe Trp Glu Val	Val Glu Asp Ile Thr	Gln		
	680		685		690
Glu Glu Arg Val	Leu Leu Leu Gln Phe	Val Thr Gly Ser Ser	Arg		
	695		700		705
Val Pro His Gly	Gly Phe Ala Asn Ile	Met Gly Gly Ser Gly	Leu		
	710		715		720
Gln Asn Phe Thr	Ile Ala Ala Val Pro	Tyr Thr Pro Asn Leu	Leu		
	725		730		735
Pro Thr Ser Ser	Thr Cys Ile Asn Met	Leu Lys Leu Pro Glu	Tyr		
	740		745		750
Pro Ser Lys Glu	Ile Leu Lys Asp Arg	Leu Leu Val Ala Leu	His		
	755		760		765
Cys Gly Ser Tyr	Gly Tyr Thr Met Ala				
	770				

<210> 2

<211> 703
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7475875CD1

<400> 2

Met	Ala	Cys	Ser	Met	Ala	Cys	Gly	Gly	Arg	Ala	Cys	Lys	Tyr	Glu	1	5	10	15
Asn	Pro	Ala	Arg	Trp	Ser	Glu	Gln	Glu	Ala	Ile	Lys	Gly	Val		20	25	30	
Tyr	Ser	Ser	Trp	Val	Thr	Asp	Asn	Ile	Leu	Ala	Met	Ala	Arg	Pro	35	40	45	
Ser	Ser	Glu	Leu	Leu	Glu	Lys	Tyr	His	Ile	Ile	Asp	Gln	Phe	Leu	50	55	60	
Ser	His	Gly	Ile	Lys	Thr	Ile	Ile	Asn	Leu	Gln	Arg	Pro	Gly	Glu	65	70	75	
His	Ala	Ser	Cys	Gly	Asn	Pro	Leu	Glu	Gln	Glu	Ser	Gly	Phe	Thr	80	85	90	
Tyr	Leu	Pro	Glu	Ala	Phe	Met	Glu	Ala	Gly	Ile	Tyr	Phe	Tyr	Asn	95	100	105	
Phe	Gly	Trp	Lys	Asp	Tyr	Gly	Val	Ala	Ser	Leu	Thr	Thr	Ile	Leu	110	115	120	
Asp	Met	Val	Lys	Val	Met	Thr	Phe	Ala	Leu	Gln	Glu	Gly	Lys	Val	125	130	135	
Ala	Ile	His	Cys	His	Ala	Gly	Leu	Gly	Arg	Thr	Gly	Val	Leu	Ile	140	145	150	
Ala	Cys	Tyr	Leu	Val	Phe	Ala	Thr	Arg	Met	Thr	Ala	Asp	Gln	Ala	155	160	165	
Ile	Ile	Phe	Val	Arg	Ala	Lys	Arg	Pro	Asn	Ser	Ile	Gln	Thr	Arg	170	175	180	
Gly	Gln	Leu	Leu	Cys	Val	Arg	Glu	Phe	Thr	Gln	Phe	Leu	Thr	Pro	185	190	195	
Leu	Arg	Asn	Ile	Phe	Ser	Cys	Cys	Asp	Pro	Lys	Ala	His	Ala	Val	200	205	210	
Thr	Leu	Pro	Gln	Tyr	Leu	Ile	Arg	Gln	Arg	His	Leu	Leu	His	Gly	215	220	225	
Tyr	Glu	Ala	Arg	Glu	Leu	Lys	His	Val	Pro	Lys	Ile	Ile	His	Leu	230	235	240	
Val	Cys	Lys	Leu	Leu	Leu	Asp	Leu	Ala	Glu	Asn	Arg	Pro	Val	Met	245	250	255	
Met	Lys	Asp	Val	Ser	Glu	Gly	Pro	Gly	Leu	Ser	Ala	Glu	Ile	Glu	260	265	270	
Lys	Thr	Met	Ser	Glu	Met	Val	Thr	Met	Gln	Leu	Asp	Lys	Glu	Leu	275	280	285	
Leu	Arg	His	Asp	Ser	Asp	Val	Ser	Asn	Pro	Pro	Asn	Pro	Thr	Ala	290	295	300	
Val	Ala	Ala	Asp	Phe	Asp	Asn	Arg	Gly	Met	Ile	Phe	Ser	Asn	Glu	305	310	315	
Gln	Gln	Phe	Asp	Pro	Leu	Trp	Lys	Arg	Arg	Asn	Val	Glu	Cys	Leu	320	325	330	
Gln	Pro	Leu	Thr	His	Leu	Lys	Arg	Arg	Leu	Ser	Tyr	Ser	Asp	Ser	335	340	345	
Asp	Leu	Lys	Arg	Ala	Glu	Asn	Leu	Leu	Glu	Gln	Gly	Glu	Thr	Pro	350	355	360	
Gln	Thr	Val	Pro	Ala	Gln	Ile	Leu	Val	Gly	His	Lys	Pro	Arg	Gln	365	370	375	
Gln	Lys	Leu	Ile	Ser	His	Cys	Tyr	Ile	Pro	Gln	Ser	Pro	Glu	Pro	380	385	390	
Asp	Leu	His	Lys	Glu	Ala	Leu	Val	Arg	Ser	Thr	Leu	Ser	Phe	Trp	395	400	405	

Ser	Gln	Ser	Lys	Phe	Gly	Gly	Leu	Glu	Gly	Leu	Lys	Asp	Asn	Gly		
				410					415					420		
Ser	Pro	Ile	Phe	His	Gly	Lys	Ile	Ile	Pro	Lys	Glu	Ala	Gln	Gln		
				425					430					435		
Ser	Gly	Ala	Phe	Ser	Ala	Asp	Val	Ser	Gly	Ser	His	Ser	Pro	Gly		
				440					445					450		
Glu	Pro	Val	Ser	Pro	Ser	Phe	Ala	Asn	Val	His	Lys	Asp	Pro	Asn		
				455					460					465		
Pro	Ala	His	Gln	Gln	Val	Ser	His	Cys	Gln	Cys	Lys	Thr	His	Gly		
				470					475					480		
Val	Gly	Ser	Pro	Gly	Ser	Val	Arg	Gln	Asn	Ser	Arg	Thr	Pro	Arg		
				485					490					495		
Ser	Pro	Leu	Asp	Cys	Gly	Ser	Ser	Pro	Lys	Ala	Gln	Phe	Leu	Val		
				500					505					510		
Glu	His	Glu	Thr	Gln	Asp	Ser	Lys	Asp	Leu	Ser	Glu	Ala	Ala	Ser		
				515					520					525		
His	Ser	Ala	Leu	Gln	Ser	Glu	Leu	Ser	Ala	Glu	Ala	Arg	Arg	Ile		
				530					535					540		
Leu	Ala	Ala	Lys	Ala	Leu	Ala	Asn	Leu	Asn	Glu	Ser	Val	Glu	Lys		
				545					550					555		
Glu	Glu	Leu	Lys	Arg	Lys	Val	Glu	Met	Trp	Gln	Lys	Glu	Leu	Asn		
				560					565					570		
Ser	Arg	Asp	Gly	Ala	Trp	Glu	Arg	Ile	Cys	Gly	Glu	Arg	Asp	Pro		
				575					580					585		
Phe	Ile	Leu	Cys	Ser	Leu	Met	Trp	Ser	Trp	Val	Glu	Gln	Leu	Lys		
				590					595					600		
Glu	Pro	Val	Ile	Thr	Lys	Glu	Asp	Val	Asp	Met	Leu	Val	Asp	Arg		
				605					610					615		
Arg	Ala	Asp	Ala	Ala	Glu	Ala	Leu	Phe	Leu	Leu	Glu	Lys	Gly	Gln		
				620					625					630		
His	Gln	Thr	Ile	Leu	Cys	Val	Leu	His	Cys	Ile	Val	Asn	Leu	Gln		
				635					640					645		
Thr	Ile	Pro	Val	Asp	Val	Glu	Glu	Ala	Phe	Leu	Ala	His	Ala	Ile		
				650					655					660		
Lys	Ala	Phe	Thr	Lys	Val	Asn	Phe	Asp	Ser	Glu	Asn	Gly	Pro	Thr		
				665					670					675		
Val	Tyr	Asn	Thr	Leu	Lys	Lys	Ile	Phe	Lys	His	Thr	Leu	Glu	Glu		
				680					685					690		
Lys	Arg	Lys	Met	Thr	Lys	Asp	Gly	Pro	Lys	Pro	Gly	Leu				
				695					700							

<210> 3

<211> 1256

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 71231882CD1

<400> 3

Met	Phe	Gly	Asp	Leu	Phe	Glu	Glu	Glu	Tyr	Ser	Thr	Val	Ser	Asn		
1				5					10					15		
Asn	Gln	Tyr	Gly	Lys	Gly	Lys	Lys	Leu	Lys	Thr	Lys	Ala	Leu	Glu		
				20					25					30		
Pro	Pro	Ala	Pro	Arg	Glu	Phe	Thr	Asn	Leu	Ser	Gly	Ile	Arg	Asn		
				35					40					45		
Gln	Gly	Gly	Thr	Cys	Tyr	Leu	Asn	Ser	Leu	Leu	Gln	Thr	Leu	His		
				50					55					60		
Phe	Thr	Pro	Glu	Phe	Arg	Glu	Ala	Leu	Phe	Ser	Leu	Gly	Pro	Glu		
				65					70					75		
Glu	Leu	Gly	Leu	Phe	Glu	Asp	Lys	Asp	Lys	Pro	Asp	Ala	Lys	Val		
				80					85					90		

Arg	Ile	Ile	Pro	Leu	Gln	Leu	Gln	Arg	Leu	Phe	Ala	Gln	Leu	Leu	
				95					100					105	
Leu	Leu	Asp	Gln	Glu	Ala	Ala	Ser	Thr	Ala	Asp	Leu	Thr	Asp	Ser	
				110					115					120	
Phe	Gly	Trp	Thr	Ser	Asn	Glu	Glu	Met	Arg	Gln	His	Asp	Val	Gln	
				125					130					135	
Glu	Leu	Asn	Arg	Ile	Leu	Phe	Ser	Ala	Leu	Glu	Thr	Ser	Leu	Val	
				140					145					150	
Gly	Thr	Ser	Gly	His	Asp	Leu	Ile	Tyr	Arg	Leu	Tyr	His	Gly	Thr	
				155					160					165	
Ile	Val	Asn	Gln	Ile	Val	Cys	Lys	Glu	Cys	Lys	Asn	Val	Ser	Glu	
				170					175					180	
Arg	Gln	Glu	Asp	Phe	Leu	Asp	Leu	Thr	Val	Ala	Val	Lys	Asn	Val	
				185					190					195	
Ser	Gly	Leu	Glu	Asp	Ala	Leu	Trp	Asn	Met	Tyr	Val	Glu	Glu	Glu	
				200					205					210	
Val	Phe	Asp	Cys	Asp	Asn	Leu	Tyr	His	Cys	Gly	Thr	Cys	Asp	Arg	
				215					220					225	
Leu	Val	Lys	Ala	Ala	Lys	Ser	Ala	Lys	Leu	Arg	Lys	Leu	Pro	Pro	
				230					235					240	
Phe	Leu	Thr	Val	Ser	Leu	Leu	Arg	Phe	Asn	Phe	Asp	Phe	Val	Lys	
				245					250					255	
Cys	Glu	Arg	Tyr	Lys	Glu	Thr	Ser	Cys	Tyr	Thr	Phe	Pro	Leu	Arg	
				260					265					270	
Ile	Asn	Leu	Lys	Pro	Phe	Cys	Glu	Gln	Ser	Glu	Leu	Asp	Asp	Leu	
				275					280					285	
Glu	Tyr	Ile	Tyr	Asp	Leu	Phe	Ser	Val	Ile	Ile	His	Lys	Gly	Gly	
				290					295					300	
Cys	Tyr	Gly	Gly	His	Tyr	His	Val	Tyr	Ile	Lys	Asp	Val	Asp	His	
				305					310					315	
Leu	Gly	Asn	Trp	Gln	Phe	Gln	Glu	Glu	Lys	Ser	Lys	Pro	Asp	Val	
				320					325					330	
Asn	Leu	Lys	Asp	Leu	Gln	Ser	Glu	Glu	Glu	Ile	Asp	His	Pro	Leu	
				335					340					345	
Met	Ile	Leu	Lys	Ala	Ile	Leu	Leu	Glu	Glu	Glu	Asn	Asn	Leu	Ile	
				350					355					360	
Pro	Val	Asp	Gln	Leu	Gly	Gln	Lys	Leu	Leu	Lys	Lys	Ile	Gly	Ile	
				365					370					375	
Ser	Trp	Asn	Lys	Lys	Tyr	Arg	Lys	Gln	His	Gly	Pro	Leu	Arg	Lys	
				380					385					390	
Phe	Leu	Gln	Leu	His	Ser	Gln	Ile	Phe	Leu	Leu	Ser	Ser	Asp	Glu	
				395					400					405	
Ser	Thr	Val	Arg	Leu	Leu	Lys	Asn	Ser	Ser	Leu	Gln	Ala	Glu	Ser	
				410					415					420	
Asp	Phe	Gln	Arg	Asn	Asp	Gln	Gln	Ile	Phe	Lys	Met	Leu	Pro	Pro	
				425					430					435	
Glu	Ser	Pro	Gly	Leu	Asn	Asn	Ser	Ile	Ser	Cys	Pro	His	Trp	Phe	
				440					445					450	
Asp	Ile	Asn	Asp	Ser	Lys	Val	Gln	Pro	Ile	Arg	Glu	Lys	Asp	Ile	
				455					460					465	
Glu	Gln	Gln	Phe	Gln	Gly	Lys	Glu	Ser	Ala	Tyr	Met	Leu	Phe	Tyr	
				470					475					480	
Arg	Lys	Ser	Gln	Leu	Gln	Arg	Pro	Pro	Glu	Ala	Arg	Ala	Asn	Pro	
				485					490					495	
Arg	Tyr	Gly	Val	Pro	Cys	His	Leu	Leu	Asn	Glu	Met	Asp	Ala	Ala	
				500					505					510	
Asn	Ile	Glu	Leu	Gln	Thr	Lys	Arg	Ala	Glu	Cys	Asp	Ser	Ala	Asn	
				515					520					525	
Asn	Thr	Phe	Glu	Leu	His	Leu	His	Leu	Gly	Pro	Gln	Tyr	His	Phe	
				530					535					540	
Phe	Asn	Gly	Ala	Leu	His	Pro	Val	Val	Ser	Gln	Thr	Glu	Ser	Val	
				545					550					555	
Trp	Asp	Leu	Thr	Phe	Asp	Lys	Arg	Lys	Thr	Leu	Gly	Asp	Leu	Arg	

Gln Ser Ile Phe	560	Gln Leu Leu Glu Phe	565	Trp Glu Gly Asp Met	570
Leu Ser Val Ala	575	Lys Leu Val Pro Ala	580	Gly Leu His Ile Tyr	585
Ser Leu Gly Gly	590	Asp Glu Leu Thr Leu	595	Cys Glu Thr Glu Ile	600
Asp Gly Glu Asp	605	Ile Phe Val Trp Asn	610	Gly Val Glu Val Gly	615
Val His Ile Gln	620	Thr Gly Ile Asp Cys	625	Glu Pro Leu Leu Leu	630
Val Leu His Leu	635	Asp Thr Ser Ser Asp	640	Gly Glu Lys Cys Cys	645
Val Ile Glu Ser	650	Pro His Val Phe Pro	655	Ala Asn Ala Glu Val	660
Thr Val Leu Thr	665	Ala Leu Ala Ile Pro	670	Ala Gly Val Ile Phe	675
Asn Ser Ala Gly	680	Cys Pro Gly Gly Glu	685	Trp Thr Ala Ile	690
Lys Glu Asp Met	695	Arg Lys Thr Phe Arg	700	Glu Gln Gly Leu Arg	705
Gly Ser Ser Ile	710	Leu Ile Gln Asp Ser	715	His Asp Asp Asn Ser	720
Leu Thr Lys Glu	725	Glu Lys Trp Val Thr	730	Ser Met Asn Glu Ile	735
Trp Leu His Val	740	Lys Asn Leu Cys Gln	745	Leu Glu Ser Glu Glu	750
Gln Val Lys Ile	755	Ser Ala Thr Val Asn	760	Thr Met Val Phe Asp	765
Arg Ile Lys Ala	770	Ile Lys Glu Leu Lys	775	Leu Met Lys Glu Leu	780
Asp Asn Ser Cys	785	Leu Arg Pro Ile Asp	790	Arg Asn Gly Lys Leu	795
Cys Pro Val Pro	800	Asp Ser Tyr Thr Leu	805	Lys Glu Ala Glu Leu	810
Met Gly Ser Ser	815	Leu Gly Leu Cys Leu	820	Gly Lys Ala Pro Ser	825
Ser Gln Leu Phe	830	Leu Phe Phe Ala Met	835	Gly Ser Asp Val Gln	840
Gly Thr Glu Met	845	Glu Ile Val Val Glu	850	Glu Thr Ile Ser Val	855
Asp Cys Leu Lys	860	Leu Met Leu Lys Lys	865	Ser Gly Leu Gln Gly	870
Ala Trp His Leu	875	Arg Lys Met Asp Trp	880	Cys Tyr Glu Ala Gly	885
Pro Leu Cys Glu	890	Glu Asn Ser Ala Arg	895	Ser Gln Leu Ile Thr	900
Gly Thr Gly Phe	905	Ser Phe Gln Pro Cys	910	Gln Asp Ala Thr Leu	915
Glu Leu Leu Ile	920	Cys Ser Gly Asp Thr	925	Leu Leu Leu Ile Glu	930
Gln Leu Pro Pro	935	Leu Gly Phe Leu Lys	940	Val Pro Ile Trp Trp	945
Gln Leu Gln Gly	950	Pro Ser Gly His Trp	955	Glu Ser His Gln Asp	960
Thr Asn Cys Thr	965	Ser Ser Trp Gly Arg	970	Val Trp Arg Ala Thr	975
Ser Gln Gly Ala	980	Ser Gly Asn Glu Pro	985	Ala Gln Val Ser Leu	990
Tyr Leu Gly Asp	995	Ile Glu Ile Ser Glu	1000	Ala Thr Leu Ala Glu	1005
Leu Lys Ser Gln	1010	Ala Met Thr Leu Pro	1015	Phe Leu Glu Phe Gly	1020
	1025		1030		1035

Val Pro Ser Pro Ala His Leu Arg Ala Trp Thr Val Glu Arg Lys
 1040 1045 1050
 Arg Pro Gly Arg Leu Leu Arg Thr Asp Arg Gln Pro Leu Arg Glu
 1055 1060 1065
 Tyr Lys Leu Gly Arg Arg Ile Glu Ile Cys Leu Glu Pro Leu Gln
 1070 1075 1080
 Lys Gly Glu Asn Leu Gly Pro Gln Asp Val Leu Leu Arg Thr Gln
 1085 1090 1095
 Val Arg Ile Pro Gly Glu Arg Thr Tyr Ala Pro Ala Leu Asp Leu
 1100 1105 1110
 Val Trp Asn Ala Ala Gln Gly Gly Thr Ala Gly Ser Leu Arg Gln
 1115 1120 1125
 Arg Val Ala Asp Phe Tyr Arg Leu Pro Val Glu Lys Ile Glu Ile
 1130 1135 1140
 Ala Lys Tyr Phe Pro Glu Lys Phe Glu Trp Leu Pro Ile Ser Ser
 1145 1150 1155
 Trp Asn Gln Gln Ile Thr Lys Arg Lys Lys Lys Lys Lys Gln Asp
 1160 1165 1170
 Tyr Leu Gln Gly Ala Pro Tyr Tyr Leu Lys Asp Gly Asp Thr Ile
 1175 1180 1185
 Gly Val Lys Asn Leu Leu Ile Asp Asp Asp Asp Phe Ser Thr
 1190 1195 1200
 Ile Arg Asp Asp Thr Gly Lys Glu Lys Gln Lys Gln Arg Ala Leu
 1205 1210 1215
 Gly Arg Arg Lys Ser Gln Glu Ala Leu His Glu Gln Ser Ser Tyr
 1220 1225 1230
 Ile Leu Ser Ser Ala Glu Thr Pro Ala Arg Pro Arg Ala Pro Glu
 1235 1240 1245
 Thr Ser Leu Ser Ile His Val Gly Ser Phe Arg
 1250 1255

<210> 4

<211> 755

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2875922CD1

<400> 4

Met Lys Lys Gln Arg Lys Ile Leu Trp Arg Lys Gly Ile His Leu
 1 5 10 15
 Ala Phe Ser Glu Lys Trp Asn Thr Gly Phe Gly Gly Phe Lys Lys
 20 25 30
 Phe Tyr Phe His Gln His Leu Cys Ile Leu Lys Ala Lys Leu Gly
 35 40 45
 Arg Pro Val Thr Trp Asn Arg Gln Leu Arg His Phe Gln Gly Arg
 50 55 60
 Lys Lys Ala Leu Gln Ile Gln Lys Thr Trp Ile Lys Asp Glu Pro
 65 70 75
 Leu Cys Ala Lys Thr Lys Phe Asn Val Ala Thr Gln Asn Val Ser
 80 85 90
 Thr Leu Ser Ser Lys Val Lys Arg Lys Asp Ala Lys His Phe Ile
 95 100 105
 Ser Ser Ser Lys Thr Leu Leu Arg Leu Gln Ala Glu Lys Leu Leu
 110 115 120
 Ser Ser Ala Lys Asn Ser Asp His Glu Tyr Cys Arg Glu Lys Asn
 125 130 135
 Leu Leu Lys Ala Val Thr Asp Phe Pro Ser Asn Ser Ala Leu Gly
 140 145 150
 Gln Ala Asn Gly His Arg Pro Arg Thr Asp Pro Gln Pro Ser Asp
 155 160 165

Phe	Pro	Met	Lys	Phe	Asn	Gly	Glu	Ser	Gln	Ser	Pro	Gly	Glu	Ser
				170					175					180
Gly	Thr	Ile	Val	Val	Thr	Leu	Asn	Asn	His	Lys	Arg	Lys	Gly	Phe
				185					190					195
Cys	Tyr	Gly	Cys	Cys	Gln	Gly	Pro	Glu	His	His	Arg	Asn	Gly	Gly
				200					205					210
Pro	Leu	Ile	Pro	Lys	Lys	Phe	Gln	Leu	Asn	Gln	His	Arg	Arg	Ile
				215					220					225
Lys	Leu	Ser	Pro	Leu	Met	Met	Tyr	Glu	Lys	Leu	Ser	Met	Ile	Arg
				230					235					240
Phe	Arg	Tyr	Arg	Ile	Leu	Arg	Ser	Gln	His	Phe	Arg	Thr	Lys	Ser
				245					250					255
Lys	Val	Cys	Lys	Leu	Arg	Lys	Ala	Gln	Arg	Ser	Trp	Val	Gln	Lys
				260					265					270
Val	Thr	Gly	Asp	His	Gln	Glu	Thr	Arg	Arg	Glu	Asn	Gly	Glu	Gly
				275					280					285
Gly	Ser	Cys	Ser	Pro	Phe	Pro	Ser	Pro	Glu	Pro	Lys	Asp	Pro	Ser
				290					295					300
Cys	Arg	His	Gln	Pro	Tyr	Phe	Pro	Asp	Met	Asp	Ser	Ser	Ala	Val
				305					310					315
Val	Lys	Gly	Thr	Asn	Ser	His	Val	Pro	Asp	Cys	His	Thr	Lys	Gly
				320					325					330
Ser	Ser	Phe	Leu	Gly	Lys	Glu	Leu	Ser	Leu	Asp	Glu	Ala	Phe	Pro
				335					340					345
Asp	Gln	Gln	Asn	Gly	Ser	Ala	Thr	Asn	Ala	Trp	Asp	Gln	Ser	Ser
				350					355					360
Cys	Ser	Ser	Pro	Lys	Trp	Glu	Cys	Thr	Glu	Leu	Ile	His	Asp	Ile
				365					370					375
Pro	Leu	Pro	Glu	His	Arg	Ser	Asn	Thr	Met	Phe	Ile	Ser	Glu	Thr
				380					385					390
Glu	Arg	Glu	Ile	Met	Thr	Leu	Gly	Gln	Glu	Asn	Gln	Thr	Ser	Ser
				395					400					405
Val	Ser	Asp	Asp	Arg	Val	Lys	Leu	Ser	Val	Ser	Gly	Ala	Asp	Thr
				410					415					420
Ser	Val	Ser	Ser	Val	Asp	Gly	Pro	Val	Ser	Gln	Lys	Ala	Val	Gln
				425					430					435
Asn	Glu	Asn	Ser	Tyr	Gln	Met	Glu	Glu	Asp	Gly	Ser	Leu	Lys	Gln
				440					445					450
Ser	Ile	Leu	Ser	Ser	Glu	Leu	Leu	Asp	His	Pro	Tyr	Cys	Lys	Ser
				455					460					465
Pro	Leu	Glu	Ala	Pro	Leu	Val	Cys	Ser	Gly	Leu	Lys	Leu	Glu	Asn
				470					475					480
Gln	Val	Gly	Gly	Gly	Lys	Asn	Ser	Gln	Lys	Ala	Ser	Pro	Val	Asp
				485					490					495
Asp	Glu	Gln	Leu	Ser	Val	Cys	Leu	Ser	Gly	Phe	Leu	Asp	Glu	Val
				500					505					510
Met	Lys	Lys	Tyr	Gly	Ser	Leu	Val	Pro	Leu	Ser	Glu	Lys	Glu	Val
				515					520					525
Leu	Gly	Arg	Leu	Lys	Asp	Val	Phe	Asn	Glu	Asp	Phe	Ser	Asn	Arg
				530					535					540
Lys	Pro	Phe	Ile	Asn	Arg	Glu	Ile	Thr	Asn	Tyr	Arg	Ala	Arg	His
				545					550					555
Gln	Lys	Cys	Asn	Phe	Arg	Ile	Phe	Tyr	Asn	Lys	His	Met	Leu	Asp
				560					565					570
Met	Asp	Asp	Leu	Ala	Thr	Leu	Asp	Gly	Gln	Asn	Trp	Leu	Asn	Asp
				575					580					585
Gln	Val	Ile	Asn	Met	Tyr	Gly	Glu	Leu	Ile	Met	Asp	Ala	Val	Pro
				590					595					600
Asp	Lys	Val	His	Phe	Phe	Asn	Ser	Phe	Phe	His	Arg	Gln	Leu	Val
				605					610					615
Thr	Lys	Gly	Tyr	Asn	Gly	Val	Lys	Arg	Trp	Thr	Lys	Lys	Val	Asp
				620					625					630
Leu	Phe	Lys	Lys	Ser	Leu	Leu	Leu	Ile	Pro	Ile	His	Leu	Glu	Val

	635		640		645
His Trp Ser Leu	Ile Thr Val Thr Leu	Ser Asn Arg Ile Ile	Ser		
	650		655		660
Phe Tyr Asp Ser	Gln Gly Ile His Phe	Lys Phe Cys Val Glu	Asn		
	665		670		675
Ile Arg Lys Tyr	Leu Leu Thr Glu Ala	Arg Glu Lys Asn Arg	Pro		
	680		685		690
Glu Phe Leu Gln	Gly Trp Gln Thr Ala	Val Thr Lys Cys Ile	Pro		
	695		700		705
Gln Gln Lys Asn	Asp Ser Asp Cys Gly	Val Phe Val Leu Gln	Tyr		
	710		715		720
Cys Lys Cys Leu	Ala Leu Glu Gln Pro	Phe Gln Phe Ser Gln	Glu		
	725		730		735
Asp Met Pro Arg	Val Arg Lys Arg Ile	Tyr Lys Glu Leu Cys	Glu		
	740		745		750
Cys Arg Leu Met	Asp				
	755				

<210> 5

<211> 1034

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 8158136CD1

<400> 5

Met Ala Pro Arg Leu	Gln Leu Glu Lys Ala	Ala Trp Arg Trp Ala	
1	5	10	15
Glu Thr Val Arg Pro	Glu Glu Val Ser	Gln Glu His Ile Glu	Thr
	20	25	30
Ala Tyr Arg Ile Trp	Leu Glu Pro Cys	Ile Arg Gly Val Cys	Arg
	35	40	45
Arg Asn Cys Lys Gly	Asn Pro Asn Cys	Leu Val Gly Ile Gly	Glu
	50	55	60
His Ile Trp Leu Gly	Glu Ile Asp Glu	Asn Ser Phe His Asn	Ile
	65	70	75
Asp Asp Pro Asn Cys	Glu Arg Arg Lys	Lys Asn Ser Phe Val	Gly
	80	85	90
Leu Thr Asn Leu Gly	Ala Thr Cys Tyr	Val Asn Thr Phe Leu	Gln
	95	100	105
Val Trp Phe Leu Asn	Leu Glu Leu Arg	Gln Ala Leu Tyr Leu	Cys
	110	115	120
Pro Ser Thr Cys Ser	Asp Tyr Met Leu	Gly Asp Gly Ile Gln	Glu
	125	130	135
Glu Lys Asp Tyr Glu	Pro Gln Thr Ile	Cys Glu His Leu Gln	Tyr
	140	145	150
Leu Phe Ala Leu Leu	Gln Asn Ser Asn	Arg Arg Tyr Ile Asp	Pro
	155	160	165
Ser Gly Phe Val Lys	Ala Leu Gly Leu	Asp Thr Gly Gln Gln	Gln
	170	175	180
Asp Ala Gln Glu Phe	Ser Lys Leu Phe	Met Ser Leu Leu Glu	Asp
	185	190	195
Thr Leu Ser Lys Gln	Lys Asn Pro Asp	Val Arg Asn Ile Val	Gln
	200	205	210
Gln Gln Phe Cys Gly	Glu Tyr Ala Tyr	Val Thr Val Cys Asn	Gln
	215	220	225
Cys Gly Arg Glu Ser	Lys Leu Leu Ser	Lys Phe Tyr Glu Leu	Glu
	230	235	240
Leu Asn Ile Gln Gly	His Lys Gln Leu	Thr Asp Cys Ile Ser	Glu
	245	250	255
Phe Leu Lys Glu Glu	Lys Leu Glu Gly	Asp Asn Arg Tyr Phe	Cys

	260		265		270
Glu Asn Cys Gln Ser Lys Gln Asn Ala Thr Arg Lys Ile Arg Leu					
	275		280		285
Leu Ser Leu Pro Cys Thr Leu Asn Leu Gln Leu Met Arg Phe Val					
	290		295		300
Phe Asp Arg Gln Thr Gly His Lys Lys Lys Leu Asn Thr Tyr Ile					
	305		310		315
Gly Phe Ser Glu Ile Leu Asp Met Glu Pro Tyr Val Glu His Lys					
	320		325		330
Gly Gly Ser Tyr Val Tyr Glu Leu Ser Ala Val Leu Ile His Arg					
	335		340		345
Gly Val Ser Ala Tyr Ser Gly His Tyr Ile Ala His Val Lys Asp					
	350		355		360
Pro Gln Ser Gly Glu Trp Tyr Lys Phe Asn Asp Glu Asp Ile Glu					
	365		370		375
Lys Met Glu Gly Lys Lys Leu Gln Leu Gly Ile Glu Glu Asp Leu					
	380		385		390
Ala Glu Pro Ser Lys Ser Gln Thr Arg Lys Pro Lys Cys Gly Lys					
	395		400		405
Gly Thr His Cys Ser Arg Asn Ala Tyr Met Leu Val Tyr Arg Leu					
	410		415		420
Gln Thr Gln Glu Lys Pro Asn Thr Thr Val Gln Val Pro Ala Phe					
	425		430		435
Leu Gln Glu Leu Val Asp Arg Asp Asn Ser Lys Phe Glu Glu Trp					
	440		445		450
Cys Ile Glu Met Ala Glu Met Arg Lys Gln Ser Val Asp Lys Gly					
	455		460		465
Lys Ala Lys His Glu Glu Val Lys Glu Leu Tyr Gln Arg Leu Pro					
	470		475		480
Ala Gly Ala Glu Pro Tyr Glu Phe Val Ser Leu Glu Trp Leu Gln					
	485		490		495
Lys Trp Leu Asp Glu Ser Thr Pro Thr Lys Pro Ile Asp Asn His					
	500		505		510
Ala Cys Leu Cys Ser His Asp Lys Leu His Pro Asp Lys Ile Ser					
	515		520		525
Ile Met Lys Arg Ile Ser Glu Tyr Ala Ala Asp Ile Phe Tyr Ser					
	530		535		540
Arg Tyr Gly Gly Gly Pro Arg Leu Thr Val Lys Ala Leu Cys Lys					
	545		550		555
Glu Cys Val Val Glu Arg Cys Arg Ile Leu Arg Leu Lys Asn Gln					
	560		565		570
Leu Asn Glu Asp Tyr Lys Thr Val Asn Asn Leu Leu Lys Ala Ala					
	575		580		585
Val Lys Gly Asp Gly Phe Trp Val Gly Lys Ser Ser Leu Arg Ser					
	590		595		600
Trp Arg Gln Leu Ala Leu Glu Gln Leu Asp Glu Gln Asp Gly Asp					
	605		610		615
Ala Glu Gln Ser Asn Gly Lys Met Asn Gly Ser Thr Leu Asn Lys					
	620		625		630
Asp Glu Ser Lys Glu Glu Arg Lys Glu Glu Glu Glu Leu Asn Phe					
	635		640		645
Asn Glu Asp Ile Leu Cys Pro His Gly Glu Leu Cys Ile Ser Glu					
	650		655		660
Asn Glu Arg Arg Leu Val Ser Lys Glu Ala Trp Ser Lys Leu Gln					
	665		670		675
Gln Tyr Phe Pro Lys Ala Pro Glu Phe Pro Ser Tyr Lys Glu Cys					
	680		685		690
Cys Ser Gln Cys Lys Ile Leu Glu Arg Glu Gly Glu Glu Asn Glu					
	695		700		705
Ala Leu His Lys Met Ile Ala Asn Glu Gln Lys Thr Ser Leu Pro					
	710		715		720
Asn Leu Phe Gln Asp Lys Asn Arg Pro Cys Leu Ser Asn Trp Pro					
	725		730		735

Glu	Asp	Thr	Asp	Val	Leu	Tyr	Ile	Val	Ser	Gln	Phe	Phe	Val	Glu	
				740					745					750	
Glu	Trp	Arg	Lys	Phe	Val	Arg	Lys	Pro	Thr	Arg	Cys	Ser	Pro	Val	
				755					760					765	
Ser	Ser	Val	Gly	Asn	Ser	Ala	Leu	Leu	Cys	Pro	His	Gly	Gly	Leu	
				770					775					780	
Met	Phe	Thr	Phe	Ala	Ser	Met	Thr	Lys	Glu	Asp	Ser	Lys	Leu	Ile	
				785					790					795	
Ala	Leu	Ile	Trp	Pro	Ser	Glu	Trp	Gln	Met	Ile	Gln	Lys	Leu	Phe	
				800					805					810	
Val	Val	Asp	His	Val	Ile	Lys	Ile	Thr	Arg	Ile	Glu	Val	Gly	Asp	
				815					820					825	
Val	Asn	Pro	Ser	Glu	Thr	Gln	Tyr	Ile	Ser	Glu	Pro	Lys	Leu	Cys	
				830					835					840	
Pro	Glu	Cys	Arg	Glu	Gly	Leu	Leu	Cys	Gln	Gln	Gln	Arg	Asp	Leu	
				845					850					855	
Arg	Glu	Tyr	Thr	Gln	Ala	Thr	Ile	Tyr	Val	His	Lys	Val	Val	Asp	
				860					865					870	
Asn	Lys	Lys	Val	Met	Lys	Asp	Ser	Ala	Pro	Glu	Leu	Asn	Val	Ser	
				875					880					885	
Ser	Ser	Glu	Thr	Glu	Glu	Asp	Lys	Glu	Glu	Ala	Lys	Pro	Asp	Gly	
				890					895					900	
Glu	Lys	Asp	Pro	Asp	Phe	Asn	Gln	Ser	Asn	Gly	Gly	Thr	Lys	Arg	
				905					910					915	
Gln	Lys	Ile	Ser	His	Gln	Asn	Tyr	Ile	Ala	Tyr	Gln	Lys	Gln	Val	
				920					925					930	
Ile	Arg	Arg	Ser	Met	Arg	His	Arg	Lys	Val	Arg	Gly	Glu	Lys	Ala	
				935					940					945	
Leu	Leu	Val	Ser	Ala	Asn	Gln	Thr	Leu	Lys	Glu	Leu	Lys	Ile	Gln	
				950					955					960	
Ile	Met	His	Ala	Phe	Ser	Val	Ala	Pro	Phe	Asp	Gln	Asn	Leu	Ser	
				965					970					975	
Ile	Asp	Gly	Lys	Ile	Leu	Ser	Asp	Asp	Cys	Ala	Thr	Leu	Gly	Thr	
				980					985					990	
Leu	Gly	Val	Ile	Pro	Glu	Ser	Val	Ile	Leu	Leu	Lys	Ala	Asp	Glu	
				995					1000					1005	
Pro	Ile	Ala	Asp	Tyr	Ala	Ala	Met	Asp	Asp	Val	Met	Gln	Val	Cys	
				1010					1015					1020	
Met	Pro	Glu	Glu	Gly	Phe	Lys	Gly	Thr	Gly	Leu	Leu	Gly	His		
				1025					1030						

<210> 6

<211> 1236

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5969491CD1

<400> 6

Met	Phe	Gly	Asp	Leu	Phe	Glu	Glu	Glu	Tyr	Ser	Thr	Val	Ser	Asn	
1				5					10					15	
Asn	Gln	Tyr	Gly	Lys	Gly	Lys	Lys	Leu	Lys	Thr	Lys	Ala	Leu	Glu	
				20					25					30	
Pro	Pro	Ala	Pro	Arg	Glu	Phe	Thr	Asn	Leu	Ser	Gly	Ile	Arg	Asn	
				35					40					45	
Gln	Gly	Gly	Thr	Cys	Tyr	Leu	Asn	Ser	Leu	Leu	Gln	Thr	Leu	His	
				50					55					60	
Phe	Thr	Pro	Glu	Phe	Arg	Glu	Ala	Leu	Phe	Ser	Leu	Gly	Pro	Glu	
				65					70					75	
Glu	Leu	Gly	Leu	Phe	Glu	Asp	Lys	Asp	Lys	Pro	Asp	Ala	Lys	Val	
				80					85					90	

Arg	Ile	Ile	Pro	Leu	Gln	Leu	Gln	Arg	Leu	Phe	Ala	Gln	Leu	Leu	
				95					100					105	
Leu	Leu	Asp	Gln	Glu	Ala	Ala	Ser	Thr	Ala	Asp	Leu	Thr	Asp	Ser	
				110					115					120	
Phe	Gly	Trp	Thr	Ser	Asn	Glu	Glu	Met	Arg	Gln	His	Asp	Val	Gln	
				125					130					135	
Glu	Leu	Asn	Arg	Ile	Leu	Phe	Ser	Ala	Leu	Glu	Thr	Ser	Leu	Val	
				140					145					150	
Gly	Thr	Ser	Gly	His	Asp	Leu	Ile	Tyr	Arg	Leu	Tyr	His	Gly	Thr	
				155					160					165	
Ile	Val	Asn	Gln	Ile	Val	Cys	Lys	Glu	Cys	Lys	Asn	Val	Ser	Glu	
				170					175					180	
Arg	Gln	Glu	Asp	Phe	Leu	Asp	Leu	Thr	Val	Ala	Val	Lys	Asn	Val	
				185					190					195	
Ser	Gly	Leu	Glu	Asp	Ala	Leu	Trp	Asn	Met	Tyr	Val	Glu	Glu	Glu	
				200					205					210	
Val	Phe	Asp	Cys	Asp	Asn	Leu	Tyr	His	Cys	Gly	Thr	Cys	Asp	Arg	
				215					220					225	
Leu	Val	Lys	Ala	Ala	Lys	Ser	Ala	Lys	Leu	Arg	Lys	Leu	Pro	Pro	
				230					235					240	
Phe	Leu	Thr	Val	Ser	Leu	Leu	Arg	Phe	Asn	Phe	Asp	Phe	Val	Lys	
				245					250					255	
Cys	Glu	Arg	Tyr	Lys	Glu	Thr	Ser	Cys	Tyr	Thr	Phe	Pro	Leu	Arg	
				260					265					270	
Ile	Asn	Leu	Lys	Pro	Phe	Cys	Glu	Gln	Ser	Glu	Leu	Asp	Asp	Leu	
				275					280					285	
Glu	Tyr	Ile	Tyr	Asp	Leu	Phe	Ser	Val	Ile	Ile	His	Lys	Gly	Gly	
				290					295					300	
Cys	Tyr	Gly	Gly	His	Tyr	His	Val	Tyr	Ile	Lys	Asp	Val	Asp	His	
				305					310					315	
Leu	Gly	Asn	Trp	Gln	Phe	Gln	Glu	Glu	Lys	Ser	Lys	Pro	Asp	Val	
				320					325					330	
Asn	Leu	Lys	Asp	Leu	Gln	Ser	Glu	Glu	Glu	Ile	Asp	His	Pro	Leu	
				335					340					345	
Met	Ile	Leu	Lys	Ala	Ile	Leu	Leu	Glu	Glu	Glu	Asn	Asn	Leu	Ile	
				350					355					360	
Pro	Val	Asp	Gln	Leu	Gly	Gln	Lys	Leu	Leu	Lys	Lys	Ile	Gly	Ile	
				365					370					375	
Ser	Trp	Asn	Lys	Lys	Tyr	Arg	Lys	Gln	His	Gly	Pro	Leu	Arg	Lys	
				380					385					390	
Phe	Leu	Gln	Leu	His	Ser	Gln	Ile	Phe	Leu	Leu	Ser	Ser	Asp	Glu	
				395					400					405	
Ser	Thr	Val	Arg	Leu	Leu	Lys	Asn	Ser	Ser	Leu	Gln	Ala	Glu	Ser	
				410					415					420	
Asp	Phe	Gln	Arg	Asn	Asp	Gln	Gln	Ile	Phe	Lys	Met	Leu	Pro	Pro	
				425					430					435	
Glu	Ser	Pro	Gly	Leu	Asn	Asn	Ser	Ile	Ser	Cys	Pro	His	Trp	Phe	
				440					445					450	
Asp	Ile	Asn	Asp	Ser	Lys	Val	Gln	Pro	Ile	Arg	Glu	Lys	Asp	Ile	
				455					460					465	
Glu	Gln	Gln	Phe	Gln	Gly	Lys	Glu	Ser	Ala	Tyr	Met	Leu	Phe	Tyr	
				470					475					480	
Arg	Lys	Ser	Gln	Leu	Gln	Arg	Pro	Pro	Glu	Ala	Arg	Ala	Asn	Pro	
				485					490					495	
Arg	Tyr	Gly	Val	Pro	Cys	His	Leu	Leu	Asn	Glu	Met	Asp	Ala	Ala	
				500					505					510	
Asn	Ile	Glu	Leu	Gln	Thr	Lys	Arg	Ala	Glu	Cys	Asp	Ser	Ala	Asn	
				515					520					525	
Asn	Thr	Phe	Glu	Leu	His	Leu	His	Leu	Gly	Pro	Gln	Tyr	His	Phe	
				530					535					540	
Phe	Asn	Gly	Ala	Leu	His	Pro	Val	Val	Ser	Gln	Thr	Glu	Ser	Val	
				545					550					555	
Trp	Asp	Leu	Thr	Phe	Asp	Lys	Arg	Lys	Thr	Leu	Gly	Asp	Leu	Arg	

Gln Ser Ile Phe	560	Gln Leu Leu Glu Phe	565	Trp Glu Gly Asp Met Val	570
Leu Ser Val Ala	575	Lys Leu Val Pro Ala	580	Gly Leu His Ile Tyr Gln	585
Ser Leu Gly Gly	590	Asp Glu Leu Thr Leu	595	Cys Glu Thr Glu Ile Ala	600
Asp Gly Glu Asp	605	Ile Phe Val Trp Asn	610	Gly Val Glu Val Gly Gly	615
Val His Ile Gln	620	Thr Gly Ile Asp Cys	625	Glu Pro Leu Leu Leu Asn	630
Val Leu His Leu	635	Asp Thr Ser Ser Asp	640	Gly Glu Lys Cys Cys Gln	645
Val Ile Glu Ser	650	Pro His Val Phe Pro	655	Ala Asn Ala Glu Val Gly	660
Thr Val Leu Thr	665	Ala Leu Ala Ile Pro	670	Ala Gly Val Ile Phe Ile	675
Asn Ser Ala Gly	680	Cys Pro Gly Gly Glu	685	Gly Trp Thr Ala Ile Pro	690
Lys Glu Asp Met	695	Arg Lys Thr Phe Arg	700	Glu Gln Gly Leu Arg Asn	705
Gly Ser Ser Ile	710	Leu Ile Gln Asp Ser	715	His Asp Asp Asn Ser Leu	720
Leu Thr Lys Glu	725	Glu Lys Trp Val Thr	730	Ser Met Asn Glu Ile Asp	735
Trp Leu His Val	740	Lys Asn Leu Cys Gln	745	Leu Glu Ser Glu Glu Lys	750
Gln Val Lys Ile	755	Ser Ala Thr Val Asn	760	Thr Met Val Phe Asp Ile	765
Arg Ile Lys Ala	770	Ile Lys Glu Leu Lys	775	Leu Met Lys Glu Leu Ala	780
Asp Asn Ser Cys	785	Leu Arg Pro Ile Asp	790	Arg Asn Gly Lys Leu Leu	795
Cys Pro Val Pro	800	Asp Ser Tyr Thr Leu	805	Lys Glu Ala Glu Leu Lys	810
Met Gly Ser Ser	815	Leu Gly Leu Cys Leu	820	Gly Lys Ala Pro Ser Ser	825
Ser Gln Leu Phe	830	Leu Phe Phe Ala Met	835	Gly Ser Asp Val Gln Pro	840
Gly Thr Glu Met	845	Glu Ile Val Val Glu	850	Glu Thr Ile Ser Val Arg	855
Asp Cys Leu Lys	860	Leu Met Leu Lys Lys	865	Ser Gly Leu Gln Gly Asp	870
Ala Trp His Leu	875	Arg Lys Met Asp Trp	880	Cys Tyr Glu Ala Gly Glu	885
Pro Leu Cys Glu	890	Glu Asp Ala Thr Leu	895	Lys Glu Leu Leu Ile Cys	900
Ser Gly Asp Thr	905	Leu Leu Leu Ile Glu	910	Gly Gln Leu Pro Pro Leu	915
Gly Phe Leu Lys	920	Val Pro Ile Trp Trp	925	Tyr Gln Leu Gln Gly Pro	930
Ser Gly His Trp	935	Glu Ser His Gln Asp	940	Gln Thr Asn Cys Thr Ser	945
Ser Trp Gly Arg	950	Val Trp Arg Ala Thr	955	Ser Ser Gln Gly Ala Ser	960
Gly Asn Glu Pro	965	Ala Gln Val Ser Leu	970	Leu Tyr Leu Gly Asp Ile	975
Glu Ile Ser Glu	980	Asp Ala Thr Leu Ala	985	Glu Leu Lys Ser Gln Ala	990
Met Thr Leu Pro	995	Pro Phe Leu Glu Phe	1000	Gly Val Pro Ser Pro Ala	1005
His Leu Arg Ala	1010	Trp Thr Val Glu Arg	1015	Lys Arg Pro Gly Arg Leu	1020
	1025		1030		1035


```

Leu Arg Thr Asp Arg Gln Pro Leu Arg Glu Tyr Lys Leu Gly Arg
1040 1045 1050
Arg Ile Glu Ile Cys Leu Glu Pro Leu Gln Lys Gly Glu Asn Leu
1055 1060 1065
Gly Pro Gln Asp Val Leu Leu Arg Thr Gln Val Arg Ile Pro Gly
1070 1075 1080
Glu Arg Thr Tyr Ala Pro Ala Leu Asp Leu Val Trp Asn Ala Ala
1085 1090 1095
Gln Gly Gly Thr Ala Gly Ser Leu Arg Gln Arg Val Ala Asp Phe
1100 1105 1110
Tyr Arg Leu Pro Val Glu Lys Ile Glu Ile Ala Lys Tyr Phe Pro
1115 1120 1125
Glu Lys Phe Glu Trp Leu Pro Ile Ser Ser Trp Asn Gln Gln Ile
1130 1135 1140
Thr Lys Arg Lys Lys Lys Lys Lys Gln Asp Tyr Leu Gln Gly Ala
1145 1150 1155
Pro Tyr Tyr Leu Lys Asp Gly Asp Thr Ile Gly Val Lys Asn Leu
1160 1165 1170
Leu Ile Asp Asp Asp Asp Asp Phe Ser Thr Ile Arg Asp Asp Thr
1175 1180 1185
Gly Lys Glu Lys Gln Lys Gln Arg Ala Leu Gly Arg Arg Lys Ser
1190 1195 1200
Gln Glu Ala Leu His Glu Gln Ser Ser Tyr Ile Leu Ser Ser Ala
1205 1210 1215
Glu Thr Pro Ala Arg Pro Arg Ala Pro Glu Thr Ser Leu Ser Ile
1220 1225 1230
His Val Gly Ser Phe Arg
1235

```

<210> 7

<211> 545

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7497367CD1

<400> 7

```

Met Leu Pro Gly Ala Trp Leu Leu Trp Thr Ser Leu Leu Leu Leu
1 5 10 15
Ala Arg Pro Ala Gln Pro Cys Pro Met Gly Cys Asp Cys Phe Val
20 25 30
Gln Glu Val Phe Cys Ser Asp Glu Glu Leu Ala Thr Val Pro Leu
35 40 45
Asp Ile Pro Pro Tyr Thr Lys Asn Ile Ile Phe Val Glu Thr Ser
50 55 60
Phe Thr Thr Leu Glu Thr Arg Ala Phe Gly Ser Asn Pro Asn Leu
65 70 75
Thr Lys Val Val Phe Leu Asn Thr Gln Leu Cys Gln Phe Arg Pro
80 85 90
Asp Ala Phe Gly Gly Leu Pro Arg Leu Glu Asp Leu Glu Val Thr
95 100 105
Gly Ser Ser Phe Leu Asn Leu Ser Thr Asn Ile Phe Ser Asn Leu
110 115 120
Thr Ser Leu Gly Lys Leu Thr Leu Asn Phe Asn Met Leu Glu Ala
125 130 135
Leu Pro Glu Gly Leu Phe Gln His Leu Ala Ala Leu Glu Ser Leu
140 145 150
His Leu Gln Gly Asn Gln Leu Gln Ala Leu Pro Arg Arg Leu Phe
155 160 165
Gln Pro Leu Thr His Leu Lys Thr Leu Asn Leu Ala Gln Asn Leu
170 175 180

```

<400> 8
Met Leu Ile Leu Thr Lys Thr Ala Gly Val Phe Phe Lys Pro Ser
1 5 10 15

Lys	Arg	Lys	Val	Tyr	Glu	Phe	Leu	Arg	Ser	Phe	Asn	Phe	His	Pro
				20					25					30
Gly	Thr	Leu	Phe	Leu	His	Lys	Ile	Val	Leu	Gly	Ile	Glu	Thr	Ser
				35					40					45
Cys	Asp	Asp	Thr	Ala	Ala	Ala	Val	Val	Asp	Glu	Thr	Gly	Asn	Val
				50					55					60
Leu	Gly	Glu	Ala	Ile	His	Ser	Gln	Thr	Glu	Val	His	Leu	Lys	Thr
				65					70					75
Gly	Gly	Ile	Val	Pro	Pro	Ala	Ala	Gln	Gln	Leu	His	Arg	Glu	Asn
				80					85					90
Ile	Gln	Arg	Ile	Val	Gln	Glu	Ala	Leu	Ser	Ala	Ser	Gly	Val	Ser
				95					100					105
Pro	Ser	Asp	Leu	Ser	Ala	Ile	Ala	Thr	Thr	Ile	Lys	Pro	Gly	Leu
				110					115					120
Ala	Leu	Ser	Leu	Gly	Val	Gly	Leu	Ser	Phe	Ser	Leu	Gln	Leu	Val
				125					130					135
Gly	Gln	Leu	Lys	Lys	Pro	Phe	Ile	Pro	Ile	His	His	Met	Glu	Ala
				140					145					150
His	Ala	Leu	Thr	Ile	Arg	Leu	Thr	Asn	Lys	Val	Glu	Phe	Pro	Phe
				155					160					165
Leu	Val	Leu	Leu	Ile	Ser	Gly	Gly	His	Cys	Leu	Leu	Ala	Leu	Val
				170					175					180
Gln	Gly	Val	Ser	Asp	Phe	Leu	Leu	Leu	Gly	Lys	Ser	Leu	Asp	Ile
				185					190					195
Ala	Pro	Gly	Asp	Met	Leu	Asp	Lys	Val	Ala	Arg	Arg	Leu	Ser	Leu
				200					205					210
Ile	Lys	His	Pro	Glu	Cys	Ser	Thr	Met	Ser	Gly	Gly	Lys	Ala	Ile
				215					220					225
Glu	His	Leu	Ala	Lys	Gln	Gly	Asn	Arg	Phe	His	Phe	Asp	Ile	Lys
				230					235					240
Pro	Pro	Leu	His	His	Ala	Lys	Asn	Cys	Asp	Phe	Ser	Phe	Thr	Gly
				245					250					255
Leu	Gln	His	Val	Thr	Asp	Lys	Ile	Ile	Met	Lys	Lys	Glu	Lys	Glu
				260					265					270
Glu	Gly	Ile	Glu	Lys	Gly	Gln	Ile	Leu	Ser	Ser	Ala	Ala	Asp	Ile
				275					280					285
Ala	Ala	Thr	Val	Gln	His	Thr	Met	Ala	Cys	His	Leu	Val	Lys	Arg
				290					295					300
Thr	His	Arg	Ala	Ile	Leu	Phe	Cys	Lys	Gln	Arg	Asp	Leu	Leu	Pro
				305					310					315
Gln	Asn	Asn	Ala	Val	Leu	Val	Ala	Ser	Gly	Gly	Val	Ala	Ser	Asn
				320					325					330
Phe	Tyr	Ile	Arg	Arg	Ala	Leu	Glu	Ile	Leu	Thr	Asn	Ala	Thr	Gln
				335					340					345
Cys	Thr	Leu	Leu	Cys	Pro	Pro	Pro	Arg	Leu	Cys	Thr	Asp	Asn	Gly
				350					355					360
Ile	Met	Ile	Ala	Trp	Asn	Gly	Ile	Glu	Arg	Leu	Arg	Ala	Gly	Leu
				365					370					375
Gly	Ile	Leu	His	Asp	Ile	Glu	Gly	Ile	Arg	Tyr	Glu	Pro	Lys	Cys
				380					385					390
Pro	Leu	Gly	Val	Asp	Ile	Ser	Lys	Glu	Val	Gly	Glu	Ala	Ser	Ile
				395					400					405
Lys	Val	Pro	Gln	Leu	Lys	Met	Glu	Ile						
				410										

<210> 9

<211> 611

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1804436CD1

<400> 9

Met	Ser	Cys	Lys	Lys	Gln	Arg	Ser	Arg	Lys	His	Ser	Val	Asn	Glu
1				5					10					15
Lys	Cys	Asn	Met	Lys	Ile	Glu	His	Tyr	Phe	Ser	Pro	Val	Ser	Lys
				20					25					30
Glu	Gln	Gln	Asn	Asn	Cys	Ser	Thr	Ser	Leu	Met	Arg	Met	Glu	Ser
				35					40					45
Arg	Gly	Asp	Pro	Arg	Ala	Thr	Thr	Asn	Thr	Gln	Ala	Gln	Arg	Phe
				50					55					60
His	Ser	Pro	Lys	Lys	Asn	Pro	Glu	Asp	Gln	Thr	Met	Pro	Gln	Asn
				65					70					75
Arg	Thr	Ile	Tyr	Val	Thr	Leu	Lys	Val	Asn	His	Arg	Arg	Asn	Gln
				80					85					90
Asp	Met	Lys	Leu	Lys	Leu	Thr	His	Ser	Glu	Asn	Ser	Ser	Leu	Tyr
				95					100					105
Met	Ala	Leu	Asn	Thr	Leu	Gln	Ala	Val	Arg	Lys	Glu	Ile	Glu	Thr
				110					115					120
His	Gln	Gly	Gln	Glu	Met	Leu	Val	Arg	Gly	Thr	Glu	Gly	Ile	Lys
				125					130					135
Glu	Tyr	Ile	Asn	Leu	Gly	Met	Pro	Leu	Ser	Cys	Phe	Pro	Glu	Gly
				140					145					150
Gly	Gln	Val	Val	Ile	Thr	Phe	Ser	Gln	Ser	Lys	Ser	Lys	Gln	Lys
				155					160					165
Glu	Asp	Asn	His	Ile	Phe	Gly	Arg	Gln	Asp	Lys	Ala	Ser	Thr	Glu
				170					175					180
Cys	Val	Lys	Phe	Tyr	Ile	His	Ala	Ile	Gly	Ile	Gly	Lys	Cys	Lys
				185					190					195
Arg	Arg	Ile	Val	Lys	Cys	Gly	Lys	Leu	His	Lys	Lys	Gly	Arg	Lys
				200					205					210
Leu	Cys	Val	Tyr	Ala	Phe	Lys	Gly	Glu	Thr	Ile	Lys	Asp	Ala	Leu
				215					220					225
Cys	Lys	Asp	Gly	Arg	Phe	Leu	Ser	Phe	Leu	Glu	Asn	Asp	Asp	Trp
				230					235					240
Lys	Leu	Ile	Glu	Asn	Asn	Asp	Thr	Ile	Leu	Glu	Ser	Thr	Gln	Pro
				245					250					255
Val	Asp	Glu	Leu	Glu	Gly	Arg	Tyr	Phe	Gln	Val	Glu	Val	Glu	Lys
				260					265					270
Arg	Met	Val	Pro	Ser	Ala	Ala	Ala	Ser	Gln	Asn	Pro	Glu	Ser	Glu
				275					280					285
Lys	Arg	Asn	Thr	Cys	Val	Leu	Arg	Glu	Gln	Ile	Val	Ala	Gln	Tyr
				290					295					300
Pro	Ser	Leu	Lys	Arg	Glu	Ser	Glu	Lys	Ile	Ile	Glu	Asn	Phe	Lys
				305					310					315
Lys	Lys	Met	Lys	Val	Lys	Asn	Gly	Glu	Thr	Leu	Phe	Glu	Leu	His
				320					325					330
Arg	Thr	Thr	Phe	Gly	Lys	Val	Thr	Lys	Asn	Ser	Ser	Ser	Ile	Lys
				335					340					345
Val	Val	Lys	Leu	Leu	Val	Arg	Leu	Ser	Asp	Ser	Val	Gly	Tyr	Leu
				350					355					360
Phe	Trp	Asp	Ser	Ala	Thr	Thr	Gly	Tyr	Ala	Thr	Cys	Phe	Val	Phe
				365					370					375
Lys	Gly	Leu	Phe	Ile	Leu	Thr	Cys	Arg	His	Val	Ile	Asp	Ser	Ile
				380					385					390
Val	Gly	Asp	Gly	Ile	Glu	Pro	Ser	Lys	Trp	Ala	Thr	Ile	Ile	Gly
				395					400					405
Gln	Cys	Val	Arg	Val	Thr	Phe	Gly	Tyr	Glu	Glu	Leu	Lys	Asp	Lys
				410					415					420
Glu	Thr	Asn	Tyr	Phe	Phe	Val	Glu	Pro	Trp	Phe	Glu	Ile	His	Asn
				425					430					435
Glu	Glu	Leu	Asp	Tyr	Ala	Val	Leu	Lys	Leu	Lys	Glu	Asn	Gly	Gln
				440					445					450
Gln	Val	Pro	Met	Glu	Leu	Tyr	Asn	Gly	Ile	Thr	Pro	Val	Pro	Leu
				455					460					465

Ser Gly Leu Ile	His Ile Ile Gly His	Pro Tyr Gly Glu Lys Lys	
	470	475	480
Gln Ile Asp Ala	Cys Ala Val Ile Pro	Gln Gly Gln Arg Ala Lys	
	485	490	495
Lys Cys Gln Glu	Arg Val Gln Ser Lys	Lys Ala Glu Ser Pro Glu	
	500	505	510
Tyr Val His Met	Tyr Thr Gln Arg Ser	Phe Gln Lys Ile Val His	
	515	520	525
Asn Pro Asp Val	Ile Thr Tyr Asp Thr	Glu Phe Phe Phe Gly Ala	
	530	535	540
Ser Gly Ser Pro	Val Phe Asp Ser Lys	Gly Ser Leu Val Ala Met	
	545	550	555
His Ala Ala Gly	Phe Ala Tyr Thr Tyr	Gln Asn Glu Thr Arg Ser	
	560	565	570
Ile Ile Glu Phe	Gly Ser Thr Met Glu	Ser Ile Leu Leu Asp Ile	
	575	580	585
Lys Gln Arg His	Lys Pro Trp Tyr Glu	Glu Val Phe Val Asn Gln	
	590	595	600
Gln Asp Val Glu	Met Met Ser Asp Glu	Asp Leu	
	605	610	

<210> 10

<211> 147

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7486358CD1

<400> 10

Met Trp Ser Leu Pro	Pro Ser Arg Ala Leu	Ser Cys Ala Pro Leu	
1	5	10	15
Leu Leu Leu Phe Ser	Phe Gln Phe Leu Val	Thr Tyr Ala Trp Arg	
	20	25	30
Phe Gln Glu Glu Glu	Glu Trp Asn Asp Gln	Lys Gln Ile Ala Val	
	35	40	45
Tyr Leu Pro Pro Thr	Leu Glu Phe Ala Val	Tyr Thr Phe Asn Lys	
	50	55	60
Gln Ser Lys Asp Trp	Tyr Ala Tyr Lys Leu	Val Pro Val Leu Ala	
	65	70	75
Ser Trp Lys Glu Gln	Gly Tyr Asp Lys Met	Thr Phe Ser Met Asn	
	80	85	90
Leu Gln Leu Gly Arg	Thr Met Cys Gly Lys	Phe Glu Asp Asp Ile	
	95	100	105
Asp Asn Cys Pro Phe	Gln Glu Ser Pro Glu	Leu Asn Asn Thr Cys	
	110	115	120
Thr Cys Phe Phe Thr	Ile Gly Ile Glu Pro	Trp Arg Thr Arg Phe	
	125	130	135
Asp Leu Trp Asn Lys	Thr Cys Ser Gly Gly	His Ser	
	140	145	

<210> 11

<211> 624

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472344CD1

<400> 11

Met Ala Leu Arg Ala Arg Ala Leu Tyr Asp Phe Arg Ser Glu Asn

1	5	10	15
Pro Gly Glu Ile Ser	Leu Arg Glu His	Glu Val Leu Ser Leu	Cys
20	25	30	
Ser Glu Gln Asp Ile	Glu Gly Trp Leu	Glu Gly Val Asn Ser	Arg
35	40	45	
Gly Asp Arg Gly Leu	Phe Pro Ala Ser	Tyr Val Gln Val Ile	Arg
50	55	60	
Ala Pro Glu Pro Gly	Pro Ala Gly Asp	Gly Gly Pro Gly Ala	Pro
65	70	75	
Ala Arg Tyr Ala Asn	Val Pro Pro Gly	Gly Phe Glu Pro Leu	Pro
80	85	90	
Val Ala Pro Pro Ala	Ser Phe Lys Pro	Pro Pro Asp Ala Phe	Gln
95	100	105	
Ala Leu Leu Gln Pro	Gln Gln Ala Pro	Pro Pro Ser Thr Phe	Gln
110	115	120	
Pro Pro Gly Ala Gly	Phe Pro Tyr Gly	Gly Gly Ala Leu Gln	Pro
125	130	135	
Ser Pro Gln Gln Leu	Tyr Gly Gly Tyr	Gln Ala Ser Gln Gly	Ser
140	145	150	
Asp Asp Asp Trp Asp	Asp Glu Trp Asp	Asp Ser Ser Thr Val	Ala
155	160	165	
Asp Glu Pro Gly Ala	Leu Gly Ser Gly	Ala Tyr Pro Asp Leu	Asp
170	175	180	
Gly Ser Ser Ser Ala	Gly Val Gly Ala	Ala Gly Arg Tyr Arg	Leu
185	190	195	
Ser Thr Arg Ser Asp	Leu Ser Leu Gly	Ser Arg Gly Gly Ser	Val
200	205	210	
Pro Pro Gln His His	Pro Ser Gly Pro	Lys Ser Ser Ala Thr	Val
215	220	225	
Ser Arg Asn Leu Asn	Arg Phe Ser Thr	Phe Val Lys Ser Gly	Gly
230	235	240	
Glu Ala Phe Val Leu	Gly Glu Ala Ser	Gly Phe Val Lys Asp	Gly
245	250	255	
Asp Lys Leu Cys Val	Val Leu Gly Pro	Tyr Gly Pro Glu Trp	Gln
260	265	270	
Glu Asn Pro Tyr Pro	Phe Gln Cys Thr	Ile Asp Asp Pro Thr	Lys
275	280	285	
Gln Thr Lys Phe Gly	Met Lys Ser Tyr	Ile Ser Tyr Lys Leu	
290	295	300	
Val Pro Thr His Thr	Gln Val Pro Val	His Arg Arg Tyr Lys	His
305	310	315	
Phe Asp Trp Leu Tyr	Ala Arg Leu Ala	Glu Lys Phe Pro Val	Ile
320	325	330	
Ser Val Pro His Leu	Pro Glu Lys Gln	Ala Thr Gly Arg Phe	Glu
335	340	345	
Glu Asp Phe Ile Ser	Lys Arg Arg Lys	Gly Leu Ile Trp Trp	Met
350	355	360	
Asn His Met Ala Ser	His Pro Val Leu	Ala Gln Cys Asp Val	Phe
365	370	375	
Gln His Phe Leu Thr	Cys Pro Ser Ser	Thr Asp Glu Lys Ala	Trp
380	385	390	
Lys Gln Gly Lys Arg	Lys Ala Glu Lys	Asp Glu Met Val Gly	Ala
395	400	405	
Asn Phe Phe Leu Thr	Leu Ser Thr Pro	Pro Ala Ala Ala Leu	Asp
410	415	420	
Leu Gln Glu Val Glu	Ser Lys Ile Asp	Gly Phe Lys Cys Phe	Thr
425	430	435	
Lys Lys Met Asp Asp	Ser Ala Leu Gln	Leu Asn His Thr Ala	Asn
440	445	450	
Glu Phe Ala Arg Lys	Gln Val Thr Gly	Phe Lys Lys Glu Tyr	Gln
455	460	465	
Lys Val Gly Gln Ser	Phe Arg Gly Leu	Ser Gln Ala Phe Glu	Leu
470	475	480	

Asp	Gln	Gln	Ala	Phe	Ser	Val	Gly	Leu	Asn	Gln	Ala	Ile	Ala	Phe	
				485					490					495	
Thr	Gly	Asp	Ala	Tyr	Asp	Ala	Ile	Gly	Glu	Leu	Phe	Ala	Glu	Gln	
				500					505					510	
Pro	Arg	Gln	Asp	Leu	Asp	Pro	Val	Met	Asp	Leu	Leu	Ala	Leu	Tyr	
				515					520					525	
Gln	Gly	His	Leu	Ala	Asn	Phe	Pro	Asp	Ile	Ile	His	Val	Gln	Lys	
				530					535					540	
Gly	Ala	Leu	Thr	Lys	Val	Lys	Glu	Ser	Arg	Arg	His	Val	Glu	Glu	
				545					550					555	
Gly	Lys	Met	Glu	Val	Gln	Lys	Ala	Asp	Gly	Ile	Gln	Asp	Arg	Cys	
				560					565					570	
Asn	Thr	Ile	Ser	Phe	Ala	Thr	Leu	Ala	Glu	Ile	His	His	Phe	His	
				575					580					585	
Gln	Ile	Arg	Val	Arg	Asp	Phe	Lys	Ser	Gln	Met	Gln	His	Phe	Leu	
				590					595					600	
Gln	Gln	Gln	Ile	Ile	Phe	Phe	Gln	Lys	Val	Thr	Gln	Lys	Leu	Glu	
				605					610					615	
Glu	Ala	Leu	His	Lys	Tyr	Asp	Ser	Val							
				620											

<210> 12

<211> 283

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7192959CD1

<400> 12

Met	Gly	Ala	Ser	Val	Ser	Arg	Gly	Arg	Ala	Ala	Arg	Val	Pro	Ala	
1				5					10					15	
Pro	Glu	Pro	Glu	Pro	Glu	Glu	Ala	Leu	Asp	Leu	Ser	Gln	Leu	Pro	
				20					25					30	
Pro	Glu	Leu	Leu	Leu	Val	Val	Leu	Ser	His	Val	Pro	Pro	Arg	Thr	
				35					40					45	
Leu	Leu	Gly	Arg	Cys	Arg	Gln	Val	Cys	Arg	Gly	Trp	Arg	Ala	Leu	
				50					55					60	
Val	Asp	Gly	Gln	Ala	Leu	Trp	Leu	Leu	Ile	Leu	Ala	Arg	Asp	His	
				65					70					75	
Gly	Ala	Thr	Gly	Arg	Ala	Leu	Leu	His	Leu	Ala	Arg	Ser	Cys	Gln	
				80					85					90	
Ser	Pro	Ala	Arg	Asn	Ala	Arg	Pro	Cys	Pro	Leu	Gly	Arg	Phe	Cys	
				95					100					105	
Ala	Arg	Arg	Pro	Ile	Gly	Arg	Asn	Leu	Ile	Arg	Asn	Pro	Cys	Gly	
				110					115					120	
Gln	Glu	Gly	Leu	Arg	Lys	Trp	Met	Val	Gln	His	Gly	Gly	Asp	Gly	
				125					130					135	
Trp	Val	Val	Glu	Glu	Asn	Arg	Thr	Thr	Val	Pro	Gly	Ala	Pro	Ser	
				140					145					150	
Gln	Thr	Cys	Phe	Val	Thr	Ser	Phe	Ser	Trp	Cys	Cys	Lys	Lys	Gln	
				155					160					165	
Val	Leu	Asp	Leu	Glu	Glu	Glu	Gly	Leu	Trp	Pro	Glu	Leu	Leu	Asp	
				170					175					180	
Ser	Gly	Arg	Ile	Glu	Ile	Cys	Val	Ser	Asp	Trp	Trp	Gly	Ala	Arg	
				185					190					195	
His	Asp	Ser	Gly	Cys	Met	Tyr	Arg	Leu	Leu	Val	Gln	Leu	Leu	Asp	
				200					205					210	
Ala	Asn	Gln	Thr	Val	Leu	Asp	Lys	Phe	Ser	Ala	Val	Pro	Asp	Pro	
				215					220					225	
Ile	Pro	Gln	Trp	Asn	Asn	Asn	Ala	Cys	Leu	His	Val	Thr	His	Val	
				230					235					240	

Phe	Ser	Asn	Ile	Lys	Met	Gly	Val	Arg	Phe	Val	Ser	Phe	Glu	His
				245					250					255
Arg	Gly	Gln	Asp	Thr	Gln	Phe	Trp	Ala	Gly	His	Tyr	Gly	Ala	Arg
				260					265					270
Val	Thr	Asn	Ser	Ser	Val	Ile	Val	Arg	Val	Arg	Leu	Ser		
				275					280					

<210> 13

<211> 142

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6169565CD1

<400> 13

Met	Ala	Asn	Gly	Lys	Lys	Phe	Leu	His	Leu	Pro	Leu	Leu	Thr	Ser
1				5					10					15
Phe	Pro	Leu	Pro	Arg	Leu	Phe	Pro	Thr	Pro	Leu	Ile	Cys	Ser	Leu
				20					25					30
Ile	Ser	Val	Val	Gly	Ile	Lys	Gly	Ile	Gln	Lys	Thr	Pro	Leu	Gln
				35					40					45
Thr	Leu	Pro	Leu	Tyr	Cys	Ser	Phe	Arg	Asp	Val	Thr	Leu	Ile	His
				50					55					60
Cys	Phe	Leu	Leu	Ile	Pro	His	Cys	Pro	Met	Pro	Leu	Leu	Ser	Arg
				65					70					75
Asp	Leu	Leu	His	Lys	Leu	Arg	Gly	Phe	Leu	His	Leu	Trp	Ala	Leu
				80					85					90
Gly	Gln	Ser	His	Pro	Tyr	Leu	Phe	Leu	Cys	Gln	Glu	Pro	Lys	Phe
				95					100					105
Ser	Leu	Pro	Glu	Val	Lys	Glu	Pro	Thr	Pro	Asp	Leu	Ser	Ile	Ile
				110					115					120
Thr	Gln	Thr	Asn	Pro	Ile	Val	Trp	Ser	Thr	Gln	Ile	Leu	Gln	Ser
				125					130					135
Trp	Arg	Pro	Thr	Thr	Pro	His								
				140										

<210> 14

<211> 354

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7494717CD1

<400> 14

Met	Ala	Glu	Ser	Pro	Thr	Glu	Glu	Ala	Ala	Thr	Ala	Gly	Ala	Gly
1				5					10					15
Ala	Ala	Gly	Pro	Gly	Ala	Ser	Ser	Val	Ala	Gly	Val	Val	Gly	Val
				20					25					30
Ser	Gly	Ser	Gly	Gly	Gly	Phe	Gly	Pro	Pro	Phe	Leu	Pro	Asp	Val
				35					40					45
Trp	Ala	Ala	Ala	Ala	Ala	Ala	Gly	Gly	Ala	Gly	Gly	Pro	Gly	Ser
				50					55					60
Gly	Leu	Ala	Pro	Leu	Pro	Gly	Leu	Pro	Pro	Ser	Ala	Ala	Ala	His
				65					70					75
Gly	Ala	Ala	Leu	Leu	Ser	His	Trp	Asp	Pro	Thr	Leu	Ser	Ser	Asp
				80					85					90
Trp	Asp	Gly	Glu	Arg	Thr	Ala	Pro	Gln	Cys	Leu	Leu	Arg	Ile	Lys
				95					100					105
Arg	Asp	Ile	Met	Ser	Ile	Tyr	Lys	Glu	Pro	Pro	Pro	Gly	Met	Phe

	110		115		120
Val Val Pro Asp Thr Val Asp Met Thr Lys Ile His Ala Leu Ile					
	125		130		135
Thr Gly Pro Phe Asp Thr Pro Tyr Glu Gly Gly Phe Phe Leu Phe					
	140		145		150
Val Phe Arg Cys Pro Pro Asp Tyr Pro Ile His Pro Pro Arg Val					
	155		160		165
Lys Leu Met Thr Thr Gly Asn Asn Thr Val Arg Phe Asn Pro Asn					
	170		175		180
Phe Tyr Arg Asn Gly Lys Val Cys Leu Ser Ile Leu Gly Thr Trp					
	185		190		195
Thr Gly Pro Ala Trp Ser Pro Ala Gln Ser Ile Ser Ser Val Leu					
	200		205		210
Ile Ser Ile Gln Ser Leu Met Thr Glu Asn Pro Tyr His Asn Glu					
	215		220		225
Pro Gly Phe Glu Gln Glu Arg His Pro Gly Asp Ser Lys Asn Tyr					
	230		235		240
Asn Glu Cys Ile Arg His Glu Thr Ile Arg Val Ala Val Cys Asp					
	245		250		255
Met Met Glu Gly Lys Cys Pro Cys Pro Glu Pro Leu Arg Gly Val					
	260		265		270
Met Glu Lys Ser Phe Leu Glu Tyr Tyr Asp Phe Tyr Glu Val Ala					
	275		280		285
Cys Lys Asp Arg Leu His Leu Gln Gly Gln Thr Met Gln Asp Pro					
	290		295		300
Phe Gly Glu Lys Arg Gly His Phe Asp Tyr Gln Ser Leu Leu Met					
	305		310		315
Arg Leu Gly Leu Ile Arg Gln Lys Val Leu Glu Arg Leu His Asn					
	320		325		330
Glu Asn Ala Glu Met Asp Ser Asp Ser Ser Ser Ser Gly Thr Glu					
	335		340		345
Thr Asp Leu His Gly Ser Leu Arg Val					
	350				

<210> 15

<211> 89

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7497510CD1

<400> 15

Met Gln Leu Gln Ala Ser Leu Ser Phe Leu Leu Ile Leu Thr Leu					
1	5		10		15
Cys Leu Glu Leu Arg Ser Glu Leu Ala Arg Asp Thr Ile Lys Asp					
	20		25		30
Leu Leu Pro Asn Val Cys Ala Phe Pro Met Glu Lys Gly Pro Cys					
	35		40		45
Gln Thr Tyr Met Thr Arg Trp Phe Phe Asn Phe Glu Thr Gly Glu					
	50		55		60
Cys Glu Leu Phe Ala Tyr Gly Gly Cys Gly Gly Asn Ser Asn Asn					
	65		70		75
Phe Leu Arg Lys Glu Lys Cys Glu Lys Phe Cys Lys Phe Thr					
	80		85		

<210> 16

<211> 419

<212> PRT

<213> Homo sapiens

<220>

<223> Incyte ID No: 7498882CD1

Met	Gly	Ala	Gly	Pro	Ser	Leu	Leu	Leu	Ala	Ala	Leu	Leu	Leu	Leu
1				5					10					15
Leu	Ser	Gly	Asp	Gly	Ala	Val	Arg	Cys	Asp	Thr	Pro	Ala	Asn	Cys
				20					25					30
Thr	Tyr	Leu	Asp	Leu	Leu	Gly	Thr	Trp	Val	Phe	Gln	Val	Gly	Ser
				35					40					45
Ser	Gly	Ser	Gln	Arg	Asp	Val	Asn	Cys	Ser	Val	Met	Gly	Pro	Gln
				50					55					60
Glu	Lys	Lys	Val	Val	Val	Tyr	Leu	Gln	Lys	Leu	Asp	Thr	Ala	Tyr
				65					70					75
Asp	Asp	Leu	Gly	Asn	Ser	Gly	His	Phe	Thr	Ile	Ile	Tyr	Asn	Gln
				80					85					90
Gly	Phe	Glu	Ile	Val	Leu	Asn	Asp	Tyr	Lys	Trp	Phe	Ala	Phe	Phe
				95					100					105
Lys	Tyr	Lys	Glu	Glu	Gly	Ser	Lys	Val	Thr	Thr	Tyr	Cys	Asn	Glu
				110					115					120
Thr	Met	Thr	Gly	Trp	Val	His	Asp	Val	Leu	Gly	Arg	Asn	Trp	Ala
				125					130					135
Cys	Phe	Thr	Gly	Lys	Lys	Val	Gly	Thr	Ala	Ser	Glu	Asn	Val	Tyr
				140					145					150
Val	Asn	Thr	Ala	His	Leu	Lys	Asn	Ser	Gln	Glu	Lys	Tyr	Ser	Asn
				155					160					165
Arg	Leu	Tyr	Lys	Tyr	Asp	His	Asn	Phe	Val	Lys	Ala	Ile	Asn	Ala
				170					175					180
Ile	Gln	Lys	Ser	Trp	Thr	Ala	Thr	Thr	Tyr	Met	Glu	Tyr	Glu	Thr
				185					190					195
Leu	Thr	Leu	Gly	Asp	Met	Ile	Arg	Arg	Ser	Gly	Gly	His	Ser	Arg
				200					205					210
Lys	Ile	Pro	Arg	Pro	Lys	Pro	Ala	Pro	Leu	Thr	Ala	Glu	Ile	Gln
				215					220					225
Gln	Lys	Ile	Leu	His	Leu	Pro	Thr	Ser	Trp	Asp	Trp	Arg	Asn	Val
				230					235					240
His	Gly	Ile	Asn	Phe	Val	Ser	Pro	Val	Arg	Asn	Gln	Gly	Cys	Glu
				245					250					255
Gly	Gly	Phe	Pro	Tyr	Leu	Ile	Ala	Gly	Lys	Tyr	Ala	Gln	Asp	Phe
				260					265					270
Gly	Leu	Val	Glu	Glu	Ala	Cys	Phe	Pro	Tyr	Thr	Gly	Thr	Asp	Ser
				275					280					285
Pro	Cys	Lys	Met	Lys	Glu	Asp	Cys	Phe	Arg	Tyr	Tyr	Ser	Ser	Glu
				290					295					300
Tyr	His	Tyr	Val	Gly	Gly	Phe	Tyr	Gly	Gly	Cys	Asn	Glu	Ala	Leu
				305					310					315
Met	Lys	Leu	Glu	Leu	Val	His	His	Gly	Pro	Met	Ala	Val	Ala	Phe
				320					325					330
Glu	Val	Tyr	Asp	Asp	Phe	Leu	His	Tyr	Lys	Lys	Gly	Ile	Tyr	His
				335					340					345
His	Thr	Gly	Leu	Arg	Asp	Pro	Phe	Asn	Pro	Phe	Glu	Leu	Thr	Asn
		</												

<211> 951

	410		415		420
Phe Thr Cys Pro	Leu Cys Arg Gln Phe	Ala Asn Ser Val Leu	Pro		
	425		430		435
Cys Tyr Pro Gly	Ser Asn Val Glu Asn	Pro Trp Gln Arg	Pro		
	440		445		450
Ser Asn Lys Ser	Ile Gln Asp Leu Ile	Lys Glu Val Glu Glu	Leu		
	455		460		465
Gln Gly Arg Pro	Gly Ala Phe Pro Ser	Glu Thr Asn Leu Ser	Lys		
	470		475		480
Glu Met Glu Ser	Val Met Lys Asp Ile	Lys Asn Thr Thr Gln	Lys		
	485		490		495
Lys Tyr Arg Asp	Tyr Ser Lys Thr Pro	Gly Ser Pro Asp Asn	Asp		
	500		505		510
Phe Leu Phe Met	Tyr Ser Val Ala Arg	Thr Asn Leu Glu Leu	Glu		
	515		520		525
Leu Ile His Arg	Gly Gly Asn Leu Cys	Ser Gly Gly Ala Ser	Thr		
	530		535		540
Ala Gly Lys Arg	Ser Cys Leu Asn Gln	Leu Phe His Val Leu	Ala		
	545		550		555
Leu His Met Arg	Leu Tyr Ser Ile Asp	Ser Glu Tyr Asn Pro	Trp		
	560		565		570
Arg Lys Leu Thr	Gln Leu Glu Glu Met	Asn Pro Gln Leu Gly	Tyr		
	575		580		585
Glu Glu Gln Gln	Pro Glu Val Pro Ile	Leu Tyr His Asp Val	Thr		
	590		595		600
Ser Leu Leu Leu	Ile Gln Ile Leu Met	Met Pro Gln Pro Leu	Arg		
	605		610		615
Lys Asp His Phe	Thr Cys Ile Val Lys	Val Leu Phe Thr Leu	Leu		
	620		625		630
Tyr Thr Gln Ala	Leu Ala Ala Leu Ser	Val Lys Cys Ser Glu	Glu		
	635		640		645
Asp Arg Ser Ala	Trp Lys His Ala Gly	Ala Leu Lys Lys Ser	Thr		
	650		655		660
Cys Asp Ala Glu	Lys Ser Tyr Glu Val	Leu Leu Ser Phe Val	Ile		
	665		670		675
Ser Glu Leu Phe	Lys Gly Lys Leu Tyr	His Glu Glu Gly Thr	Gln		
	680		685		690
Glu Cys Ala Met	Val Asn Pro Ile Ala	Trp Ser Pro Glu Ser	Met		
	695		700		705
Glu Lys Cys Leu	Gln Asp Phe Cys Leu	Pro Phe Leu Arg Ile	Thr		
	710		715		720
Ser Leu Leu Gln	His His Leu Phe Gly	Glu Asp Leu Pro Ser	Cys		
	725		730		735
Gln Glu Glu Glu	Glu Phe Ser Val Leu	Ala Ser Cys Leu Gly	Leu		
	740		745		750
Leu Pro Thr Phe	Tyr Gln Thr Glu His	Pro Phe Ile Ser Ala	Ser		
	755		760		765
Cys Leu Asp Trp	Pro Val Pro Ala Phe	Asp Ile Ile Thr Gln	Trp		
	770		775		780
Cys Phe Glu Ile	Lys Ser Phe Thr Glu	Arg His Ala Glu Gln	Gly		
	785		790		795
Lys Ala Leu Leu	Ile Gln Glu Ser Lys	Trp Lys Leu Pro His	Leu		
	800		805		810
Leu Gln Leu Pro	Glu Asn Tyr Asn Thr	Ile Phe Gln Tyr Tyr	His		
	815		820		825
Arg Lys Thr Cys	Ser Val Cys Thr Lys	Val Pro Lys Asp Pro	Ala		
	830		835		840
Val Cys Leu Val	Cys Gly Thr Phe Val	Cys Leu Lys Gly Leu	Cys		
	845		850		855
Cys Lys Gln Gln	Ser Tyr Cys Glu Cys	Val Leu His Ser Gln	Asn		
	860		865		870
Cys Gly Ala Gly	Thr Gly Ile Phe Leu	Leu Ile Asn Ala Ser	Val		
	875		880		885

Ile	Ile	Ile	Ile	Arg	Gly	His	Arg	Phe	Cys	Leu	Trp	Gly	Ser	Val	
				890					895					900	
Tyr	Leu	Asp	Ala	His	Gly	Glu	Glu	Asp	Arg	Asp	Leu	Arg	Arg	Gly	
				905					910					915	
Lys	Pro	Leu	Tyr	Ile	Cys	Lys	Glu	Arg	Tyr	Lys	Val	Leu	Glu	Gln	
				920					925					930	
Gln	Trp	Ile	Ser	His	Thr	Phe	Asp	His	Ile	Asn	Lys	Arg	Trp	Gly	
				935					940					945	
Pro	His	Tyr	Asn	Gly	Leu										
				950											

<210> .18

<211> 668

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7102342CD1

<400> 18

Met	Phe	Arg	Leu	Trp	Leu	Leu	Leu	Ala	Gly	Leu	Cys	Gly	Leu	Leu	
1				5					10					15	
Ala	Ser	Arg	Pro	Gly	Phe	Gln	Asn	Ser	Leu	Leu	Gln	Ile	Val	Ile	
				20					25					30	
Pro	Glu	Lys	Ile	Gln	Thr	Asn	Thr	Asn	Asp	Ser	Ser	Glu	Ile	Glu	
				35					40					45	
Tyr	Glu	Gln	Ile	Ser	Tyr	Ile	Ile	Pro	Ile	Asp	Glu	Lys	Leu	Tyr	
				50					55					60	
Thr	Val	His	Leu	Lys	Gln	Arg	Tyr	Phe	Leu	Ala	Asp	Asn	Phe	Met	
				65					70					75	
Ile	Tyr	Leu	Tyr	Asn	Gln	Gly	Ser	Met	Asn	Thr	Tyr	Ser	Ser	Asp	
				80					85					90	
Ile	Gln	Thr	Gln	Cys	Tyr	Tyr	Gln	Gly	Asn	Ile	Glu	Gly	Tyr	Pro	
				95					100					105	
Asp	Ser	Met	Val	Thr	Leu	Ser	Thr	Cys	Ser	Gly	Leu	Arg	Gly	Ile	
				110					115					120	
Leu	Gln	Phe	Glu	Asn	Val	Ser	Tyr	Gly	Ile	Glu	Pro	Leu	Glu	Ser	
				125					130					135	
Ala	Val	Glu	Phe	Gln	His	Val	Leu	Tyr	Lys	Leu	Lys	Asn	Glu	Asp	
				140					145					150	
Asn	Asp	Ile	Ala	Ile	Phe	Ile	Asp	Arg	Ser	Leu	Lys	Glu	Gln	Pro	
				155					160					165	
Met	Asp	Asp	Asn	Ile	Phe	Ile	Ser	Glu	Lys	Ser	Glu	Pro	Ala	Val	
				170					175					180	
Pro	Asp	Leu	Phe	Pro	Leu	Tyr	Leu	Glu	Met	His	Ile	Val	Val	Asp	
				185					190					195	
Lys	Thr	Leu	Tyr	Asp	Tyr	Trp	Gly	Ser	Asp	Ser	Met	Ile	Val	Thr	
				200					205					210	
Asn	Lys	Val	Ile	Glu	Ile	Val	Gly	Leu	Ala	Asn	Ser	Met	Phe	Thr	
				215					220					225	
Gln	Phe	Lys	Val	Thr	Ile	Val	Leu	Ser	Ser	Leu	Glu	Leu	Trp	Ser	
				230					235					240	
Asp	Glu	Asn	Lys	Ile	Ser	Thr	Val	Gly	Glu	Ala	Asp	Glu	Leu	Leu	
				245					250					255	
Gln	Lys	Phe	Leu	Glu	Trp	Lys	Gln	Ser	Tyr	Leu	Asn	Leu	Arg	Pro	
				260					265					270	
His	Asp	Ile	Ala	Tyr	Leu	Leu	Ile	Tyr	Met	Asp	Tyr	Pro	Arg	Tyr	
				275					280					285	
Leu	Gly	Ala	Val	Phe	Pro	Gly	Thr	Met	Cys	Ile	Thr	Arg	Tyr	Ser	
				290					295					300	
Ala	Gly	Val	Ala	Leu	Gln	Cys	Gly	Pro	Ala	Ser	Cys	Cys	Asp	Phe	
				305					310					315	

```

Arg Thr Cys Val Leu Lys Asp Gly Ala Lys Cys Tyr Lys Gly Leu
320 325 330
Cys Cys Lys Asp Cys Gln Ile Leu Gln Ser Gly Val Glu Cys Arg
335 340 345
Pro Lys Ala His Pro Glu Cys Asp Ile Ala Glu Asn Cys Asn Gly
350 355 360
Ser Ser Pro Glu Cys Gly Pro Asp Ile Thr Leu Ile Asn Gly Leu
365 370 375
Ser Cys Lys Asn Asn Lys Phe Ile Cys Tyr Asp Gly Asp Cys His
380 385 390
Asp Leu Asp Ala Arg Cys Glu Ser Val Phe Gly Lys Gly Ser Arg
395 400 405
Asn Ala Pro Phe Ala Cys Tyr Glu Glu Ile Gln Ser Gln Ser Asp
410 415 420
Arg Phe Gly Asn Cys Gly Arg Asp Arg Asn Asn Lys Tyr Val Phe
425 430 435
Cys Gly Trp Arg Asn Leu Ile Cys Gly Arg Leu Val Cys Thr Tyr
440 445 450
Pro Thr Arg Lys Pro Phe His Gln Glu Asn Gly Asp Val Ile Tyr
455 460 465
Ala Phe Val Arg Asp Ser Val Cys Ile Thr Val Asp Tyr Lys Leu
470 475 480
Pro Arg Thr Val Pro Asp Pro Leu Ala Val Lys Asn Gly Ser Gln
485 490 495
Cys Asp Ile Gly Arg Val Cys Val Asn Arg Glu Cys Val Glu Ser
500 505 510
Arg Ile Ile Lys Ala Ser Ala His Val Cys Ser Gln Gln Cys Ser
515 520 525
Gly His Gly Val Cys Asp Ser Arg Asn Lys Cys His Cys Ser Pro
530 535 540
Gly Tyr Lys Pro Pro Asn Cys Gln Ile Arg Ser Lys Gly Phe Ser
545 550 555
Ile Phe Pro Glu Glu Asp Met Gly Ser Ile Met Glu Arg Ala Ser
560 565 570
Gly Lys Thr Glu Asn Thr Trp Leu Leu Gly Phe Leu Ile Ala Leu
575 580 585
Pro Ile Leu Ile Val Thr Thr Ala Ile Val Leu Ala Arg Lys Gln
590 595 600
Leu Lys Lys Trp Phe Ala Lys Glu Glu Glu Phe Pro Ser Ser Glu
605 610 615
Ser Lys Ser Glu Gly Ser Thr Gln Thr Tyr Ala Ser Gln Ser Ser
620 625 630
Ser Glu Gly Ser Thr Gln Thr Tyr Ala Ser Gln Thr Arg Ser Glu
635 640 645
Ser Ser Ser Gln Ala Asp Thr Ser Lys Ser Lys Ser Gln Asp Ser
650 655 660
Thr Gln Thr Gln Ser Ser Ser Asn
665

```

<210> 19

<211> 206

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4169939CD1

<400> 19

```

Met Met Leu Arg Leu Leu Ser Ser Leu Leu Leu Val Ala Val Ala
1 5 10 15
Ser Gly Tyr Gly Pro Pro Ser Ser His Ser Ser Ser Arg Val Val
20 25 30

```

```

His Gly Glu Asp Ala Ile Pro Ile Asn Ser Glu Glu Leu Phe Val
      35                      40                      45
His Pro Leu Trp Asn Arg Ser Cys Val Ala Cys Gly Asn Asp Ile
      50                      55                      60
Ala Leu Ile Lys Ser Arg Ser Ala Gln Leu Gly Asp Ala Val
      65                      70                      75
Gln Leu Ala Ser Leu Pro Pro Ala Gly Asp Ile Leu Pro Asn Lys
      80                      85                      90
Thr Pro Cys Tyr Ile Thr Gly Trp Gly Arg Leu Tyr Thr Asn Gly
      95                      100                     105
Pro Leu Pro Asp Lys Leu Gln Gln Ala Arg Leu Pro Val Val Asp
     110                      115                     120
Tyr Lys His Cys Ser Arg Trp Asn Trp Trp Gly Ser Thr Val Lys
     125                      130                     135
Lys Thr Met Val Cys Ala Gly Gly Tyr Ile Arg Ser Gly Cys Asn
     140                      145                     150
Gly Asp Ser Gly Gly Pro Leu Asn Cys Pro Thr Glu Asp Gly Gly
     155                      160                     165
Trp Gln Val His Gly Val Thr Ser Phe Val Ser Gly Phe Gly Cys
     170                      175                     180
Asn Phe Ile Trp Lys Pro Thr Val Phe Thr Arg Val Ser Ala Phe
     185                      190                     195
Ile Asp Trp Ile Glu Glu Thr Ile Ala Ser His
     200                      205

```

<210> 20

<211> 267

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6539977CD1

<400> 20

```

Met Ser Leu Arg Val Leu Gly Ser Gly Thr Trp Pro Ser Ala Pro
  1      5      10
Lys Met Phe Leu Leu Leu Thr Ala Leu Gln Val Leu Ala Ile Ala
      20      25      30
Met Thr Arg Ser Gln Glu Asp Glu Asn Lys Ile Ile Gly Gly Tyr
      35      40      45
Thr Cys Thr Arg Ser Ser Gln Pro Trp Gln Ala Ala Leu Leu Ala
      50      55      60
Gly Pro Arg Arg Arg Phe Leu Cys Gly Gly Ala Leu Leu Ser Gly
      65      70      75
Gln Trp Val Ile Thr Ala Ala His Cys Gly Arg Pro Ile Leu Gln
      80      85      90
Val Ala Leu Gly Lys His Asn Leu Arg Arg Trp Glu Ala Thr Gln
      95     100     105
Gln Val Leu Arg Val Val Arg Gln Val Thr His Pro Asn Tyr Asn
     110     115     120
Ser Arg Thr His Asp Asn Asp Leu Met Leu Leu Gln Leu Gln
     125     130     135
Pro Ala Arg Ile Gly Arg Ala Val Arg Pro Ile Glu Val Thr Gln
     140     145     150
Ala Cys Ala Ser Pro Gly Thr Ser Cys Arg Val Ser Gly Trp Gly
     155     160     165
Thr Ile Ser Ser Pro Ile Ala Arg Tyr Pro Ala Ser Leu Gln Cys
     170     175     180
Val Asn Ile Asn Ile Ser Pro Asp Glu Val Cys Gln Lys Ala Tyr
     185     190     195
Pro Arg Thr Ile Thr Pro Gly Met Val Cys Ala Gly Val Pro Gln
     200     205     210

```

Gly	Gly	Lys	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Val	
				215					220					225	
Cys	Arg	Gly	Gln	Leu	Gln	Gly	Leu	Val	Ser	Trp	Gly	Met	Glu	Arg	
				230					235					240	
Cys	Ala	Leu	Pro	Gly	Tyr	Pro	Gly	Val	Tyr	Thr	Asn	Leu	Cys	Lys	
				245					250					255	
Tyr	Arg	Ser	Trp	Ile	Glu	Glu	Thr	Met	Arg	Asp	Lys				
				260					265						

<210> 21
 <211> 86
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7675588CD1

Met	Gly	Leu	Ser	Gly	Leu	Leu	Pro	Ile	Leu	Val	Pro	Phe	Ile	Leu	
1				5					10					15	
Leu	Gly	Asp	Ile	Gln	Glu	Pro	Gly	His	Ala	Glu	Gly	Ile	Leu	Gly	
				20					25					30	
Lys	Pro	Cys	Pro	Lys	Ile	Lys	Val	Glu	Cys	Glu	Val	Glu	Glu	Ile	
				35					40					45	
Asp	Gln	Cys	Thr	Lys	Pro	Arg	Asp	Cys	Pro	Glu	Asn	Met	Lys	Cys	
				50					55					60	
Cys	Pro	Phe	Ser	Arg	Gly	Lys	Lys	Cys	Leu	Asp	Phe	Arg	Lys	Val	
				65					70					75	
Ser	Leu	Thr	Leu	Tyr	His	Lys	Glu	Glu	Leu	Glu					
				80					85						

<210> 22
 <211> 232
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6244077CD1

Met	Ala	Asn	Tyr	Tyr	Glu	Val	Leu	Gly	Val	Gln	Ala	Ser	Ala	Ser	
1				5					10					15	
Pro	Glu	Asp	Ile	Lys	Lys	Ala	Tyr	Arg	Lys	Leu	Ala	Leu	Arg	Trp	
				20					25					30	
His	Pro	Asp	Lys	Asn	Pro	Asp	Asn	Lys	Glu	Glu	Ala	Glu	Lys	Lys	
				35					40					45	
Phe	Lys	Leu	Val	Ser	Glu	Ala	Tyr	Glu	Val	Leu	Ser	Asp	Ser	Lys	
				50					55					60	
Lys	Arg	Ser	Leu	Tyr	Asp	Arg	Ala	Gly	Cys	Asp	Ser	Trp	Arg	Ala	
				65					70					75	
Gly	Gly	Gly	Ala	Ser	Thr	Pro	Tyr	His	Ser	Pro	Phe	Asp	Thr	Gly	
				80					85					90	
Tyr	Thr	Phe	Arg	Asn	Pro	Glu	Asp	Ile	Phe	Arg	Glu	Phe	Phe	Gly	
				95					100					105	
Gly	Leu	Asp	Pro	Phe	Ser	Phe	Glu	Phe	Trp	Asp	Ser	Pro	Phe	Asn	
				110					115					120	
Ser	Asp	Arg	Gly	Gly	Arg	Gly	His	Gly	Leu	Arg	Gly	Ala	Phe	Ser	
				125					130					135	
Ala	Gly	Phe	Gly	Glu	Phe	Pro	Ala	Phe	Met	Glu	Ala	Phe	Ser	Ser	
				140					145					150	
Phe	Asn	Met	Leu	Gly	Cys	Ser	Gly	Gly	Ser	His	Thr	Thr	Phe	Ser	

	155		160		165
Ser Thr Ser Phe	Gly Gly Ser Ser Ser	Gly Ser Ser Gly Phe	Lys		
	170		175		180
Ser Val Met Ser	Ser Thr Glu Met Ile	Asn Gly His Lys Val	Thr		
	185		190		195
Thr Lys Arg Ile	Val Glu Asn Gly Gln	Glu Arg Val Glu Val	Glu		
	200		205		210
Glu Asp Gly Gln	Leu Lys Ser Val Thr	Val Asn Gly Lys Glu	Gln		
	215		220		225
Leu Lys Trp Met	Asp Ser Lys				
	230				

<210> 23
 <211> 237
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7498404CD1

<400> 23	
Met Ala Ser Leu Glu Val Ser Arg Ser Pro Arg Arg Ser Arg Arg	
1 5 10 15	
Glu Leu Glu Val Arg Ser Pro Arg Gln Asn Lys Tyr Ser Val Leu	
20 25 30	
Leu Pro Thr Tyr Asn Glu Arg Glu Asn Leu Pro Leu Ile Val Trp	
35 40 45	
Leu Leu Val Lys Ser Phe Ser Glu Ser Gly Ile Asn Tyr Glu Ile	
50 55 60	
Ile Ile Ile Asp Asp Gly Ser Pro Asp Gly Thr Arg Asp Val Ala	
65 70 75	
Glu Gln Leu Glu Lys Ile Tyr Gly Ser Asp Arg Ile Leu Leu Arg	
80 85 90	
Pro Arg Glu Lys Lys Leu Gly Leu Gly Thr Ala Tyr Ile His Gly	
95 100 105	
Met Lys His Ala Thr Gly Asn Tyr Ile Ile Ile Met Asp Ala Asp	
110 115 120	
Leu Ser His His Pro Lys Phe Ile Pro Glu Phe Ile Arg Lys Gln	
125 130 135	
Lys Glu Gly Asn Phe Asp Ile Val Ser Gly Thr Arg Tyr Lys Gly	
140 145 150	
Asn Gly Gly Val Tyr Gly Trp Asp Leu Lys Arg Lys Ile Ile Arg	
155 160 165	
Leu Tyr Arg Lys Glu Val Leu Glu Lys Leu Ile Glu Lys Cys Val	
170 175 180	
Ser Lys Gly Tyr Val Phe Gln Met Glu Met Ile Val Arg Ala Arg	
185 190 195	
Gln Leu Asn Tyr Thr Ile Gly Glu Val Pro Ile Ser Phe Val Asp	
200 205 210	
Arg Val Tyr Gly Glu Ser Lys Leu Gly Gly Asn Glu Ile Val Ser	
215 220 225	
Phe Leu Lys Gly Leu Leu Thr Leu Phe Ala Thr Thr	
230 235	

<210> 24
 <211> 146
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7391748CD1

<400> 24

```

Met Leu Leu Gln Leu Ser Arg Arg Val Arg Arg Asn Arg Asn Val
 1          5          10          15
Asn Pro Val Ala Leu Pro Arg Ala Gln Glu Gly Leu Arg Pro Gly
          20          25          30
Thr Leu Cys Thr Val Ala Gly Trp Gly Arg Val Ser Met Arg Arg
          35          40          45
Gly Thr Asp Thr Leu Arg Glu Val Gln Leu Arg Val Gln Arg Asp
          50          55          60
Arg Gln Cys Leu Arg Ile Phe Gly Ser Tyr Asp Pro Arg Arg Gln
          65          70          75
Ile Cys Val Gly Asp Arg Arg Glu Arg Lys Ala Ala Phe Lys Gly
          80          85          90
Asp Ser Gly Gly Pro Leu Leu Cys Asn Asn Val Ala His Gly Ile
          95          100          105
Val Ser Tyr Gly Lys Ser Ser Gly Val Pro Pro Glu Val Phe Thr
          110          115          120
Arg Val Ser Ser Phe Leu Pro Trp Ile Arg Thr Thr Met Arg Ser
          125          130          135
Phe Lys Leu Leu Asp Gln Met Glu Thr Pro Leu
          140          145

```

<210> 25

<211> 696

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7499780CD1

<400> 25

```

Met Thr Ser Ser Gly Pro Gly Pro Arg Phe Leu Leu Leu Leu Pro
 1          5          10          15
Leu Leu Leu Pro Pro Ala Ala Ser Ala Ser Asp Arg Pro Arg Gly
          20          25          30
Arg Asp Pro Val Asn Pro Glu Lys Leu Leu Val Ile Thr Val Ala
          35          40          45
Thr Ala Glu Thr Glu Gly Tyr Leu Arg Phe Leu Arg Ser Ala Glu
          50          55          60
Phe Phe Asn Tyr Thr Val Arg Thr Leu Gly Leu Gly Glu Glu Trp
          65          70          75
Arg Gly Gly Asp Val Ala Arg Thr Val Gly Gly Gly Gln Lys Val
          80          85          90
Arg Trp Leu Lys Lys Glu Met Glu Lys Tyr Ala Asp Arg Glu Asp
          95          100          105
Met Ile Ile Met Phe Val Asp Ser Tyr Asp Val Ile Leu Ala Gly
          110          115          120
Ser Pro Thr Glu Leu Leu Lys Lys Phe Val Gln Ser Gly Ser Arg
          125          130          135
Leu Leu Phe Ser Ala Glu Ser Phe Cys Trp Pro Glu Trp Gly Leu
          140          145          150
Ala Glu Gln Tyr Pro Glu Val Gly Thr Gly Lys Arg Phe Leu Asn
          155          160          165
Ser Gly Gly Phe Ile Gly Phe Ala Thr Thr Ile His Gln Ile Val
          170          175          180
Arg Gln Trp Lys Tyr Lys Asp Asp Asp Asp Asp Gln Leu Phe Tyr
          185          190          195
Thr Arg Leu Tyr Leu Asp Pro Gly Leu Arg Glu Lys Leu Ser Leu
          200          205          210
Asn Leu Asp His Lys Ser Arg Ile Phe Gln Asn Leu Asn Gly Ala
          215          220          225
Leu Asp Glu Val Val Leu Lys Phe Asp Arg Asn Arg Val Arg Ile

```

	230		235		240
Arg Asn Val Ala	Tyr Asp Thr Leu Pro	Ile Val Val His Gly	Asn		
	245		250		255
Gly Pro Thr Lys	Leu Gln Leu Asn Tyr	Leu Gly Asn Tyr Val	Pro		
	260		265		270
Asn Gly Trp Thr	Pro Glu Gly Gly Cys	Gly Phe Cys Asn Gln	Asp		
	275		280		285
Arg Arg Thr Leu	Pro Gly Gly Gln Glu	Val Phe His Glu Pro	His		
	290		295		300
Ile Ala Asp Ser	Trp Pro Gln Leu Gln	Asp His Phe Ser Ala	Val		
	305		310		315
Lys Leu Val Gly	Pro Glu Glu Ala Leu	Ser Pro Gly Glu Ala	Arg		
	320		325		330
Asp Met Ala Met	Asp Leu Cys Arg Gln	Asp Pro Glu Cys Glu	Phe		
	335		340		345
Tyr Phe Ser Leu	Asp Ala Asp Ala Val	Leu Thr Asn Leu Gln	Thr		
	350		355		360
Leu Arg Ile Leu	Ile Glu Glu Asn Arg	Lys Val Ile Ala Pro	Met		
	365		370		375
Leu Ser Arg His	Gly Lys Leu Trp Ser	Asn Phe Trp Gly Ala	Leu		
	380		385		390
Ser Pro Asp Glu	Tyr Tyr Ala Arg Ser	Glu Asp Tyr Val Glu	Leu		
	395		400		405
Val Gln Arg Lys	Arg Val Gly Val Trp	Asn Val Pro Tyr Ile	Ser		
	410		415		420
Gln Ala Tyr Val	Ile Arg Gly Asp Thr	Leu Arg Met Glu Leu	Pro		
	425		430		435
Gln Arg Asp Val	Phe Ser Gly Ser Asp	Thr Asp Pro Asp Met	Ala		
	440		445		450
Phe Cys Lys Ser	Phe Arg Asp Lys Gly	Ile Phe Leu His Leu	Ser		
	455		460		465
Asn Gln His Glu	Phe Gly Arg Leu Leu	Ala Thr Ser Arg Tyr	Asp		
	470		475		480
Thr Glu His Leu	His Pro Asp Leu Trp	Gln Ile Phe Asp Asn	Pro		
	485		490		495
Val Asp Trp Lys	Glu Gln Tyr Ile His	Glu Asn Tyr Ser Arg	Ala		
	500		505		510
Leu Glu Gly Glu	Gly Ile Val Glu Gln	Pro Cys Pro Asp Val	Tyr		
	515		520		525
Trp Phe Pro Leu	Leu Ser Glu Gln Met	Cys Asp Glu Leu Val	Ala		
	530		535		540
Glu Met Glu His	Tyr Gly Gln Trp Ser	Gly Gly Arg His Glu	Asp		
	545		550		555
Ser Arg Leu Ala	Gly Gly Tyr Glu Asn	Val Pro Thr Val Asp	Ile		
	560		565		570
His Met Lys Gln	Val Gly Tyr Glu Asp	Gln Trp Leu Gln Leu	Leu		
	575		580		585
Arg Thr Tyr Val	Gly Pro Met Thr Glu	Ser Leu Phe Pro Gly	Tyr		
	590		595		600
His Thr Lys Ala	Arg Ala Val Met Asn	Phe Val Val Arg Tyr	Arg		
	605		610		615
Pro Asp Glu Gln	Pro Ser Leu Arg Pro	His His Asp Ser Ser	Thr		
	620		625		630
Phe Thr Leu Asn	Val Ala Leu Asn His	Lys Gly Leu Asp Tyr	Glu		
	635		640		645
Gly Gly Gly Cys	Arg Phe Leu Arg Tyr	Asp Cys Val Ile Ser	Ser		
	650		655		660
Pro Arg Lys Gly	Trp Ala Leu Leu His	Pro Gly Arg Leu Thr	His		
	665		670		675
Tyr His Glu Gly	Leu Pro Thr Thr Trp	Gly Thr Arg Tyr Ile	Met		
	680		685		690
Val Ser Phe Val	Asp Pro				
	695				

<210> 26
 <211> 630
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7499881CD1

<400> 26
 Met Ala Tyr Gln Ile Val Leu Glu Leu His Phe Ser His Cys Ala
 1 5 10 15
 Ala Met Gly Ala Ala Leu Leu Met Leu Ile Glu Asn Ala Leu Ile
 20 25 30
 Thr Gln Ser Arg Leu Met Leu Leu Glu Ser Val Leu Ile Phe Phe
 35 40 45
 Asn Leu Leu Ala Val Leu Ser Tyr Leu Lys Phe Phe Asn Cys Gln
 50 55 60
 Lys His Ser Pro Phe Ser Leu Ser Trp Trp Phe Trp Leu Thr Leu
 65 70 75
 Thr Gly Val Ala Cys Ser Cys Ala Val Gly Ile Lys Tyr Met Gly
 80 85 90
 Val Phe Thr Tyr Val Leu Val Leu Gly Val Ala Ala Val His Ala
 95 100 105
 Trp His Leu Leu Gly Asp Gln Thr Leu Ser Asn Val Gly Ala Asp
 110 115 120
 Val Gln Cys Cys Met Arg Pro Ala Cys Met Gly Gln Met Arg Met
 125 130 135
 Ser Gln Gly Val Cys Val Phe Cys His Leu Leu Ala Arg Ala Val
 140 145 150
 Ala Leu Leu Val Ile Pro Val Val Leu Tyr Leu Leu Phe Phe Tyr
 155 160 165
 Val His Leu Ile Leu Val Phe Arg Ser Gly Pro His Asp Gln Ile
 170 175 180
 Met Ser Ser Ala Phe Gln Ala Ser Leu Glu Gly Gly Leu Ala Arg
 185 190 195
 Ile Thr Gln Gly Gln Pro Leu Glu Val Ala Phe Gly Ser Gln Val
 200 205 210
 Thr Leu Arg Asn Val Phe Gly Lys Pro Val Pro Cys Trp Leu His
 215 220 225
 Ser His Gln Asp Thr Tyr Pro Met Ile Tyr Glu Asn Gly Arg Gly
 230 235 240
 Ser Ser His Gln Gln Gln Val Thr Cys Tyr Pro Phe Lys Asp Val
 245 250 255
 Asn Asn Trp Trp Ile Val Lys Asp Pro Arg Arg His Gln Leu Val
 260 265 270
 Val Ser Ser Pro Pro Arg Pro Val Arg His Gly Asp Met Val Gln
 275 280 285
 Leu Val His Gly Met Thr Thr Arg Ser Leu Asn Thr His Asp Val
 290 295 300
 Ala Ala Pro Leu Ser Pro His Ser Gln Glu Val Ser Cys Tyr Ile
 305 310 315
 Asp Tyr Asn Ile Ser Met Pro Ala Gln Asn Leu Trp Arg Leu Glu
 320 325 330
 Ile Val Asn Arg Gly Ser Asp Thr Asp Val Trp Lys Thr Ile Leu
 335 340 345
 Ser Glu Val Arg Phe Val His Val Asn Thr Ser Ala Val Leu Lys
 350 355 360
 Leu Ser Gly Ala His Leu Pro Asp Trp Gly Tyr Arg Gln Leu Glu
 365 370 375
 Ile Val Gly Glu Lys Leu Ser Arg Gly Tyr His Gly Ser Thr Val
 380 385 390
 Trp Asn Val Glu Glu His Arg Tyr Gly Ala Ser Gln Glu Gln Arg

	395		400		405
Glu Arg Glu Arg	Glu Leu His Ser Pro	Ala Gln Val Asp Val	Ser		
	410		415		420
Arg Asn Leu Ser	Phe Met Ala Arg Phe	Ser Glu Leu Gln Trp	Arg		
	425		430		435
Met Leu Ala Leu	Arg Ser Asp Asp Ser	Glu His Lys Tyr Ser	Ser		
	440		445		450
Ser Pro Leu Glu	Trp Val Thr Leu Asp	Thr Asn Ile Ala Tyr	Trp		
	455		460		465
Leu His Pro Arg	Thr Ser Ala Gln Ile	His Leu Leu Gly Asn	Ile		
	470		475		480
Val Ile Trp Val	Ser Gly Ser Leu Ala	Leu Ala Ile Tyr Ala	Leu		
	485		490		495
Leu Ser Leu Trp	Tyr Leu Leu Arg Arg	Arg Arg Asn Val His	Asp		
	500		505		510
Leu Pro Gln Asp	Ala Trp Leu Arg Trp	Val Leu Ala Gly Ala	Leu		
	515		520		525
Cys Ala Gly Gly	Trp Ala Val Asn Tyr	Leu Pro Phe Phe Leu	Met		
	530		535		540
Glu Lys Thr Leu	Phe Leu Tyr His Tyr	Leu Pro Ala Leu Thr	Phe		
	545		550		555
Gln Ile Leu Leu	Leu Pro Val Val Leu	Gln His Ile Ser Asp	His		
	560		565		570
Leu Cys Arg Ser	Gln Leu Gln Arg Ser	Ile Phe Ser Ala Leu	Val		
	575		580		585
Val Ala Trp Tyr	Ser Ser Ala Cys His	Val Ser Asn Thr Leu	Arg		
	590		595		600
Pro Leu Thr Tyr	Gly Asp Lys Ser Leu	Ser Pro His Glu Leu	Lys		
	605		610		615
Ala Leu Arg Trp	Lys Asp Ser Trp Asp	Ile Leu Ile Arg Lys	His		
	620		625		630

<210> 27

<211> 242

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7488579CD1

<400> 27

Met Leu Leu Leu Ala	Pro Gln Met Leu Asn	Leu Leu Leu Leu Ala	
1	5	10	15
Leu Pro Val Leu Ala	Ser Arg Ala Tyr	Ala Ala Pro Ala Pro	Gly
	20	25	30
Gln Ala Leu Gln Arg	Val Gly Ile Val	Gly Gly Gln Glu Ala	Pro
	35	40	45
Arg Ser Lys Trp Pro	Trp Gln Val Ser	Leu Arg Val His Gly	Pro
	50	55	60
Tyr Trp Met His Phe	Cys Gly Gly Ser	Leu Ile His Pro Gln	Trp
	65	70	75
Val Leu Thr Ala Ala	His Cys Val Gly	Pro Asp Val Lys Asp	Leu
	80	85	90
Ala Ala Leu Arg Val	Gln Leu Arg Glu	Gln His Leu Tyr Tyr	Gln
	95	100	105
Asp Gln Leu Leu Pro	Val Ser Arg Ile	Ile Val His Pro Gln	Phe
	110	115	120
Tyr Ile Ile Gln Thr	Gly Ala Asp Ile	Ala Leu Leu Glu Leu	Glu
	125	130	135
Glu Pro Val Asn Ile	Ser Ser His Ile	His Thr Val Thr Leu	Pro
	140	145	150

Pro Ala Ser Glu Thr Phe Pro Pro Gly Met Pro Cys Trp Val Thr
 155 160 165
 Gly Trp Gly Asp Val Asp Asn Asn Val His Leu Pro Pro Pro Tyr
 170 175 180
 Pro Leu Lys Glu Val Glu Val Pro Val Val Glu Asn His Leu Cys
 185 190 195
 Asn Ala Glu Tyr His Thr Gly Leu His Thr Gly His Ser Phe Gln
 200 205 210
 Ile Val Arg Asp Asp Met Leu Cys Ala Gly Ser Glu Asn His Asp
 215 220 225
 Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys Lys Val Asn
 230 235 240
 Gly Thr

<210> 28
 <211> 48
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7510521CD1

<400> 28
 Met Gly Ala Gly Pro Ser Leu Leu Leu Ala Ala Leu Leu Leu Leu
 1 5 10 15
 Leu Ser Gly Asp Gly Ala Val Arg Cys Asp Thr Pro Ala Asn Cys
 20 25 30
 Thr Tyr Leu Asp Leu Leu Gly Thr Trp Val Phe Gln Asp His Lys
 35 40 45
 Lys Lys Lys

<210> 29
 <211> 4384
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7994355CB1

<400> 29
 agaggcggcg gcggtgtct gtaggtggc gggtccggcag cagcccgcc cccggacgca 60
 ggacgtggcc ccaggcagcc ctgcagctc agtgcctag ccggggcaag ccgcgtctc 120
 cgctgtctgg acgggcccag gcgagatgta gggctctggg ccgggaggcc gccgggtggg 180
 cggctgatcg cggaggatcg cggagggcgc gccgaggatg gagagagcga tggagcaact 240
 caaccgcctg acgcgctcgc tgcgccgcgc gcgcaccgtg gagttgccg aggagatgag 300
 gtctcactat gttccccagg ctggtcttga actcctgggc tcaagtgatc cgcaccctgc 360
 tcccccaaaa gtgttgggat tacagatgtg agccaccgcg cctggcctgg ggataatctt 420
 ctgaaaaaaa tgataatgaa actgctgttt atacattaat gccaatggtt atggctgatc 480
 aacacaggtc tgtttctgaa ctactatcaa attcaaaatt tgatgtcaat tatgcattcg 540
 gacgtgtgaa aagaagcttg cttcacattg cagcaaatg tggatcggtg gaatgcttgg 600
 ttttgcgtgt aaagaaagga gcaaatccta actatcaaga tatttcaggc tgtacacccc 660
 ttcatttggc agcaagaaat ggtcatggtc agagagatac agcacagatc ctactattac 720
 gaggagccaa atatctgcca gataaaaaat gagtaactcc tctggattta tgtgtacagg 780
 gtggatatgg agagacttgt gaagtattaa ttcaatatca cccgaggctt tttcagacta 840
 ttattcaaat gacacagaat gaagacctcc gagaaaacat gttacggcaa gttctggagc 900
 atttgtctca gcaaagtga agccagtacc taaagattct aacaagcctt gctgaagttg 960
 ctacaacaaa tggtcataaa ctgcttagcc tctctagcaa ttatgatgct caaatgaaga 1020
 gccttttaag gattgtgaga atgtttgtc acgtctttcg aattgggtcca tccctcccca 1080
 gtaatggaat tgatatgggc tacaatggga ataaaactcc aagaagccag gtgttcaagc 1140

```

ctctggaatt gctttggcac tcgttagatg aatgggctagt tttaatagcc acagaattga 1200
tgaaaaaacaa aagagactca acagagatca cttctatttt actgaaacaa aaaggccaag 1260
atcaagatgc tgcttccatt cctccatttg aacctccagg acctgggagc tatgaaaatc 1320
tgtccactgg cacaaggga tctaaaccag atgctcttgc agggagacag gaagccagt 1380
cagattgtca ggtgttatt tctatgacag ctaaccggct aagtgtgtgc attcaagctt 1440
tttacctgtg ctgttcttgt cagatgcctc cgggaatgac ttcacctgtt ttcattgaat 1500
ttgtctgcaa acatgatgaa gttttaaaat gctttgttaa tagaaatccc aaaattatat 1560
ttgaccactt tcactttctc cttgaatgtc ctgagttgat gtcaagattc atgcatatca 1620
taaaagcaca gccttttaaa gatcgctgtg aatgggttcta tgaacatttg cattcaggac 1680
agccagattc agatatgggtg cacaggccag tgaatgaaaa tgatatcctg ctgggttcaca 1740
gagattctat ttttagagt agctgtgaag ttgtgtcaaa agcaaattgt gcaaagctaa 1800
agcaagggat tgctgtacgg ttccatggag aagaaggcat gggccaaggt gttgtgctg 1860
agtgggttga tattctgtcc aatgagatag tcaatcctga ttatgcattg tttaccagt 1920
cagctgatgg aacaactttt cagcctaata gcaactctta tgtaaactct gatcacttga 1980
actattttct gtttgcgtgg cagatcttgg gattagcgtt gaaccacagg cagctgggtca 2040
atatttactt cacacgatcc ttctacaagc acattcttgg tattcctgta aattaccaag 2100
atgtggcacc cattgatcca gaatatgcca aaaatttgca atggatttta gataatgata 2160
taagtgtatc ggtcttagaa ctaacttttt ctgttgagac tgatgtgttt ggagcaatgg 2220
aagaggtgcc tttgaaacct' ggggggtggga gtattcttgt gacacaaaat aataaagcgg 2280
agtacgtcca gcttgttact gaacttcgaa tgacaagagc cattcagcct cagatcaatg 2340
cttttttaca gggctttcat atgttcattc caccctccct catacagctt tttgatgaat 2400
atgaattgga gctactgctt tctggcatgc cagaaattga tgtgagtgat tggataaaaa 2460
atacagaata cacaagtggc tatgaaagag aagatccagt tattcagtgg ttctgggaag 2520
ttgtagaaga cattactcaa gaggagagag ttcttctctt acagtttgtt acgggcagtt 2580
ccagggtccc acatggtggg ttgtctaata tcatgggtgg aagtggattg caaaacttta 2640
caatcgctgc tgtgccatat actccaaatc ttttaccac ttcaagcaca tgcataca 2700
tgctcaagtt acctgaatac ccaagtaaag aaatactcaa ggacagactt cttgtggcac 2760
tacattgtgg cagctatggt tacacaatgg cataatgaag tctggaaaac tctctgact 2820
actgatgcac aattcagaat ggcagaagta atttgggaaa atgtcaacaa aaaagcagcc 2880
taaatgcaac ccataggcag ggtgatgtct tccaatttat aaaggatcat caggttttct 2940
gtttctctct tttccctttt atgttttctc ttgttgtgat acaattagaa aatataaaat 3000
cacagtagat tttattttttt' aaaatgctaa ctgaaagtaa tagagactgt cctttttcat 3060
aattaatttt atccaagatt gtattaaggg aaaatctgat tctacattcc acctctgcta 3120
tgtaactgtc ttgttaaaag ggtgttttct cctaatttct gatataattat atgaggtcat 3180
ccagctgggtg tgttcttttg catgtaaact gccatttata ttttagaaaa ctattgtata 3240
gaatggattt agattgtcta taaagccaca aatacgtatt ttgccacagt gtattctata 3300
ttgcaatgat ttttttagca ttttaatat ttaatatata ttgtaaaatt tagactgatg 3360
atactaacag ttgatgaaat gacatataat ttatatatga aagcttacgc tatattgtat 3420
gaattatttg catctttcag tggccagttt tccatatgta tatattatgg tctcaatgtt 3480
tttcttacgc ctcattttaa tttataatga aggtaaaatt aaaatgtatt ttaccacgtt 3540
tcttttcatt acttttatct gtgagctctg acacatctga aaaagtaatc tgatgtgcaa 3600
attataattt aaatatgtta atttttttgc ttcttaaatt tgcttttcat cattaaaatg 3660
tcaagttcaa gtgatatgtg cctaatatca cttggatgtt ggtgggtttt tgaatttttg 3720
ggtggttaat cagtttttat ttgaaaagac gtacttgaat agttacagca tatgtttgaa 3780
caggaagtag gaacatgcat acacgaagaa atgctaacgg aaggatttgt tatgtttagg 3840
atcttccctt ggaaactaaa aatagaatat taatgacatt actgtttgta gaatgacata 3900
tgcagatttt ctcataagca gtcatttgtt ttgccagtaa tgtttgagag acatgtaagt 3960
tgaaagtttt gctaaattat aaagctcctt taattcgttg gttttgattc tcttattctc 4020
ttgtcttttc taaatgttaa caaaatatat cttaacagat tacatgaaat ttaggaatta 4080
tttaaaagtt accattagct ctaaaattaa gattcggatg ctttatttat agtaactgaa 4140
gctaataatg ttttatgttt tgattttttg aaatttaatt gtagaagtca ctgccttctg 4200
agttttcaaa tagataacca cctttaatat tacactgctt ataatactaa tgtttacaga 4260
tatgtttctg tttataacca tataatacat tggctttgtc atattagttt tttttgcaag 4320
tagttatgta aaagagatag ataataaaat attaaataac tgagaaaaaa aaaaaaaaaa 4380
aaaa

```

<210> 30

<211> 4007

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475875CB1

<400> 30

```

atggcatgtt ccatggcgctg tggcggtaga gcttgcaagt atgagaaccc agcccgctgg 60
agttagcagg agcaagccat taaggggggtt tactcatcct gggtcactga taatatactg 120
gccatggccc gcccatcctc tgagctcctg gagaagtacc acatcattga tcagttcctc 180
agccatggca taaaaacaat aatcaacctc cagcgccctg gtgagcatgc tagctgtggg 240
aaccctctgg aacaagaaag tggcttcaca taccttctctg aggttttcat ggaggttggc 300
atttactttc acaatttcgg atggaaggat tatggtgtag cgtctcttac tactatccta 360
gatatggtga aggtgatgac atttgcctta caggaaggaa aagtagctat ccattgtcat 420
gcagggttgg gtcgaacagg tgttttaata gcctgttact tagtttttgc aacgagaatg 480
actgctgacc aagcaattat atttgtgcgg gcaaagcgac ccaattccat acaaaccaga 540
ggacagctcc tctgtgtaag ggaatttact cagtttctaa ctctctccg caatatattc 600
tcttgctgtg atcccaaagc acatgctgtc accttacctc aatatctaat tcgccagcgt 660
catctgcttc atggttatga ggcacgactt ctgaaacacg tgccaaaaat tatccacctc 720
gtttgcaaat tgctgctgga cttagcggag aacaggccag tgatgatgaa ggatgtgtcc 780
gaaggacctg gtctctctgc tgaaatagaa aagacaatgt ctgagatggg caccatgcag 840
ctggataaag agttactgag gcatgacagt gatgtgtcca acccgctaa ccccatgca 900
gtggcagcag attttgacaa tcgaggcatg attttctcca atgagcaaca gtttgacct 960
ctttggaaaa ggcggaatgt tgagtgccct caaccctga ctcatctgaa aaggcggtc 1020
agctacagtg actcagattt aaagagggcc gagaacctcc tggagcaagg ggagactcca 1080
cagacagtgc ctgccagat cttggttggc cacaagccca ggcagcagaa gctcataagc 1140
cattgtttaca tcccacagtc tccagaacca gacttacaca aggaagcctt ggttcgcagc 1200
acactttctt tctggagtca gtcaaagttt ggaggcctgg aaggactcaa agataatggg 1260
tcaccaattt tccatggaaa gatcattcca aaggaagcac agcagagtgg agctttctct 1320
gcagatggtt caggttcaca cagccctggg gagccagttt caccagctt tgcaaatgtc 1380
cataaggatc caaacctgc tcaccagcaa gtgtctcact gtcagtgtaa aactcatggt 1440
gttgggagcc ctggctctgt caggcagaac agcaggacac cccgaagccc tctggactgt 1500
ggctccagtc ccaaagcaca gttcttggtt gaacatgaaa cccaggacag taaagatctg 1560
tctgaagcag cttcacactc tgcattacag tctgaattga gtgctgaggc aagaagaata 1620
ctggcgggcca aagccctagc aaatttaaat gaatctgtag aaaaggagga actaaaaagg 1680
aaggtagaaa tgtggcagaa agagcttaat tcccagatg gagcttggga aagaatatgt 1740
ggcgagaggg accctttcat cctatgcagc ttgatgtggt cttgggtgga gcaaatgaag 1800
gagcctgtaa tcaccaaaga ggatgtggac atgttggttg acagcgagc agatgccgca 1860
gaagcacttt ttttattaga gaaggacag caccagacta ttctctgcgt gttgcactgc 1920
atagtgaacc tgcagacaat tcccgtagg gtggaggaag ctttccttgc ccatgccatt 1980
aaggcattca ctaaggttaa ttttgattct gaaaatggac caacagttta caacacctg 2040
aagaaaatat ttaagcacac gctggaagaa aaaagaaaaa tgacaaaaga tggccctaag 2100
cctggcctct agcttttact cgtggtgaat atttcagacc taaagatcca gatagtatct 2160
ctgttcatat gtgaataagt tgaagattgt ggggctactt tttctcatag cactttattt 2220
tgaatgttgt tagtttgtgc tgagaatggt cgtccgtatt tgaaccaatt atttatttta 2280
aaatatattt aagctacatt tttgttttga aaaattgcca taaatttggg gccactttct 2340
tttattttatt tgactgagtt aatattattg tattaacatt ttaagtatat ggtgtttaca 2400
ttcttatttc tttggcattt tggaaataat cataacttgt ctttccaaaa taaccatttt 2460
cttgatggaa ctcttcctag agtttttacc aaatagctaa ctttagtagt aaaacctcat 2520
tgtgtatcca tccccccaca gatgaactaa gaaagtcacc aagtgtctta agctgtttta 2580
tatttgttac gaagaaggct attgctacaa tattttttaa ggtttctttt ttaactttga 2640
aattttttgt ttttctttt ctttttataa atgtaacaga gggtttcaaa gcatattatt 2700
tttcagagag atttagtttt actttaatgg agtgactgtg aagtggttgg gattttttgc 2760
ttgtagaaaag tagacttgct ctttgtcaga tttccaaaca accttgccag ccttggctgt 2820
caaaaggagg caggagcagt tctcaacaca ccaagcetta tccccactcc cttgggttgc 2880
tgctgagcca aatagcatct ttacagagga agtgggatca gaggcaggaa gtgtggaaag 2940
ttgctaagaa gcagggttg cctctgtcct cccggggact ccacagggat attcgtgcag 3000
ggcaggggct ctgtgccagc cctgctctct cagatgccac agccactctg cagaggtgac 3060
tcttgagct ggaggaagtc aaaactgggc cactgtttgt actgatggtg tattagcatg 3120
agcagcgtgg ccctggcccc acactcccaa atctgccact ccatagacct acttgcctca 3180
aggctttata tttggctgct ttcttacaat gagaattaag atttttaaac tgaagttgac 3240
catacaggtt gcattagccc taactggctt catgtaagaa ggggtactgc cttaaactagt 3300
tcctgttaag ctgaaccata aattatcagt tgaagccata cttttattta aattaatata 3360
cgtagatacc agaggccaag ccacagagag gataatagtt cttcccaata aaggtgatat 3420
taatcagact aatttcgaac taaagaagtt actgcttaaa gacggaattt caggggaagc 3480
aagactcatt tagaacaat gaaatttctc cagtcttaca tttctgaatt gacttctagc 3540
acatcaaaaa tatttcagtc attatcagtc tcattaactg aaatgccaaa tgctaaatgc 3600

```


agtgttcttt	cacactgttt	taattttctt	gggaaattga	gtccagtga	tgttaaatgga	3660
gtgggttgcc	catccctgaa	atgtcttatt	ttcaagtgcc	tggcctggga	aagaagggga	3720
agaaacaatt	gcattatatt	caaagataca	ctataaaaaat	agagttttta	ccaaaaaaag	3780
atgtttgttc	tcactctcagt	aggcctcatt	tgggcaagt	acccacaggt	cttttggcga	3840
gtttgtctatt	tgccctgttga	aatacttggt	tcaacttaga	gaacagttat	gatgtgacca	3900
tagcatggca	caactaaaaa	tctaagcctg	aaacctgaaa	aaagagatat	gacaagggaa	3960
attaatcagg	ctatacataa	gtattgtatt	tatttgaata	aaaataa		4007

<210> 31

<211> 4524

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 71231882CB1

<400> 31

ccgccctcgc	caacatggcg	gcgcccagtt	ggggcggggt	cgttcgcttc	gcgttttggc	60
cagggcgggg	gtctgggctt	taggcaggta	gtatttagtt	tcacaatgtt	tggggacctg	120
tttgaagagg	agtattccac	tgtgtctaat	aatcagtatg	gaaaaggga	gaaattaaag	180
actaaagctt	tggagccacc	tgctcctaga	gaattcacca	atttaagcgg	aatcagaaat	240
caggggtggaa	cctgttacct	caattccctt	cttcagactc	ttcatttcac	acctgaattc	300
agagaagctc	tattttctct	tggcccagaa	gagcttggtt	tgtttgaaga	taaggataaa	360
cccgatgcaa	aggttcgaat	catcccttta	cagttacagc	gcttgtttgc	tcagctttctg	420
ctcttagacc	aggaagctgc	atccacagca	gacctcactg	acagctttgg	gtggaccagt	480
aatgaggaaa	tgaggcaaca	tgatgtgcag	gaactgaatc	gaatcctctt	cagcgctttg	540
gaaacttctt	tagttgggac	ctccggtcat	gacctcatct	atcgtctgta	ccatggaacc	600
attgttaacc	agattgtttg	taaagaatgt	aagaacgtta	gcgagaggca	ggaagacttc	660
ttagatctaa	cagtagcagt	caaaaatgta	tccggtttgg	aagatgctct	ctggaacatg	720
tatgtagaag	aggaagtttt	tgattgtgac	aacttgtacc	actgtggaac	ttgtgacagg	780
ctgggttaaag	cagcaaaagc	ggccaaatta	ctttagactg	ctccttttct	tactgtttca	840
ttactaagat	ttaattttga	ttttgtgaaa	tgcgaaacgt	acaaggaaac	tagctgttat	900
acattccctc	tccggattaa	tctcaagccc	ttttgtgaac	agagtgaatt	ggatgactta	960
gaatatatat	atgacctctt	ctcagttatt	atacaciaag	gtggctgcta	cggaggccat	1020
taccatgtat	atattaaaga	tgttgatcat	ttgggaaact	ggcagtttca	agaggaaaaa	1080
agtaaaccag	atgtgaatct	gaaagatctc	cagagtgaag	aagagattga	tcattccactg	1140
atgattctaa	aagcaatctt	attagaggag	gagaataatc	taattcctgt	tgatcagctg	1200
ggccagaaaac	ttttgaaaaa	gataggaata	tcttggaaac	agaagtacag	aaaacagcat	1260
ggaccattgc	ggaagttctt	acagctccat	tctcagatat	ttctactcag	ttcagatgaa	1320
agtacagttc	gtctcttgaa	gaatagttct	ctccaggctg	agtctgattt	ccaaagggaat	1380
gaccagcaaa	ttttcaagat	gcttctctca	gaatccccag	gtttaaacia	tagcatctcc	1440
tgtccccact	ggtttgatat	aaatgattct	aaagtccagc	caatcaggga	aaaggatatt	1500
gaacagcaat	ttcagggtaa	agaaagtgcc	tacatgttgt	tttatcgga	atcccagttg	1560
cagagacccc	ctgaagctcg	agctaataca	agatatgggg	ttccatgtca	tttactgaat	1620
gaaatggatg	cagctaaccat	tgaactgcaa	accaaaaggg	cagaatgtga	ttctgcaaac	1680
aatacttttg	aattgcatct	tcacctgggc	cctcagtatc	atttcttcaa	tggggctctg	1740
cacccagtag	tctctcaaac	agaaagcgtg	tgggatttga	cctttgataa	aagaaaaact	1800
ttaggagatc	tccggcagtc	aatatttcag	ctgttagaat	tttgggaagg	agacatgggt	1860
cttagtggtg	caaagcttgt	accagcagga	cttcacattt	accagtcact	tggcggggat	1920
gaactgacac	tgtgtgaaac	tgaaattgct	gatggggaag	acatctttgt	gtggaatggg	1980
gtggagggtg	gtggagtcca	cattcaaact	ggtattgact	gcgaacctct	acttttaaat	2040
gttcttcac	tagacacaag	cagtgatgga	gaaaagtgtt	gtcagggtgat	agaattctcca	2100
catgtctttc	cagctaattg	agaagtgggc	actgtcctca	cagccttagc	aatcccagca	2160
ggtgtcatct	tcatacaacg	tgctggatgt	ccagggtggg	agggttggac	ggccatcccc	2220
aaggaagaca	tgagggaagc	gttcaggggg	caagggtcca	gaaatggaag	ctcaatttta	2280
attcaggatt	ctcatgatga	taacagcttg	ttgaccaagg	aagagaaatg	ggtcactagt	2340
atgaatgaga	ttgactggct	ccacgttaaa	aatttatgcc	agttagaatc	tgaagagaag	2400
caagttaaaa	tatcagcaac	tgtaaacaca	atgggtgttg	atattcgaat	taaagcata	2460
aaggaattaa	aattaatgaa	ggaactagct	gacaactagc	gttttgagacc	tattgataga	2520
aatgggaagc	ttctttgtcc	agtgcgggac	agctatactt	tgaagggaagc	agaattgaag	2580
atgggaagtt	cattgggact	gtgtcttgga	aaagcaccaa	gttcgtctca	gttggttctg	2640
ttttttgcaa	tggggagtga	cgttcaacct	gggacagaaa	tggaaatcgt	agtagaagaa	2700

```

acaatatctg tgagagattg tttaaagtta atgctgaaga aatctggcct acaaggagat 2760
gcctggcatt tacgaaaaat ggattgggtgc tatgaagctg gagagccttt atgtgaagaa 2820
aattcagcca gaagccaaact cattaccctt ggaactggct tctcgttcca gccttgccag 2880
gatgcaacac tgaaagaact tctgatatgt tctggagata ctttgctttt aattgaagga 2940
caacttcctc ctctgggttt cctgaaggtg cccatctggg ggtaccagct tcagggtccc 3000
tcaggacact gggagagtca tcaggaccag accaactgta cttcgtcttg gggcagagtt 3060
tgagagacca cttccagcca aggtgcttct ggggaacgagc ctgcgcaagt ttctctcctc 3120
tacttgggag acatagagat ctcaagaat gccacgctgg cggagctgaa gtctcaggcc 3180
atgaccttgc ctcttttctt ggagttcggg gtcccgtccc cagcccacct cagagcctgg 3240
acgggtggaga ggaagcgccc aggcaggctt ttacgaactg accggcagcc actcaggga 3300
tataaactag gacggagaat tgagatctgc ttagagcccc ttcagaaagg cgaaaacttg 3360
ggcccccagg acgtgctgct gaggacacag gtgcgcatcc ctggtgagag gacctatgcc 3420
cctgccctgg acctgggtg gaaacggggc cagggtggga ctgccggctc cctgaggcag 3480
agagttgccg atttctatcg tcttcccggt gagaagattg aaattgccaa atactttccc 3540
gaaaagtctg agtggcttcc gatattagc tggaaccaac aaataacca gagggaaaa 3600
aaaaaaaaac aagattatct gcaaggggca ccgtattact tgaaagacgg agatactatt 3660
ggtgttaaga atctcctgat tgacgacgat gatgatttca gtacaatcag agatgacact 3720
ggaaaagaaa agcagaaaac acgggccctg gggagaagga aaagccaaga agccctccat 3780
gagcagagca cctacatcct ctccagtcca gagacgctg cccggccccg agccccgga 3840
acttctctct ccatccacgt ggggagcttc agataaccgc gccgctgcac ggctctactc 3900
ccgatgaact ctccggctga tgcacaaaac gtgggtttcc tgggcatggg gactggctgc 3960
ctggcgcttc caatcccaa tcctctgctt cctttgagca cagggacggc tcctctgagg 4020
cctggccagt gcatgtagtc acttagctct gcaacacgtg gcagccacgg gggctgggtg 4080
agctctggat gtcgcccacc cagctgccag taggtgctgg gctctctcac acagcaccg 4140
gccccagctg cctttttttt tcttttaacc agaaaatgca caacgtgtgc gtgaaccgca 4200
ggtatggagg cagcggcatg ccgttgctcc gctgtgggag gtgtgtgggg tcaggccagc 4260
cactttcctc cgtgttcaga tgactctcgt tcgccctgac cggcttctca cagtgtctca 4320
ggccactgcg ccaccgcgct ggtgctgagc agaagcgggc agaagtgggg tctgctttca 4380
ggacttcatt tcccccactc gttccggccc cgcattgctc acgtctgccc tttggtctga 4440
gttaaaactg cgatgctgaa aagtgcgagc tctttccacg aggaggagcc acacagggtg 4500
gcctccgagg gtgagtcgct ctgc 4524

```

<210> 32

<211> 3250

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2875922CB1

<400> 32

```

cgccggctgt ggccgcggcg aacaggccgg acgctcgtc gctggcgggg cttcccttgg 60
agctgacgaa ggcgggggtcc ggggtccagg ctgctgccgc gacggggccg gcggcggggc 120
agctgccagg aacgagggta gcagctgcat ccagatctca ttatgcatca gaaaaatgaa 180
aaaacagagg aaaattctat ggaggaaaagg aatccactta gccttttctg agaaatggaa 240
tactgggttt ggaggtttta agaagtttta ttttcaccaa cacttgtgca ttctgaaagc 300
taagctggga aggccagtta cttggaatag acagttaga catttccagg gtagaaagaa 360
agctcttcaa atccagaaaa cgtggatcaa ggatgaaccc ctttgtgcta agaccaagtt 420
caatgtggct actcaaaatg ttagtacttt gtccctctaa gtgaaaagaa aggacgctaa 480
acacttcatt tctcctcaa agactctcct gagactccaa gcagagaagc tggtgtcatc 540
agcaaaagaa tctgacctg aatactgcag agagaaaaat ctcttgaagg cagtacttga 600
ctttccatca aatagtgtt taggtcaggc caatggtcac agacctagga cagaccaca 660
accttctgac tttcccatga agttcaatgg ggagagccaa agtccagggt agagtggcac 720
gattgtggtc acctgaaca accataagag aaagggtctt tgttacggct gctgccaaag 780
gccggagcac cacaggaatg ggggaccctt gattccaaa aagttccaac ttaaccaaca 840
tagaaggata aaattatctc ctcttatgat gtatgagaaa ttatccatga ttagatttctg 900
gtacaggatt ctcatatccc agcacttcac aacaaaagc aaggtttgca agctaagaaa 960
agcccagcga agctgggtac agaaagtac tggggacat caagagaccc gtagggagaa 1020
cgggtgagggt ggcagttgca gccatttcc tttcccagaa cctaaagacc cttcttctgc 1080
gcatcagccg tactttccag atatggacag cagtgtgtgt gtgaagggga cgaactctca 1140
tgtgcctgat tgccacacta aaggaagctc tttcttgggc aaggagctta gtttagacga 1200
agcattccct gaccaacaga atggcagttc cacaacgcc tgggaccagt catcctgttc 1260

```

```

ttctcctaag tgggagtgtg cagagctgat tcatgacatc cccttaccag aacatcggtc 1320
taataccatg ttcatttcag aaactgaaag agaaattatg actctgggtc aggaaaatca 1380
gacaagttct gtcagtgatg acagagtaaa actgtcagtg tctggagcag atacatctgt 1440
gagtgcgtg gatgggctg tgtcccaaaa ggctgttcaa aatgagaact cataccagat 1500
ggaggaggat ggatctctca agcagagcat tcttagttct gagttgctgg accaccctta 1560
ctgtaaaagt ccactggagg ctcccttggt gtgcagtggg ctcaaactag aaaatcaagt 1620
aggaggtgga aagaacagtc agaaagcctc tccagtggat gatgaacagc tgtcagtctg 1680
tctttctgga ttcctagatg aggttatgaa gaagtatggc agtttggttc cactcagtga 1740
aaaagaagtc cttggaagat taaaagatgt ctttaatgaa gacttttcta atagaaaacc 1800
atztatcaat agggaaataa caaactatcg ggccagacat caaaaatgta acttccgtat 1860
cttctataat aaacacatgc tggatatgga cgacctggcg actctggatg gtcagaactg 1920
gctgaatgac cagggtcatta atatgtatgg tgagctgata atggatgcag tcccagacaa 1980
agttcacttc ttcaacagct tttttcatag acagctggta accaaaggat ataatggagt 2040
aaaaagatgg actaaaaagg tggatttggt taaaaagagt cttctgttga ttcctattca 2100
cctggaagtc cactggtctc tcattactgt gacactctct aatcgaatta tttcatttta 2160
tgattcccaa ggcattcatt ttaagttttg tgtagagaat ataagaaagt atttgctgac 2220
tgaagccaga gaaaaaata gacctgaatt tcttcagggt tggcagactg ctgttacgaa 2280
gtgtattcca caacagaaaa acgacagtga cgttggagtc tttgtgctcc agtactgcaa 2340
gtgcctcgcc ttagagcagc ctttccagtt ttcacaagaa gacatgcccc gagtgcggaa 2400
gaggatttac aaggagctat gtgagtgcgc gctcatggac tgaaactcag cagggactct 2460
gggaagtctg accaagttag agcagatggg ttgttacttg aatctccaaa cacttagttg 2520
aatttttaca gatatttcag atcagtggtg ttgggccact attgttacct caaatattat 2580
ttttgccctt attcatttct ccagctacca tgtactattg tttaatgttc agtttggttt 2640
catttttaat tttatggttc tgtgcgtccc ccatatttaa tatttattat tcaaacgcat 2700
gcatatagac agagcatgca gtgaagagta ttaaaaaaaa aagcttagta gatttggtgc 2760
agcttttgaa acttagttag acgtgaactg aatacagggt tcaaatttac tcccagaacc 2820
taaaaatgca agatgttttt gatacaacat aactctgaga atagtaagtg ttccttgggg 2880
cattaagggt agctgggggt ggttttgaca aatccagtc tgttttactt taccagcggc 2940
aactttcacc aacttccccct ctccaagtga gtcttagaga gtgcagtcca ttccttttga 3000
agggtgagat ggaagtgggc gtaaaactgac tgggtgtctc tgtttctgga ggcacacttg 3060
taagcacagt ggctgctttg ggaggagtaa ggtgtgagaa aaagcaacct tggaggccag 3120
taacaatgac cgtggtttta cgtggttataa ggaattataa tacgtggcat acatctcata 3180
aaggcttttg ctgggatatt gaattccctg aatttttctg ttttcgacct gttaaaaaaa 3240
tcttaacatc                                     3250

```

<210> 33

<211> 3834

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 8158136CB1

<400> 33

```

gcggccgggtc gtgcgggtcg ggcgcgggcg ggcgcgggcg cagtggcgcg cacaggtgat 60
tgactggcca gctgcctgaa ggagcgccag gtctctcttg ctggcagggt gcgaagccca 120
ttggggcgggc ggtgcagacc gcgcgggcggt ctgcggcggt ctggctcggg aggcgttcct 180
ggggccaagg ccatggcccc gcggtctgag ctggagaagg cggcctggcg ctggcgagg 240
acggtgcggc ccgaggagggt gtgcaggag cacatcgaga ccgcttaccg catctggctg 300
gagccctgca ttcgcggcgt gtgcagacga aactgcaaag gaaatccgaa ttgcttggtt 360
ggtattgggt agcatatttg gttaggagaa atagatgaaa atagttttca taacatcgat 420
gatcccaact gtgagaggag aaaaaagaac tcatttgggt gcctgactaa ccttgaggcc 480
acttgttatg tcaacacatt tcttcaagtg tggtttctca acttgagct tcggcaggca 540
ctctacttat gtccaagcac ttgtagttag tacatgctgg gagacggcat ccaagaagaa 600
aaagattatg agcctcaaac aatttgtgag catctccagt acttgtttgc cttgttgcaa 660
aacagtaata ggcgatacat tgatccatca ggatttggtt aagccttggg cctggacact 720
ggacaacagc aggatgctca agaattttca aagctcttta tgtctctatt ggaagatact 780
ttgtctaaac aaaagaatcc agatgtgcgc aatattgttc aacagcagtt ctgtgtgagaa 840
tatgcctatg taactgtttg caaccagtggt ggcagagagt ctaagctttt gtcaaaattt 900
tatgagctgg agttaaatat ccaaggccac aaacagttta cagattgtat ctcggaattt 960
ttgaagggaag aaaaattaga aggagaqaat cgctattttt gcgagaactg tcaaagcaaa 1020
cagaatgcaa caagaaagat tcgacttctt agccttctct gcactctgaa cttgcagcta 1080

```

```

atgcggttttg tctttgacag gcaaactgga cataagaaaa agctgaatac ctacattggc 1140
ttctcagaaa ttttggatat ggagccttat gtggaacata aaggtgggtc ctacgtgtat 1200
gaactcagcg cagtcctcat acacagagga gtgagtgcct attctggcca ctacatcgcc 1260
cacgtgaaag atccacagtc tggatgaatgg tataagttta atgatgaaga catagaaaag 1320
atggagggga agaaattaca actagggatt gaggaagatc tagcagaacc ttctaagtc 1380
cagacacgta aacccaagtg tggcaaagga actcattgct ctgaaatgc atatatgttg 1440
gtttatagac tgcaaactca agaaaagccc aacactactg ttcaagttcc agcctttctt 1500
caagagctgg tagatcggga taattccaaa tttgaggagt ggtgtattga aatggctgag 1560
atgcgtaagc aaagtgtgga taaaggaaaa gcaaaacacg aagaggttaa ggagctgtac 1620
caaaggttac ctgctggagc tgagccctat gagtttgcct ctctggaatg gctgcaaaag 1680
tggttggatg aatcaacacc taccaaacct attgataatc acgcttgctt gtgttcccat 1740
gacaagcttc acccgataa aatatcaatt atgaagagga tatctgaata tgcagctgac 1800
attttctata gtagatagg aggaggtcca agactaactg tgaaagccct gtgtaaggaa 1860
tgtgtagtag aacgttgtcg catattgcgt ctgaagaacc aactaaatga agattataaa 1920
actgttaata atctgtgaa agcagcagta aagggcgatg gattttgggt ggggaagtcc 1980
tccttgcgga gttggcgcca gctagctctt gaacagctgg atgagcaaga tgggtgatgca 2040
gaacaaagca acggaaagat gaacggtagc accttaaata aagatgaatc aaaggaagaa 2100
agaaaagaag aggaggaatt aaattttaat gaagatatc tgtgtccaca tgggtgagtt 2160
tgcatactcg aaaatgaaag aaggcttggt tctaaagagg cttggagcaa actgcagcag 2220
tactttccaa aggtcctga gtttccaagt tacaagagt gctgttcaca gtgcaagatt 2280
ttagaaagag aaggggaaga aaatgaagcc ttacataaga tgattgcaaa cgagcaaaag 2340
acttctctcc caaatttggt ccaggataaa aacagaccgt gtctcagtaa ctggccagag 2400
gatacggatg tcctctacat cgtgtctcag ttctttgtag aagagtggcg gaaatttggt 2460
agaaagccta caagatgcag ccctgtgtca tcagttggga acagtgcctt tttgtgtccc 2520
cacgggggccc tcattgtttac atttgccttc atgaccaaag aagattctaa acttatagct 2580
ctcatatggc ccagttagtg gcaaatgata caaaagctct ttgttgtgga tcatgtaatt 2640
aaaatcacga gaattgaagt gggagatgta aacccttcag aaacacagta tatttctgag 2700
cccaaactct gtccagaatg cagagaaggg ttattgtgtc agcagcagag ggacctgcgt 2760
gaatacactc aagccaccat ctatgtccat aaagtgtgg ataataaaaa ggtgatgaag 2820
gattcggtc cggaactgaa tgtgagtagt tctgaaacag aggaggacaa ggaagaagct 2880
aaaccagatg gagaaaaaga tccagatttt aatcaaagca atggtggaac aaagcgcaa 2940
aagatatccc atcaaaatta tatagcctat caaaagcaag ttattcgccg aagtatgca 3000
catagaaaag tctgtggtga gaaagcactt ctggttctg ctaatcagac gttaaaagaa 3060
ttgaaaattc agatcatgca tgcattttca gttgtcctt ttgaccagaa tttgtcaatt 3120
gatggaaaga ttttaagtga tgactgtgcc accctaggca cccttggcgt cattcctgaa 3180
tctgtcattt tattgaaggc tgatgaacca attgcagatt atgctgcaat ggatgatgtc 3240
atgcaagttt gtatgccaga agaaggggtt aaagggtactg gtcttcttgg acattaatct 3300
ttgaatactt gctgactgct aagaaatgac cagaggggaa gaggagtgtg acatgttagg 3360
gcattaaagc aaaggtggat ttaagaatta aaccattaca tgccccttcc aaaaggcaga 3420
aatccattca aacgtgactg tcccaaatgc cttatgtcaa ataaagcaga ttgcactgat 3480
ggacatcaga cttgaaggaa atgtttccaa ttttatattt aaggggggtg gtgggtggga 3540
gggggcaagt aaagacggaa caagttagt agcagtaata gtaaactcatg tttacatatg 3600
agatttatag tctgtggagg ggaataaagt tctgttatat ttccttgctc gagtttcata 3660
ccagatgcgt tgggtccataa aggattgtat caagtagatg ggacaacatt ctgctctgaa 3720
cgaaaagtaa ttttagagac ataacctgct taccaatgcc tgtctttgat tcatattcta 3780
ctttcaataa agcatgaaag tgaaaaaaa aaaaaaaaaa aaaaaaaa aaaa 3834

```

<210> 34

<211> 4493

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5969491CB1

<400> 34

```

gcgcccagtt ggggcggggt cgttcgcttc gcgtttggcc aggcgggggt ctgggcttta 60
ggcaggtagt atttagtttc acaatgtttg gggacctggt tgaagaggag tattccactg 120
tgtctaataa tcagtatgga aaagggaaga aattaaagac taaagctttg gagccacctg 180
ctcctagaga attcaccaat ttaagcggaa tcagaaatca ggggtggaacc tgttacctca 240
attcccttct tcagactctt catttcacac ctgaattcag agaagctcta ttttctcttg 300
gccagaaga gcttggtttg tttgaagata aggataaacc cgatgcaaag gttcgaatca 360

```

tccctttaca	gttacagcgc	ttgtttgctc	agctttctgt	cttagaccag	gaagctgcat	420
ccacagcaga	cctcactgac	agctttgggt	ggaccagtaa	tgaggaaatg	aggcaacatg	480
atgtgcagga	actgaatcga	atcctcttca	gcgctttgga	aacttcttta	gttgggacct	540
cgggtcatga	cctcatctat	cgtctgtacc	atggaacat	tgtaaccag	attgtttgta	600
aagaatgtaa	gaacgttagc	gagaggcagg	aagacttctt	agatctaaca	gtagcagtca	660
aaaatgtatc	cggtttgga	gatgctctct	ggaacatgta	tgtagaagag	gaagtttttg	720
attgtgacaa	cttgtaaccac	tgtggaactt	gtgacaggct	ggttaaagca	gcaaagtcgg	780
ccaaattacg	taagctgcct	ccttttctta	ctgtttcatt	actaagattt	aatttttgatt	840
ttgtgaaatg	cgaacgctac	aaggaaacta	gctgtttatac	attccctctc	cggattaatc	900
tcaagccctt	ttgtgaacag	agtgaattgg	atgacttaga	atatatatat	gacctcttct	960
cagttattat	acacaaaggt	ggctgctacg	gaggccatta	ccatgtatat	attaaagatg	1020
ttgatcattt	gggaaactgg	cagtttcaag	gagaaaaaag	taaaccagat	gtgaacttga	1080
aagatctcca	gagtgaagaa	gagattgatac	atccactgat	gattctaaaa	gcaatcttat	1140
tagaggagga	gaataatcta	attcctgttg	atcagctggg	ccagaaactt	ttgaaaaaga	1200
taggaatatac	ttggaacaag	aagtacagaa	aacagcatgg	accattgcgg	aagttcttac	1260
agctccattc	tcagatattt	ctactcagtt	cagatgaaag	tacagttcgt	ctcttgaaga	1320
atagttctct	ccaggctgag	tctgatttcc	aaaggaatga	ccagcaaatt	ttcaagatgc	1380
ttcctccaga	atccccaggt	ttaaacaata	gcactctctg	tccccactgg	tttgatataa	1440
atgattctaa	agtccagcca	atcagggaaa	aggatattga	acagcaattt	cagggttaaag	1500
aaagtgccta	catgttggtt	tatcggaaat	cccagttgca	gagacccctt	gaagctcgag	1560
ctaataccaag	atatgggggt	ccatgtcatt	tactgaatga	aatggatgca	gctaacattg	1620
aactgcaaac	caaaaaggga	gaatgtgatt	ctgcaaacaa	tacttttgaa	ttgcatcttc	1680
aacctggggcc	tcagtatcat	ttcttcaatg	gggctctgca	cccagtagtc	tctcaaacag	1740
aaagcgtgtg	ggatttgacc	tttgataaaa	gaaaaacttt	aggagatctc	cggcagtcga	1800
tatttcagct	tttagaattt	tgggaaggag	caattgttct	tagtggttga	aagcttgta	1860
cagcaggact	tcacatttac	cagtcaactg	gcggggatga	actgacactg	tgtgaaactg	1920
aaattgctga	tggggaagac	atctttgtgt	ggaatggggg	ggaggttggt	ggagtccaca	1980
ttcaaactgg	tattgactgc	gaacctctac	ttttaaatgt	tcttcattct	gacacaagca	2040
gtgatggaga	aaagtgttgt	caggtgatag	aatctccaca	tgtctttcca	gctaattgcag	2100
aagtgggcac	tgtcctcaca	gccttagcaa	tcccagcagg	tgtcatcttc	atcaacagtg	2160
ctggatgtcc	aggtggggag	ggttggacgg	ccatcccca	ggaagacatg	aggaagacgt	2220
tcaggagca	agggctcaga	aatggaaagt	caattttaat	tcaggattct	catgatgtat	2280
acagcttggt	gaccaaggaa	gagaaatggg	tcaactagat	gaatgagatt	gactggctcc	2340
acgttaaaaa	tttatgccag	ttagaatctg	aagagaagca	agttaaaata	tcagcaactg	2400
ttaacacaat	ggtgtttgat	attcgaatta	aagccataaa	ggaattaaaa	ttaatgaagg	2460
aactagctga	caacagctgt	ttgagacctt	ttgatagaaa	tgggaagctt	ctttgtccag	2520
tgccggacag	ctatactttg	aaggaagcag	aattgaagat	gggaagtcca	ttgggactgt	2580
gtcttgga	agcaccaagt	tcgtctcagt	tgttcctggt	ttttgcaatg	gggagtgcag	2640
ttcaacctgg	gacagaaatg	gaaatcgtag	tagaagaaac	aatatctctg	agagattggt	2700
taaagttaat	gctgaagaaa	tctggcctac	aaggagatgc	ctggcattta	cgaaaaatgg	2760
attgggtgcta	tgaagctgga	gagcctttat	gtgaagaaga	tgcaacactg	aaagaacttc	2820
tgatatgttc	tggagatact	ttgcttttaa	ttgaaggaca	acttcctcct	ctgggtttcc	2880
tgaaggtgcc	catctggtgg	taccagcttc	agggctccct	aggacactgg	gagagtcata	2940
aggaccagac	caactgtact	tcgtcttggt	gcagagtttg	gagagccact	tccagccaag	3000
gtgcttctgg	gaacgagcct	gcgcaagttt	ctctcctcta	cttgggagac	atagagatct	3060
cagaagatgc	cacgctggcg	gagctgaagt	ctcaggccat	gaccttgctt	cctttcctgg	3120
agttcgggtg	cccgctccca	gcccacctca	gagcctggac	ggtggagagg	aagcgcccag	3180
gcaggctttt	acgaactgac	cggcagccac	tcagggaata	taaaactagga	cggagaattg	3240
agatctgctt	agagcccctt	cagaaaggcg	aaaacttggg	cccccaggac	gtgctgctga	3300
ggacacaggt	gcgcacccct	ggtgagagga	cctatgcccc	tgccctggac	ctggtgtgga	3360
acggcgccca	gggtgggact	gccggctccc	tagggcagag	agttgccgat	ttctatcgtc	3420
ttcccgtgga	gaagattgaa	attgccaatt	actttcccga	aaagtctcag	tggcttccga	3480
tatctagctg	gaaccaacaa	ataaccaaga	ggaaaaagaa	aaaaaaacaa	gattatttgc	3540
aaggggcacc	gtattacttg	aaagacggag	atactatttg	tgtaagaat	ctcctgattg	3600
acgacgatga	tgatttcagt	acaatcagag	atgacactgg	aaaagaaaag	cagaaacaac	3660
gggcccctgg	gagaaggaaa	agccaagaag	ccctccatga	gcagagcagc	tacatcctct	3720
ccagtgcaga	gacgcctgcc	cggccccgag	ccccggaaac	ttctctctcc	atccacgtgg	3780
ggagcttcag	ataaccgcgc	cgtctgcagg	ctctactccc	gatgaactct	cgggtgatg	3840
ccacaacagt	gggttctctg	ggcatgggga	ctggctgcct	ggcgcttoca	atccccaatc	3900
ctctgcttcc	tttgagcaca	gggacggctc	ctctgagggc	tggccagtgc	atgtagtac	3960
ttagctctgc	aacacgtggc	agccacgggg	gctggtgcag	ctctggatgt	cgccaccca	4020
gctgccagta	ggtgctgggc	tctctcacac	agcaccgggc	cccagctgcc	tttttttttc	4080
ttttaaccag	aaaatgcaca	acgtgtgcgt	gaaccgcagg	tatggaggca	gcggcatgcc	4140

```

gttgcctccg tgtgggaggt gtgtgggggtc aggccagcca ctttctccg tgttcagatg 4200
actctcgttc gccctgaccg gcttctcaca gtgtctcagg ccactgcgcc accgcgctgg 4260
tgctgagcag aagcgggcag aagtgggggtc tgctttcagg acttcatttc cccactcgt 4320
tccggccccc gcatgctcca cgtctgccct ttggtctgag ttaaaactgc gatgctgaaa 4380
agtgcgagct ctttccacga ggaggagcca cacagggtgg cctccgaggg tgagtgcctc 4440
tgctaagcaa gggcagccgc tgcacgtcag cccgcaggcc aagggtccag ctt 4493

```

<210> 35

<211> 2921

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7497367CB1

<400> 35

```

ctggggctgg attgagctga ccacaggcca caccagactc ctctctgctc ctgaggaaga 60
cagggcagcc cggcgccacc cgctcgccgc tcacgaagat gctccctgga gctgggctgc 120
tctggacctc cctcctgctc ctggccaggg ctgcccagcc ctgtcccatg ggttgtgact 180
gcttcgtcca ggaggtgttc tgctcagatg aggagcttgc caccgtcccg ctggacatcc 240
cgccatatac gaaaaacatc atctttgttg agacctcgtt caccacattg gaaaccagag 300
cttttggcag taacccaac ttgaccaagg tggcttccct caacactcag ctctgccagt 360
ttagggccgga tgcctttggg gggctgcccc ggctggagga cctggagggtc acaggcagta 420
gcttcttgaa cctcagcacc aacatcttct ccaacctgac ctgctggggc aagctcacc 480
tcaacttcaa catgctggag gctctgcccg agggcttttt ccagcacctg gctgccctgg 540
agtccctcca cctgcagggg aaccagctcc aggcctgccc caggaggctc ttccagctc 600
tgacctatct gaagacactc aacctggccc agaacctcct ggcccagctc ccggaggagc 660
tgttccaccc actcaccagc ctgcagaccc tgaagctgag caacaacggc ctctctggtc 720
tccccagggg tgtgtttggc aaactgggca gcctgcagga gctcttctg gacagcaaca 780
acatctcgga gctgccccct caggtgttct cccagctctt ctgcctagag aggtctgtggc 840
tgcaacgcaa cgccatcacg cacctgcccg tctccatctt tgcctccctg ggtaactga 900
cctttctgag cttgcagtgg aacatgcttc gggctcctgc tgcggcctc tttgccaca 960
ccccatgctt ggttggcctg tctctgacct ataaccagct ggagactgtc gctgagggca 1020
cctttgcccc cctgtccaac ctgcgttccc tcatgtcttc atacaatgcc attaccacc 1080
tcccagctgg catcttcaga gacctggagg agttgggtcaa actctacctg ggcagcaaca 1140
accttacggc gctgcaccca gccctcttcc agaacctgtc caagctggag ctgctcagcc 1200
tctccaagaa ccagctgacc aacttccgg agggcatctt cgacaccaac tacaacctgt 1260
tcaacctggc cctgcacggg aacctctggc agtgcgactg ccacctggcc tacctcttca 1320
actggctgca gcagtacacc gatcggtccc tgaacatcca gacctactgc gctggccctg 1380
cctacctcaa aggccagggtg gtgcccgcct tgaatgagaa gcagctggtg tgtcccgta 1440
cccgggacca cttgggcttc caggtcacgt ggccggacga aagcaaggca gggggcagct 1500
gggatctggc tgtgcaggaa agggcagccc ggagccagtg cacctacagc aaccccagg 1560
gcaccgtggt gctgcgctgt gaccaggccc agtgtcgctg gctgaacgtc cagctctctc 1620
ctcggcaggg ctccctggga ctgcagtaca atgctagtca ggagtgggac ctgaggtcga 1680
gctgcgggtc tctgcggctc accgtgtcta tcgaggctcg ggcagcaggg ccctagtagc 1740
agcgcataca ggagctgggg aagggggcct ctggggcctg accaggcgac aggtaggggc 1800
ggaggggagc tgagtctccg aagccttggc ttttccatg caagggacag ggttacatcc 1860
ccaaggtgag ggggtggagt ctgggtctgt ccaactaaca gggctctctc ctctcttcc 1920
ttcatcgctt ctctggagt gtgcggccta acaaggccat ccttatgctt tgcaaaagc 1980
cctcaaaagc tgcaccacag cctggagaat aaaatatcct cagccctgat gcctccccat 2040
tatgtaacac ccaaccgctc tcacctacac cctgaggtct attcactgca tccagtgat 2100
acaaagtggg ggcactgcc ttctgacatc tggctcaaaa gccagtgctc tgtttccatt 2160
tatttccctg gaatttcatt taaaattggt atagagaaaa aaaggatgtg acagaagcag 2220
agatgaccag aaagcacagg ggcagggttc tgactggcgt gtgggagacc ctgtggccgg 2280
caccacctc cacacgagga ctaagctctg atttttttat cttgcccata ttctacctc 2340
aggggtctag ggagtcgcgc cttacaaatc ataaattctc atcagatggg ttttatttga 2400
cctgtatat catgacttat ttttaatctg actatggcat aacattacaa gacgaggcaa 2460
aaatatttaa cccccaaata tatttctttg cctaccttg aacttgccct gcagagtctc 2520
ttgtgaggag aatccacatc ctataaagaa gccctttcc cctttgtttt ccttcccttc 2580
tttccagctc aggagatcat caactaagag ccaggcacc cttttaagtc gataaagaa 2640
agtttacaac ctgctctctc tctctctgaa gtctgctgag agcttccct gcacaataaa 2700
acttggcctc cacaatcctt tatcttaacc tgaacattcc tttccattga tcccaggctc 2760

```

```

tcctcaacac tcagctctgc cagtttaggc cggatgcctt tgggggggctg cccaggctgg 2820
aggacctgga ggtcacaggc agtagcttct tgaacctcca ggtcctccag tttaggccgg 2880
atgccttttg ggggctgccc aggctggagg acctggaggt c 2921

```

<210> 36

<211> 2572

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7632424CB1

<400> 36

```

tagtgaccta tagaacacta ttagtcgcat gccgcgtcgt aagctcgggtg cagcgataag 60
ggcagtcgac agtcttttagt agggaaagga gacaagtgcct agctactgcc gcccaagtgg 120
aaggaaattat ctatagagta agtatgctaa tcttgactaa gactgcagga gtttttttta 180
aaccatcaaa aaggaaagtt tatgaatttt taagaagttt taattttcat cctggaacac 240
tattttcttca taaaatagta ttgggaattg aaactagttg tgatgataca gcagctgctg 300
tgggtggatga aactggaaat gtgttgggag aagcaatata ttcccaaact gaagttcatt 360
taaaaacagg tgggattggt cctccagcag ctcaacagct tcacagagaa aatattcaac 420
gaatagtaca agaagctctt tctgccagtg gagtctctcc aagtgcctc tcagcaattg 480
caactaccat aaaaccagga cttgctttta gccctgggag gggcttatca tttagcttac 540
agctggtagg acagttaaaa aagccattca ttcccattca tcatatggag gctcatgcac 600
ttactattag gttgaccaat aaagtagaat ttctttttt agttcttttg atttctggag 660
gtcactgtct gttggcatta gttcaaggag tttcagattt tctgcttctt ggaaagtctt 720
tggacatagc accaggtgac atgcttgaca aggtggcaag aagactttct ttaataaaac 780
atccagagtg ctccaccatg agtgggtgga aagccataga acatttggcc aaacaaggaa 840
atagatttca ttttgacatc aaacctcctc tgcatcatgc taaaaattgt gatttttctt 900
ttactggact tcaacacggt actgataaaa taataatgaa aaaggaaaaa gaggaaggta 960
ttgagaaggg gcaaatcctg tcttcagcag cagacattgc tgccacagta cagcacacaa 1020
tggcatgtca tcttgtgaaa agaacacatc gggctattct gttttgtaag cagagagact 1080
tgttacctca aaataatgca gtactgggtg catctgggtg tgtcgcaagt aacttctata 1140
tccgcagagc tctggaaatt ttaacaaacg caacacagtg cactttgttg tgtcctctc 1200
ccagactatg cactgataat ggcattatga ttgcatggaa tgggtattgaa agactacgtg 1260
ctggcttggg cattttacat gacatagaag gcatccgcta tgaacaaaaa tgtcctcttg 1320
gagtagacat atcaaaagaa gttggagaag ctccataaaa agtaccacaa ttaaaaatgg 1380
agatatgatt tctgctgttc aaaaaagtc ctaaaagggtc tcactctctg acctcagctg 1440
gagtacagta gccagatcac aactcactgc aacctgact tcctgaactc aagaaatcct 1500
cctgccttag cctcttgaat agccgggact acaggtgtgc atgtccatgc ccagccaact 1560
ttatttctat tttttgtaga gacaggctct tgccatgttg cccgggctgg tcctgaactg 1620
ctgaattcaa gtgatcctcc caccttggcc tccagaagtg ctgggattat ggggtgtgagc 1680
caccatgcct agccaaaatg tttcttaagg tatacathtt gggctttaga agacttatac 1740
atttgaataa tttattacta aatatctcaa agtattacaa taaatgttac catgtgagct 1800
actttgaatc aggcctcttg cacaccaatt taaaaatggt aactcttgat atatacata 1860
gttataccac tcatgtcagt caataaattt taaggtttaa gtgcaggcct ttgtttacag 1920
aaatccta at tttttgaaac cataactctg acctgacact aaattcctgt agacatgcta 1980
aggaaaatct gcttagtata gagatcaaga acttccattc aaaaagatta ttcagttatg 2040
ttatttgc at attaccattg ttaaaaaataa aaaaattttt aaaagatggc tcaggaatcc 2100
tttcattcta aaatgtttta ttagccttct gcaggctctg ggttggcttt ctggacatgt 2160
ttcccaagtt ggttcaaaac cactattctt ggtcttgata catttgatgat atctgatgta 2220
attgtagatg aaaaagaaaa aatattatgg gaatccttgg tcttttttgt tgctttgaag 2280
agtcccatga agttagttac aggaggtaat aaagtcaaag aatcttttgg ttgaatttta 2340
ggttttaatt ttcttcgggtg ctcttccctt aaaaatttag cacaattaag tgtctctcct 2400
gagagcccta ggtatagtct cattacttct acttctatgt agttctattt tctttacata 2460
actaaaacat actaaagaaa cagatcagta cttccagaca tttagatgtc aggggcaaaa 2520
aatttgtggg gaccactata tggtttgcta gcttgttatg tcaagtttagc ac 2572

```

<210> 37

<211> 3758

<212> DNA

<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1804436CB1

<400> 37

```
ggagctctgg agccttttgc tctcacaata cgagcgggaa ctgcgttgag cgctggattc 60
caggccgaggt gctggcgagg cgcgcagttc tctgctgttt aaaaagtatc cctgtgcttt 120
gaagatactg ctataatttg aaaatttgaa attagtgttt cagctgaacc atccgttcat 180
cttcaagcca tcatgagctg taagaagcag aggtcacgga agcactcagt caatgaaaaa 240
tgtaatatga aaatcgagca ctatttttct cgggtctcta aagagcaaca gaataattgc 300
agtactttct taatgaggat ggagtctaga ggagacccaa gagccacaac taataccag 360
gctcaaagat tccattcacc taagaaaaat ccagaagacc agaccatgcc ccaaatagg 420
acaatatatg ttacettgaa ggtaaaaccac aggagaaacc aagatatgaa acttaagctc 480
acacatagtg agaatagtag cttatatatg gctctcaaca ctctccaggc tgtcagaaaa 540
gagatagaaa ctaccaagg ccaagaaatg cttgtgctgt gcacagaagg aatcaaagag 600
tacataaacc ttggaatgcc cctcagttgt ttccctgaag gtggccagggt ggtcattaca 660
ttttcccaaa gtaaaagtaa gcagaaggaa gataaccaca tatttggcag gcaggacaaa 720
gcatcgactg aatgtgtcaa attttacatt catgcaattg gaattgggaa gtgtaaaaga 780
aggattgtta aatgtgggaa gcttcacaaa aagggcgca aactctgtgt ttatgctttc 840
aaaggagaaa ccatcaagga tgcactgtgc aaggatggca gatttctttc ctttctggag 900
aatgatgatt ggaaactcat tgaaaaaat gacaccattt tagaaagcac ccagccagtt 960
gatgaattag aaggcagata ctttcagggt gaggttgaga aaagaatgggt cccagtgca 1020
gcagcttctc agaatcctga gtcagagaaa agaaacacct gtgtgttgag agaacaaatc 1080
gtggctcagt accccagttt gaaaagagaa agtgaaaaaa tcattgaaaa cttcaagaaa 1140
aaaatgaaag taaaaaatgg ggaacatta tttgaattgc atagaacaac gtttgggaaa 1200
gtacacaaaa attcttcttc gattaaaatg gtgaaacttc ttgtacgtct cagtgaactc 1260
gttggttact tattctggga cagtgcact acgggttacg ccacctgctt tgtttttaa 1320
ggattgttca ttttaacttg tcggcatgta atagatagca ttgtgggaga cggaatagag 1380
ccaagtaagt gggcaaccat aattggtcaa tgtgtaaggg tgacatttgg ttatgaagag 1440
ctaaaagaca aggaacaaa ctactttttt gttgaacctt ggtttgagat acataatgaa 1500
gagcttgact atgctgtcct gaaactgaag gaaaatggac aacaagtacc tatggaacta 1560
tataatggaa ttactctgtt gccacttagt gggttgatac atattattgg ccatccatat 1620
ggagaaaaaa agcagattga gcttctgtct gtgatccctc agggtcagcg agcaagaaa 1680
tgtcaggaac gtgttcagtc taaaaaagca gaaagtccag agtatgtcca tatgtatact 1740
caaagaagtt tccagaaaat agttcacaa cctgatgtga ttacctatga cactgaattt 1800
ttctttgggg ctccggctc ccctgtgttt gattcaaaaag gttcattgggt ggccatgcat 1860
gctgctggct ttgcttatac ttacaaaaat gagactcgta gtatcattga gtttggctct 1920
accatggaat ccactcctct tgatattaag caaagacata aacctggga tgaagaagta 1980
tttgtaaatc agcaggatgt agaaatgatg agtgatgagg acttgtgaga attcagctc 2040
ctggatttaa gggaattggct tatggagttg ttatttcata ggcattgaaa atggttttct 2100
aaactccaaa atgggtcatct tatcaataat aataatattg accatttctt atctgccagg 2160
catttttcta agcacatgaa gaaattagtc ctaacaacac tatgagatgg actataactt 2220
gcccacattt tttttttttt tttgagactg agtctcactc tgtcgctctg gctggagtac 2280
agtgggtgca tctcagctca ctgcaacttc cacctcccag gttcaagcga ttcttatgct 2340
tcagtctcct gagcagctgg gattacaggg aaacgccacc acaccagctt aaaattttt 2400
tttttttttg tttttttagt agagacaggg ttccaccatg ttggtcaggc gggctctgaa 2460
ctcctgacct cgtgatccac ctgcctcggc cttccaaagt gctgggatta caagtttgag 2520
ccactgcacc tggctaactt gccctatttt aaagtcaagc aatgggaaga ataacaagat 2580
tatatagtaa tcagtttcat gacactaaaa gtcatatagt catagggttt tttcatcttt 2640
catatctttg cctaaattca tttgtacag tgcaggaacc aaaacttgtt catctcatga 2700
ttccctacat ctgacataag gaaagtaagt gctcagaaaa atgtgcagggt caataagtgt 2760
caaaagttgg ggctgcaatt aatgctaaca taagagctaa atgcttgatt agaaatgatc 2820
tcaaaacctt ttagaatttc caaaatcttc atattactga aactgtcgga atatatgggt 2880
cctgaaattc agaagatgat agtcactctt cccatattta taggctatta aggcaaggga 2940
tatcttaaac atcatattac tttatttaga tttctactac tccaattatt aatgttatgt 3000
atctctcatt gttttacttc ttcattggtat tatgaagact atatagatga ttcaaccaag 3060
cctgcaaatc tccctcttgt ggaattccac tggacccaat ctgttttcca tttccattgc 3120
aataactact aagccatata atatcaagca ccctccctct aggtccaggg actatcacag 3180
aagaagcagg catgtaagat ttaaggact ggtttcgagg ggttcagagt agggaaacag 3240
cctgttgcat tgtaagagtg atgtcatctt gaagagcagc tggcatgatg actgctgttt 3300
gactcctgca taccaagata ttctgcagca atgtctttaa acagtgcagg tagtacagat 3360
aacccctcat aaagatgctt atctaacctc cccagtgttc aggtgtttca caagaaagtc 3420
tgagatatga ctagctacac gttttgcca aaatgcttgt tatataaagg gtacttttgg 3480
```



```

gaggggtgagt gccgccattt agtgggtgct agaaacattg cttctgtttg taagttccta 3540
ttaaattgttt ctttctgaga aaaaaaagta tatgttttaa aattgacaag gtgcagtcaa 3600
tttctaggaa caggtgacca ctttttcaga gatgaagtgt ttatattaat taaggagcac 3660
ttgggtttctg tatctaataa tagaactgac ttagaagtag cagtaggtga tctccctcta 3720
aagtccgggg gttctgcgcc tgggatctgc caccagct 3758

```

<210> 38

<211> 1036

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7486358CB1

<400> 38

```

cgggcgggcg agccccggcg ccagagcatt ggggtgtctg tacttaggca ggtgctgtgg 60
agccttccta tgagtcccca tgcagccatg cttctgttct actcacagca cttttactcc 120
tgccagctcc tccttttagt ccttgacca tctgtttttg tttccccacc ccactggcat 180
gtccagggat cacttccac tcacacctct tgtacttcta ttcttatttc agggactgat 240
cctgggacac ccgaggaagt ggggtgtctc ctctagccca ctgaggcact agggagcagt 300
ggtggctgga cctcatgcc tgcctctgcc cactggggtc ccttatcttt ctaaccacc 360
cccattggtg ctgttacttc tccctgaaca agagatgatt gaaacagctt tgtctggagg 420
aggggaagcc acatttggga gggacagatt ataagccag agatgaggcc tggaggagg 480
agtttccacc tgagctcaag agtttaaggg ccagaactgt gtgcagtagg ctcctgtggg 540
agtccaggct gggaaggctg gaggttagga gtgcttctct tgccctgcgc catgtggagt 600
ctgccgccga gcagggtctt gtctgtgctg ccactgtctc ttctcttcag cttccagttc 660
ctggttacct atgcttggcg tttccaagag gaagaggagt ggaatgacca aaaacaaatt 720
gctgtttatc tcctccac cctggagttt gccgtgtaca cattcaacaa gcagagcaag 780
gactggtatg cctacaagct ggtgcctgtc ctggcttctt ggaaggagca ggggttatgat 840
aagatgacat tctccatgaa tctgcaactg ggcagaacca tgtgtgggaa atttgaagat 900
gacattgaca actgcccttt tcaagagagc ccagagctga acaataacctg cacctgtctc 960
ttcaccattg gaatagagcc ctggaggaca cggtttgacc tctggaacaa gacgtgctca 1020
ggcgggcatt cctgag 1036

```

<210> 39

<211> 3651

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472344CB1

<400> 39

```

ggaggcgctg cgagcggagc cgcgcgagg ggcgcaccgg ctggtccggg cagcggggggt 60
ttgccgcctt cggggctcca gtccgcgcgc cagtgtctga tgcagtaccg cggggccctc 120
aggtgggcct cggctcggga cgccgggagt cgggaccgcc agtcggggcg ccgggacat 180
ggcgctgcgc gcccgggcgc tgtacgactt caggctcggg aaccccgagg agatctcgct 240
gcgagagcac gaggtgtgta gcctgtgcag cgagcaggac atcgagggtt ggctcgaggg 300
ggtcaacagc cgcgcgacc ggggcctctt cccggcctcc tatgtgcagg tgatecgcg 360
ccccagacct gggcgggcg gagacggcg cccgggcgcc ccggcccgct acgccaatgt 420
gcccccggg ggcttcgagc cctgcctgt cgcgcccccc gcctccttca agcgccgcc 480
tgacgccttc caggcgctgc tgcagccaca gcaggcgcc cctccgagca cttccagcc 540
gcccggcgcg ggcttcccg acggcggggg cgccctgcag ccgtcgctc agcagctcta 600
cggcggttac caggccagcc aaggcagcga tgatgactgg gacgacgagt gggacgacag 660
ctccacgggt gcggacgagc cgggcgtctt gggcagcgga gcatacccg accctcagcg 720
ctcgtcttcg gcgggtgtgg ggcagcgcg ccgctaccgc ctgtccacgc gctccagact 780
gtccctgggc tcccgggcg gctcgggtcc cccgcagcac cacccgctcg ggcccaagag 840
ctcgccacc gtgagccgca acctcaatcg cttctccacc ttctgtcaagt ccggcgggga 900
ggccttcgtg ctgggggagg cgtcaggctt cgtgaaggac ggggacaagc tgtgcgtgg 960
gctggggccc tatggcccc agtggcagga gaaccctac ccgttccagt gcaccatcga 1020
cgacccacc aagcagacca agttcaagg catgaagagc tacatctcct acaagctggt 1080

```

```

gcccacgcac acgcagggtgc cgggtgcatcg gcgctacaag cacttcgact ggctgtacgc 1140
gcgcctggcg gagaagttcc cggatcatctc cgtgccccac ctgcccagaga agcaggccac 1200
cggccgcttc gaggaaggact tcatctcttaa gcgcagggaag ggccctgatct ggtggatgaa 1260
ccacatggcc agccacccag tgctggcgca gtgcgacgtc ttccagcact tcctgacgtg 1320
ccccagcagc accgacgaga aagcctggaa gcagggcaag aggaaggccg agaaggacga 1380
gatgggtgggc gccaaacttct tctgaccct tagcacgcc cccgcccgtg cccttgacct 1440
gcaggaggtg gagagcaaga tagacggctt caagtgcctt accaagaaga tggacgacag 1500
cgcgctgcag ctcaaccaca cggccaacga gttcgcgcgc aagcagggtga cgggcttcaa 1560
aaaggagtat cagaaggtgg gccagtcctt ccgcggcctc agccaggcct ttgagctgga 1620
ccagcaggcc ttctcggtgg gcctgaacca ggctatcgcc ttcaccggag atgcctatga 1680
cgccattggc gagctcttcg cggagcagcc caggcaggac ctggatcccg tcatggacct 1740
attagcgctg tatcaggggc atctggctaa ctccccggac atcatccacg ttcagaaagg 1800
agctctttacc aaagtcaagg agagtaggcg acacgtggag gaaggggaaga tggaggtgca 1860
gaaggctgac ggcattcagg atcgctgtaa cactatttct tttgccactt tggctgaaat 1920
tcaccacttc catcaaattc gagtgcagaga ctttaaatac cagatgcagc atttcttaca 1980
acaacaaata atatttttcc aaaaagttac ccagaagtgg gaagaagctc ttcacaaata 2040
tgatagtgtt taatgactgg acgttggaat atggactttt tcagttcaag gataatttct 2100
acagcagaat aaaaactgct gtcaaagagc tattgccagc tatcagtggt ggtacaagga 2160
cggttttgtg ttcatctgaa acccagctga atttataatt atgtaggaaa taaacagtta 2220
atatggttat ataatagaaa cagtaccaca cattgttaact aaattatact atgtatgcct 2280
acactaccat tgtaactttt ggaataatga ttatactatt tgcccttattg ctttttgaag 2340
tatgggtatt ttagtgcata cttttagtag ctcaaaaccc atgaagggtc tcaaagaagc 2400
tggctggata caagcctgct gtggatgcct ttttactctc atagattggg attacctaaa 2460
ttcaacctat tctctgttta caaactccaa ctagagcagc tatgcgacct tgtgccttta 2520
gactcttggg ttttcatttc tccccgtccc tccccacct ttttaaagta agccacagct 2580
ttcttgattg aaagagtga aggccagtg atataatgac aaactgatga taaccttata 2640
ttggcagtag ggggtggggg gggcggtggg gtgggacgat cagctgtcat caatttgcac 2700
agcaagtatt atctcctgat aagatgctgg tgaatgcagg ggagtgcgat tcattgctca 2760
tctttggata tgaagtctgt tagggaagaa acagtgccac tattccctta gatgcaacag 2820
tagcatagcc tcttcaccca ggctcccaa aagcttggcg tgaagatttc agcaaacatg 2880
tcttacaaca tgaggaggag gagtctaaat cagtcagggg ataaaagtat cgaatcattg 2940
acaacacaca cttggcttta gttcttagga gggttttgtt tttgtttttg tttttaggtt 3000
gaagattttc ttttaattt cagttttttg aaaaaaatg gatctacact gttaactgat 3060
tgagactcca ctgtgattca cttgtttact taaaaacttt tcagggatgt ctgtaaattt 3120
cagtgttaat atgtcatgaa aagtgggtg gattgatcta aggagggacc agaaataatt 3180
tttgctattc caaatactga aggaaaaaga taattgatgt atactatgtt ttaaaaaaaa 3240
aaaagggtatt gatgagcccc cccccccag gacatttaac cttaaaaattt attttaaatg 3300
tattctttta ttattataag ggaaatacag atggctgata aataccaaaa agattcaaaa 3360
gcagcttaat ttaaaaagca caaagagatt ctggcttaca gtgcccacat ctcaatgttt 3420
ttatagttgc tgagctaaact aatgtgatta ttgagtttac agatttaaaa attgtcactg 3480
ttagagtatc tactgttttt atgaagtcaa acttatgctg cctcagaaat ccctgggtac 3540
tgaaatggta acatggaagt gaagaggtca ctttgaaata ttggtgagtc acaaagatta 3600
aagaaaagga tcagtttgca gatactcaga aaagggtatt gaaataatta c 3651

```

<210> 40

<211> 1989

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7192959CB1

<400> 40

```

ggggtctccg gatatcgggg ctatcaagga cgggtctgggg gatgtgggac ttgcgtacgg 60
ggcccaggga tgggtgaagac ggggctgcgg ctggcagaat cagggcaggc agaggctgcg 120
cgccgcggag gctgctggtc cctgacgcct tgacccttcc tctcccagc ccacagccgg 180
gcttggcggg gggaccatgg gcgcctcggt ctccaggggc cgggcccggc ggggtcccgcg 240
ggcggagccg gaaccgaag aggcgctgga cctgagccaa ctaccccag agctgcttct 300
ggtgggtgctg agccacgtcc ccccgcgcac gctgctcggg cgctgccgcc aagtgtgccg 360
gggctggcga gccctgggtg acggccaggc cctgtggctg ctgatectgg ccgcgacca 420
cggcgccacc ggccgcgcgc tgctgcacct cgcccgagc tgccagctc ccgcccgtaa 480
cgccaggcct tgccccctgg gccgcttctg cgcgcgcaga cccatcggac gcaaccttat 540

```

```

tcgcaacccc tgccggccaag aaggcctccg aaagtggatg gtgcaacacg gtgggggacgg 600
ctgggttggtg gaggaanaaca ggacaaccgt gcctggggcc ccttctcaga cgtgcttcgt 660
gacttcattc agctgggtgtt gcaagaagca ggtcttgga ctagaggagg aggggtctgtg 720
gccagaactg ctggatagtg gcaggattga gatttgtgtc tctgactggt ggggagccccg 780
acacgacagc ggctgtatgt acagactcct cgtccaactt ctagacgcca accagactgt 840
tctagataaa ttctctgctg tgccctgatcc catcccgag tggaacaaca atgcctgcct 900
tcacgtcacc cacgtgttct ccaacatcaa gatgggctgc cgcttctgtg ctttcgaaca 960
ccggggccag gacacacagt tctgggctgg ccactatgga gcccgtgtga ccaactccag 1020
tgtgatcgtg cgagtcctgc tgtcctagtc cagcactacc cttcttgcaa gacagcctga 1080
ctgtgccttc cagggcctgg gaccattggc tgggacccct cattaaccaa ccaagcactt 1140
gtacctccct ggcatactgg gaattcctgg gtccaatcaa aggcctgac gggccctgtc 1200
ttcaggttct agaaactacc agaagaagct tcttccatct tatctaccta ctgcagctgc 1260
tgctttgctg gggggcccta ctgtactgag gggaaaccca caagtgaagca tgggggaatg 1320
cccactctgg agaagaattc ttgcccctcc ctctcttctc ctcccaacca aaacctccg 1380
atccagcccc aggcctctct gtaccatccc ctcccctccg caccactgac ctcttgtctc 1440
ctattgtttc tgccacaggg acttcccttg tcccttcagg gaagccctca actcatctct 1500
gaacttagaa tctcatcctt agggctcaga gactccagc cctaagggtg tgaacctcac 1560
ctcctggggc ttccgagcta cagggtcagg accaggtcat gcagctgaaa gtctacgcta 1620
aaaaaanaaa gtttgggcca ggcacagtgg ctcatgcctg taatcccagc actatgggag 1680
gctgaggcag gcggatcacc tgaggctcagg agttcaagac cagcctggcc aacatagtga 1740
aactgtgtct gtgaaggcag gcattgttgt tccactgcgg gatgggatca ggcacagcag 1800
agaatttatc tagaacagtc tgggtggcgt ctagaagttg gacgaggagt ctgtacatac 1860
agccgctgtc gtgtcgggct cccacaccag cagagacaca aatctcaatc ctgccactat 1920
ccagcagttc tggccacaga cctcctcctc ctaggtccaa gacctgcttc ttgcaacacc 1980
agctgaatg

```

<210> 41

<211> 1629

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6169565CB1

<400> 41

```

cgggcgctag ggcgctggca atgtgtagcg gtcacgctgc gcgtaaccac cacacccgcc 60
gcgcttaatg cgccgctaca gggcgcgctc cattcgccat tcaggctgcg caactgttgg 120
gaagggcgat cggtcggggc ctcttcgcta ttacgccagc tggcgaaagg gggatgtgct 180
gcaaggcgat taagttgggt aacgccaggt tttcccagtc acgacgttgt aaaacgacgg 240
ccagtgaatt gaatttaggt gacactatag aagactcaaa ttactccttc ctcccctcta 300
ggtttcctga tttgtaacct ttctaaactc ggtataaagg gggacattaa agtaaagtgc 360
cttatattct tttgcaatac tgccctggcc caggccaagg gcaaccgttg gcctgaaaac 420
ggcacattcg accaccaaat ttttaagggt ttagataact tcatttatcg aaatggcaaa 480
tggcaagaag ttccctcatt tacctttgct cagtccttc cctctgcaa gattgttccc 540
cacacctcta atttgcctc taatctctgt tgttggcata aagggcattc agaaaactcc 600
cctccaaaca ctccccttat attgttccct tcgggacgtc accctcatac attgttccct 660
gttaatccca cattgcccc agcctcttct aagtagagac ctgctgcaca aattgcgtgg 720
cttccctcac ctttgggctc ttggccaaag ccacccctac ttatttttat gccaggaacc 780
caaattctcc ctacctgaag ttaaggagcc aacacctgac cttagcatta taactcaaac 840
aaatcctatt gtttgggtcaa cacaattct gcagtcgtgg cgccccacca ctcccatta 900
aaatttcatt aaagatcct tctcattaga tcacagtcca acagtattct gtcagccctg 960
agaggcttcg gggactcaag cccatcatct ccactgtaga actgccttct tgctcttct 1020
ttccttcctc tttctattcc aatacttget gccaaaacc gggatgagtc ttggggctga 1080
gaactcttgc aaccaggaa gcagtgggcc ccgacagctc atccctggct aattcctgga 1140
tctgagggt tctctggctt cccgcctagt cctcaactc ttctctttt ttaaaagtga 1200
tttgcctgag gtaattgagt gacaggagag gacttcgagg ctgtggccag ggctaccct 1260
ggatgggttc tcaaaactcc cggacccaga attttgctc caaccgctgg atctgggcaa 1320
tttactttct gttttcccc aacggctcca gtttgagagg tcctttgctg gttccttcta 1380
aaacatccaa caccagacac taattagtaa ggtttgccc cactgtctc tcctgtgcta 1440
cccaggagac tcaggtccgc agtcccggtc tcctggggac cattaggacc atcagagacc 1500
ttgggagggt gacgttcagg gatgcctgac ctttccctc agtctacct ctcttgagag 1560
gatctccctg ctctgcctt tttctgggga tgcctagtct aacggcagac accctggcct 1620

```

ttccatcgc

1629

<210> 42

<211> 3166

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7494717CB1

<400> 42

tcttgacag	cgcctctcgc	tggttggggc	gggggtgggc	ggagccagca	ccgtctgggc	60
tgtggaagcg	gaggggtggg	gacactctgg	cccggttctc	ggtggtgcgg	gagcggggcg	120
gagcagcggc	cgctctggtc	ggcggacgtg	ctgccagagta	gtcccgaag	cgaagcagcg	180
atggcggaga	gtccgactga	ggaggcggca	acggcgggcg	ccggggcggc	gggccccggg	240
gcgagcagcg	ttgctgggtg	tggtggcggt	agcggcagcg	gcggcggggt	cgggcccgcct	300
ttcctgcccg	atgtgtgggc	ggcggcggcg	gcagcgggcg	gggcccgggg	cccggggagc	360
ggcctggctc	cgctgcccgg	gctcccgcgc	tcagccgctg	cccacggggc	cgcgctgctt	420
agccactggg	accccacgct	cagctccgac	tgggacggcg	agcgcaccgc	gccgcagtgt	480
ctactccgga	tcaagcggga	tatcatgtcc	atltataagg	agcctcctcc	aggaatgttc	540
gttgtacctg	atactgttga	catgactaag	attcatgcat	tgatcacagg	cccatttgac	600
actccttatg	aagggggttt	cttccctgtt	gtgtttcggt	gtccgcccga	ctatcccatc	660
cacccacctc	gggtcaaact	gatgacaacg	ggcaataaca	cagtgaaggt	taaccccaac	720
ttctaccgca	atgggaaagt	ctgcttgagt	attctaggta	catggactgg	acctgcctgg	780
agcccagccc	agagcatctc	ctcagtgtct	atctctatcc	agtccctgat	gactgagaac	840
ccctatcaca	atgagcccgg	ctttgaacag	gagagacatc	caggagacag	caaaaactat	900
aatgaatgta	tccggcacga	gaccatcaga	gttgacgtct	gtgacatgat	ggaaggaaag	960
tgtccctgtc	ctgaacccct	acgaggggtg	atggagaagt	cctttctgga	gtattacgac	1020
ttctacgagg	tggcctgcaa	agatcgctg	caccttcaag	gccaaactat	gcaggaccct	1080
tttgagagag	agcggggcca	ctttgactac	cagtccctct	tgatgcgcct	gggactgata	1140
cgtcagaaag	tgetggagag	gctccataat	gagaatgcag	aaatggactc	tgatagcagt	1200
tcactctggga	cacagacaga	ccttcatggg	agcctgaggg	tttagaccct	gtccccatct	1260
ccccctcccc	cactcaagag	tcccagcaga	atcccttccc	cccaccccag	ggatggagag	1320
gcactgtgta	tctccctcca	gactcgaagt	catcctgcaa	gatggcaaga	accaagcaag	1380
ctccgatccc	aggggtgtgg	agtggggggc	tgttcccggg	ctgacctcct	tggcactgga	1440
gcactctggg	cttcgttcat	ccattcatcc	cgtatcaggg	gccaaaggtac	ctttacagga	1500
gcacctaag	cgagggcctt	tggcaaaaa	aaaacaacca	acacacctct	ccacagggcc	1560
agtcctctag	ggataagtgg	aagatggaaa	ttgcaattcc	aagagggagt	gtgcccaaat	1620
gatttatggg	gatacctgga	agggagcttg	gggtgggggc	tgtctgtgac	acttaagcag	1680
tctgggtggg	tgtctatttg	tctgtcttca	gtcttgaagc	agggcttccc	aatgcccttt	1740
tcctccctgc	cttccctccc	ccattatttc	ccacaggcca	gcataatttt	gtttttccta	1800
atttatagtc	actgttctag	acagacccaa	gagaaggaac	agtgggtggg	tctaggctgc	1860
tgatcagtaa	gctttacct	gcacctgagc	acctttctcc	cctccccctc	ttcctcacc	1920
ttttctagat	gtaagacaga	aagtaaatgt	gactgggact	taaccaaggt	cttggtaaa	1980
cctgcattgg	accgtaagaa	gctgaaaata	ctgtttgttc	ccgcaatcac	tgatttgaaa	2040
agttcccaac	acaggcagct	gctgtgtata	tgggattaga	gccactacat	agaatagtct	2100
cttacagatt	ttcataaata	ctagtcacaa	taagggtatt	tttcttgggg	gtggagtaag	2160
ggggagactg	atgctagtcc	ttgttgtatt	ttgttgggct	gtccttgtgt	attttcaccc	2220
cagcctgtag	tcctcctcac	ttcaacccca	gggatttttg	gggagcaagg	gtagccaatg	2280
gcagaggggg	ttggggctgg	gactctggag	gtcctcctcc	ttctttctct	tccttcggcc	2340
tccccctgac	ccccagctgc	tcttgctact	gtctctgatg	gggtatttgc	tggctttgtt	2400
gcttctctat	ctgtatttag	ctgcagtgat	ccttttagctg	gttggctcag	aaaaaaaaaa	2460
atgtgcttta	ggtgccctgt	aatcctgggc	atcaaggga	tecatccttc	ccctttttga	2520
tatgttctcc	ccgtacttcc	agatttattg	ttatggctcc	cagtgggtat	tggcgattct	2580
tgtgatgcag	ggcctcagtc	agtgtccagc	catgcataag	ggagaggata	gtgtgtacct	2640
gccctgccct	ctgctatgaa	ggtctctgcc	ttgtggatca	tgggactccc	cttggaggat	2700
ctgtgcaaa	gggggctggg	cacaaaggag	aatgtcctat	ttgggagggc	aggaagcaaa	2760
ggaactggac	agggattggg	gggcttgggg	aacggaagtt	tatcttggat	acccttgaag	2820
aggctgggtc	tcttcacatg	aagatcgaaa	agggaccctg	cttccaattt	ccctcttcca	2880
ttcctcgagc	tactccaggg	cttagaagaa	tgctcttggg	ctgtgggtcc	agtgttgtct	2940
gtcatccatt	taagtgttcc	cactttcaag	tgacaatcct	ctccttggcc	ctgccatagg	3000
gcagagcatg	tctggcatag	cagcctgact	tttatgccct	aatcttgagt	tgaggaaata	3060

tatgcacagg agtcaaagag atgtctttat atctgactgt atataaatga agtttttttg 3120
 ttttttttgt tttccttttt ggtgcataaa gtttggtttg gcagaa 3166

<210> 43

<211> 399

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7497510CB1

<400> 43

ctcctccctt tggctccctt tctccctaaa atggcaaccc caggctgagc cactggctga 60
 gtggcaccat gcagcttcag gcctctctct cgtttctcct gattctcact ctctgcctag 120
 agcttcgata agaactagca cgagacacta tcaaggatct cctcccaaat gtatgcgctt 180
 ttcctatgga aaagggccct tgtcaaacct acatgacgag atgggttttc aactttgaaa 240
 ctgggtgaatg tgagttattt gcttacggag gctgaggagg caacagcaac aactttttga 300
 ggaaagaaaa atgtgagaaa ttctgcaagt tcacctgatt ttctaacaag aacacagccc 360
 tccatggatt cgggattgct ctgagggccca tagaaggca 399

<210> 44

<211> 1811

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7498882CB1

<400> 44

ggtttccggg ccggcgctact atttcaaggc gcgcgcctcg tgggtggactc accgctagcc 60
 cgcagcgctc ggcttcctgg taattcttca cctcttttct cagctccctg cagcatgggt 120
 gctgggcccct ccttgctgct cgccgccttc ctgctgcttc tctccggcga cggcgccgtg 180
 cgctgcgaca cacctgccaa ctgcacctat cttgacctgc tgggcacctg ggtcttccag 240
 gtgggctcca gcggttccca gcgcgatgtc aactgctcgg ttatgggacc acaagaaaaa 300
 aaagtagtggt tgtaccttca gaagctggat acagcatatg atgaccttgg caattctggc 360
 catttcacca tcatttacaa ccaaggcctt gagatttgtt tgaatgacta caagtgggtt 420
 gcctttttta agtataaaga agagggcagc aaggtgacca cttactgcaa cgagacaatg 480
 actgggtggg tgcataatgt gttgggcccg aactgggctt gtttcaccgg aaagaagggtg 540
 ggaactgcct ctgagaatgt gtatgtcaac acagcacacc ttaagaattc tcaggaaaaag 600
 tattctaata ggctctacaa gtatgatcac aactttgtga aagctatcaa tgccattcag 660
 aagtcttgga ctgcaactac atacatggaa tatgagactc ttacctggg agatatgatt 720
 aggagaagtg gtggccacag tcgaaaaatc ccaaggccca aacctgcacc actgactgct 780
 gaaatacagc aaaagatttt gcatttgcca acatcttggg actggagaaa tgttcatggt 840
 atcaattttg tcagtcctgt tcgaaaccaa ggctgtgaag gcggcttccc ataccttatt 900
 gcaggaaagt acgcccaga ttttgggctg gtggaagaag cttgcttccc ctacacaggc 960
 actgattctc catgcaaaat gaaggaagac tgctttcgtt attactcctc tgagtaccac 1020
 tatgtaggag gtttctatgg aggtgcaat gaagccctga tgaagcttga gttgggtccat 1080
 catgggcccc tggcagttgc ttttgaagta tatgatgact tcctccacta caaaaagggtg 1140
 atctaccacc aactggtct aagagaccct ttcaaccctt ttgagctgac taatcatgct 1200
 gttctgcttg tgggctatgg cactgactca gcctctggga tggattactg gattgttaaa 1260
 aacagctggg gcaccggctg ggggtgagaat ggctacttcc ggatccgcag aggaactgat 1320
 gagtgtgcaa ttgagagcat agcagtgga gccacaccaa ttctaaatt gtagggtatg 1380
 ccttccagta tttcataatg atctgcatca gttgtaaagg ggaattggta tattcacaga 1440
 ctgtagactt tcagcagcaa tctcagaagc ttacaaatag atttccatga agatatttgt 1500
 cttcagaatt aaaactgcc ttaattttta tatacctttc aatcgccac tggccatttt 1560
 tttctaagta ttcaattaag tgggaatttt ctggaagatg gtcagctatg aagtaataga 1620
 gtttgcttaa tcatttgtaa ttcaaacatg ctatattttt taaaatcaat gtgaaaacat 1680
 agacttattt ttaaattgta ccaatcacaa gaaaataatg gcaataatta tcaaaacttt 1740
 taaaatagat gctcatattt ttaaaataaa gttttaaaaa taactgcaaa aaaaaaaaaa 1800
 aggggggggc g 1811

<210> 45
 <211> 4407
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5524205CB1

<400> 45
 cagtgtgctg gaaagtggac agatatactg cattttttaa acagagtggg aaattttcctg 60
 gaaatccctg gcctccat at aagaaaagga catcactcca tcctagctat aaaggtctta 120
 tgagactttg cactgtaaaa ctttacacat tgtgctattc actctgcttt acaagatttt 180
 gatggatcat caaaatctgt cagaacatgt actctgcatg gttttatatac tgattgaatt 240
 aggacttgaa aattctgctg aagaagaatc agatgaagag gcatcagtgg gtggaccaga 300
 acgttgctcat gacagtttgt ttctggcag taacttagtg tcaaacatgc gacactttat 360
 aaactatgtt agagtaagag ttccagagac tgctcctgaa gtaaagagag actcacctgc 420
 aagtactagc tctgataact tgggttcttt acaaaattct ggtacagctc aagttttcag 480
 tttagtagca gaacgtagaa agaaatttca ggaaatcatc aatcgcagta gcagtgaagc 540
 aaatcagggtg gttcgtccca aaacttcaag taaatgggtc gtcctgggtt cagctccaca 600
 gttaactaca gccatttttg aaattaaaga aagcatattg tctttgctaa ttaaacttca 660
 ccacaaaactc tcaggaaaac aaaactccta ctatcctcct tggcttgatg acatagaaat 720
 tttaatccaa ccagaaattc cttaatacag tcatggagat ggtataactg ccgtggaaag 780
 aattttacta aaagctgcat cgcaagtag aatgaacaaa cgcacattg aagagatatg 840
 tagaaaagtg acccctcctg taccacctaa aaaagtcact gcagcagaga agaaaacatt 900
 ggacaaagaa gaaagggcag agaaggctag agagaggcag cagaaattgc ttgaggagtt 960
 tgcttcacga cagaaaagct ttatggaaac tgcaatggat gttgattctc ctgagatga 1020
 tattcctatg gagatcacca cggcagaacc acagggttcc gaggcagtat atgactgtgt 1080
 tatttgtgga cagagtggcc cctcctctga agatcgacct actggattag ttgtactgtt 1140
 acaagcatcc tcagttttgg ggcagtgcg tgacaatgtt gagccaaaaa agttgccgat 1200
 cagtgaagag gagcagattt acccttggga tacctgtgca gccgttcattg atgtgaggct 1260
 ttcattatta cagcgttatt ttaaggatag ttcattgtctc ttggcagtat caattggctg 1320
 ggaaggaggt gtttatgtac aaacctgtgg tcacacatta catatagatt gtcataaact 1380
 ttacatcgaa tcattacgga atgaccaggt tcttcagggc ttctcgtggg acaaaggaga 1440
 attcacgtgt ccactctgta ggcagtttgc taacagtgtt cttccattgt atcctggaag 1500
 caatgtggaa aataaccctt ggcaacgtcc tagcaacaaa agcatacaag atctcataaa 1560
 ggaagtggag gagctgcagg gacgaccggg agctttccca tcagaaacaa atttaagtaa 1620
 agaaatggaa tctgtaatga aagatataaa aaataccact cagaagaaat atagagacta 1680
 tagcaagacc cggggtcac cagacaatga ttttctcttt atgtactctg ttgctagaac 1740
 caatttagaa cttgaattga ttcacgagg aggcaatttg tggtcagggt gtgcaagcac 1800
 agctggcaaa aggtcttgtt taaatcagct gtttcatgta ttagccttgc acatggcgtt 1860
 ttatagcatt gactctgagt ataaccctg gagaaagctc acccagttag aagagatgaa 1920
 tccacagctg ggatatgaag aacaacagcc tgagggttcca attctttatc atgatgtaac 1980
 atcccttttg ctcatccaga tcttaatgat gccacaaccc ttacgcaaaag accactttac 2040
 ctgcattgtg aaggtaactt ttaccctact gtacacacag gctcttgca cactctcagt 2100
 taaatgcagc gaagaagata ggtcagcctg gaaacacgcg ggagctctca aaaagagtag 2160
 atgtgatgca gaaaagtctt acgaagtatt actgagcttt gtgataagtg aactatttaa 2220
 aggaaagtta taccatgaag aaggaactca ggaatgtgca atggttaacc ctattgcttg 2280
 gtctcctgaa tccatggaaa aatgcttaca ggacttctgc ttaccttttc tcagaatcac 2340
 cagccttctt cagcaccacc tttttgggga agatttacct agctgccagg aagaagaaga 2400
 attttcagtt cttgccagct gcctgggact tctgccaacg ttttaccaaa cagaacatcc 2460
 atttcactagt gcctcctgtc tggattggcc agttccagca tttgatatta taactcagtg 2520
 gtgttttgag ataaaatcat ttactgaaag acatgcagaa caaggaaagg ccttgcttat 2580
 ccaagagtca aaatggaaat taccacacct actacagttg cctgagaatt ataaccacct 2640
 ttttcagtac taccacagaa aaacctgtag tgtctgcacc aagggttcta aagatcctgc 2700
 tgtttgcctt gtgtgtggta cttttgtatg cctgaaagga ctttgctgca agcaacaaag 2760
 ttactgtgaa tgtgtactgc actctcagaa ctgtgggtgca ggaacaggta ttttcccttt 2820
 gatcaatgca tcggttaatta tcatcattcg aggtcaccgc ttctgcctct ggggttccgt 2880
 gtatttggat gctcatggag aggaagaccg ggactcttagg cgaggcaaac ctctctacat 2940
 ttgtaaggaa agatacaaa ttcttagaca acagtggatt tctcatactt ttgatcacat 3000
 caataaaaga tggggtccac attacaatgg gctgtgactc tccacctcag cattgcatcg 3060
 tatcatcatt ttcgctacga atttattttt caacaataag ctttaactta atttggggga 3120
 ttaacacttt tgctgaggga gaaaaagaaa acatacatta tgaagccttt ccaaaattag 3180

gtgcttggtgta	atcacggttaa	tgggtataatt	tttttttttt	aatatctgga	gaacattaat	3240
aacaagtttaa	attattcttt	agtggtcatt	ttttaagtgc	acaattaata	agaagcacaa	3300
cttggttcaca	aactcattca	gaaatgattc	tcccaacaat	gcataatcagc	tattcattga	3360
tacttagagt	gggtgtgatt	tatttgacat	tttactgctt	ctttctgtct	gtgtgtttta	3420
atttgcatct	gccaaagcata	atgcatcttt	tttctctgct	cattcttgtg	ttgattggag	3480
aatttttctg	tatgtaatta	gaaaaaatg	taaaacatga	tttatgtgaa	atactgtata	3540
gtaaaagttg	gtctaatagt	agaacttta	aattttttct	tattgtgagg	aatctgttaa	3600
aagtttaaaag	ctttgctgaa	aactgaattc	attctcagga	atttcataaa	tcttctcccc	3660
aggtaaataa	ttgaaatagc	tgtaaaaata	gtagatagct	gctgttaata	taatacagta	3720
cattttgggg	ggcataatgtg	tggttggggg	gtccttaaaa	atcaaaattt	gccatttcag	3780
ttggatgaat	tactagaggt	aataacaaat	cttactataa	aatcaagagg	tttaagaaca	3840
tacactgggc	agatgttgat	tccgtgcatg	cccacctttt	attaccaaac	aagggtttgt	3900
ttatatgatt	gtattagaaa	tgctcagact	tccccagaaa	tgaaccataa	attttggaa	3960
ttcctttcag	ctcaagaggt	tcagctatat	tgtattttgtg	cagtgtaatc	actactattt	4020
ctgctcggtt	tcctaaaagg	aaaaaaaagg	cgcagtggtg	atgaccctca	tgaatgagcc	4080
acgcttctgc	attcttctta	gaaactgctg	tgaaaaacaa	tttatgtttg	cagggtttaa	4140
aaatcagtaa	aaatgggaat	gattgagcta	aaacccactc	tatgagaagg	aagattactg	4200
aaaagcatgt	gacatattgc	tacaaagatt	ttttttccta	aatgattcag	taattgaatg	4260
attatttta	atatagtgtc	atcaagcaat	cccttggtact	ttggacttcc	atggcttggt	4320
atataaaatt	acattttttac	atgtaaaaat	aaactaaaca	aatctaata	taaaatataa	4380
acataatgtc	agatccatgt	tctatac				4407

<210> 46

<211> 2177

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7102342CB1

<400> 46

agcggaattc	ccgacttccc	aacggcttcc	cgctggcagc	cccgaagccg	caccatgttc	60
cgcctctggt	tgctgctggc	cggtctctgc	ggcctcctgg	cgtcaagacc	cggttttcaa	120
aattcacttc	tacagatcgt	aattccagag	aaaatccaaa	caaatacaaa	tgacagtcca	180
gaaatagaat	atgaacaaat	atcctatat	attccaatag	atgagaaaact	gtacactgtg	240
caccttaaac	aaagatattt	tttagcagat	aattttatga	tctatttcta	caatcaagga	300
tctatgaata	cttattcttc	agatattcag	actcaatgct	actatcaagg	aaatattgaa	360
ggatattccag	attccatggt	cacactcagc	acgtgctctg	gactaagagg	aatactgcaa	420
tttgaataatg	tttcttatgg	aattgagcct	ctggaatctg	cagttgaatt	tcagcatggt	480
ctttacaaat	taaagaatga	agacaatgat	attgcaattt	ttattgacag	aagcctgaaa	540
gaacaaccaa	tggatgacaa	cattttttata	agtgaataat	cagaaccagc	tggtccagat	600
ttatttcttc	ttatctaga	aatgcataat	gtggtggaca	aaactttgta	tgattactgg	660
ggctctgata	gcatgatagt	aacaaataaa	gtcatcgaaa	ttgttggcct	tgcaaattca	720
atgttaccce	aatttaaagt	tactattgtg	ctgtcatcat	tggagttatg	gtcagatgaa	780
aataagattt	ctacagttgg	tgaggcagat	gaattattgc	aaaaattttt	agaatggaaa	840
caatcttate	ttaacctaa	gctcatgat	attgcataat	tactaattta	tatggattat	900
cctcggttatt	tgaggagcag	gtttcctgga	acaatgtgta	ttactcggtta	ttctgcagga	960
gttgcatgtc	aatgtggacc	tgcaagctgt	tgtgattttc	gaacttgtgt	actgaaagac	1020
ggagcaaaat	gttataaagg	actgtgctgc	aaagactgtc	aaattttaca	atcaggcggt	1080
gaatgtaggc	cgaaagcaca	tctgaatgt	gacatcgctg	aaaattgtaa	tggaagctca	1140
ccagaatgtg	gtcctgacat	aactttaatc	aatggacttt	catgcaaaaa	taataagttt	1200
atttgttatg	acgggagactg	ccatgatctc	gatgcacggt	gtgagagtgt	atttggaaaa	1260
ggttcaagaa	atgctccatt	tgctgctat	gaagaaatac	aatctcaatc	agacagattt	1320
gggaactgtg	gtagggatag	aaataacaaa	tatgtgttct	gtggatggag	gaatcttata	1380
tgtggaagat	tagtttgtac	ctaccctact	cgaagccctt	tccatcaaga	aaatgggtgat	1440
gtgattttatg	ctttcgtagc	agattctgta	tgcataactg	tagactacaa	attgcctcga	1500
acagttccag	atccactggc	tgtcaaaaa	ggctctcagt	gtgatattgg	gaggggtttgt	1560
gtaaatcgtg	aatgtgtaga	atcaaggata	ataaggctt	cagcacatgt	ttgttcacaa	1620
cagtgttctg	gacatggagt	gtgtgattcc	agaaacaagt	gccattgttc	gccaggctat	1680
aagcctccaa	actgccaaat	acgttccaaa	ggattttcca	tatttctctga	ggaagatatg	1740
ggttcaatca	tggaaagagc	atctgggaag	actgaaaaca	cctggcttct	aggtttctct	1800
attgctcttc	ctattctcat	tgtaaacaac	gcaatagttt	tggcaaggaa	acagttgaaa	1860

```

aagtggttcg ccaaggaaga ggaattccca agtagcgaat ctaaatacgga aggtagcaca 1920
cagacatatg ccagccaatc cagctcagaa ggcagcactc agacatatgc cagccaaacc 1980
agatcagaaa gcagcagtca agctgatact agcaaatacca aatcacagga cagtacccaa 2040
acacaaagca gtagtaacta gtgattcctt cagaaggcaa cggataacat cgagagtctc 2100
gctaagaaat gaaaattctg tctttccttc cgtggtcaca gctgaaagaa acaataaatt 2160
gagtgtggat ctaaaaaa                                     2177

```

<210> 47

<211> 703

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4169939CB1

<400> 47

```

atgatgctcc ggctgctcag ttccctcctc cttgtggcgg ttgcctcagg ctatggccca 60
ccttcctctc actcttcag cgcgcttgct catggtgagg atgcgatccc catcaactct 120
gaggagctgt ttgtgcatcc actctggaac cgctcgtgtg tggcctgtgg caatgacatc 180
gccctcatca agctctcacg cagcgcacag ctgggagatg ccgtccagct cgctcactc 240
cctcccgctg gtgacatcct tcccaacaag acaccctgct acatcacagg ctggggccgt 300
ctctatacca atggggccact ccagacaag ctgcagcagg cccggctgcc cgtggtggac 360
tataagcact gctccagggt gaactggtgg ggttccaccg tgaagaaaac catggtgtgt 420
gctggagggt acatccgctc cggctgcaac ggtgactctg gaggaccct caactgcccc 480
acagaggatg gtggtggca ggtccacggt gtgaccagct ttgtttctgg ctttggctgc 540
aacttcatct ggaagccac ggtgttcaact cgagtctccg ccttcatcga ctggattgag 600
gagaccatag caagccacta gaaccaaggc ccagctggca gtgctgatcg atccacatc 660
ctgaataaag aataaagatc tctcagaaaa aaaaaaaaaa aaa 703

```

<210> 48

<211> 1295

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6539977CB1

<400> 48

```

ggggaatgcc agttctggca ccaaccttcc tgtccctgc tggggcctct gctcccccat 60
ctctcaggag tcgaaagtga gaaagcaaga catcaaggag ggacctgtgc cctgctccac 120
atcctccac ctgcgcggcg cagagcctgc agggcccgcc cccctcgtct ctggtcccta 180
cctctctgct gtgtcttcat gtccctgagg gtcttggggt ctgggacctg gccctcagcc 240
cctaaaaatgt tctcctctgct gacagcactt caagtccctg ctatagccat gacacggagc 300
caagaggatg agaacaagat aattggtggc tatacgtgca cccggagctc ccagccgtgg 360
caggcggccc tgetggcggg tcccaggcgc cgcttctct gcgaggcgc cctgctttca 420
ggccagtggg tcatcactgc tgetcactgc ggccgcccga tcttcagggt tgccctgggc 480
aagcacaacc tgaggagggt ggaggccacc cagcagggtc tgcgcggtgt tcgtcagggtg 540
acgcacccca actacaactc ccggaccac gacaacgacc tcatgctgct gcagctacag 600
cagcccgcac ggatcgggag ggcagtcagg cccattgagg tcaccaggc ctgtgccagc 660
cccgggacct cctgcgaggt gtcaggctgg ggaactatat ccagcccat cgccaggtag 720
ccgcctctc tgcaatgcgt gaacatcaac atctccccg atgagggtgt ccagaaggcc 780
tattctagaa ccatcacgcc tggcatggct tgtgcaggag ttccccaggg cgggaaggac 840
tcttgtcagg gtgactctgg gggaccctg gtgtgcagag gacagctcca gggcctcgtg 900
tcttggggaa tggagcgctg cgccctgcct ggctaccccg gtgtctacac caacctgtgc 960
aagtacagaa gctggattga ggaaacgat cgggacaaat gatggtcttc acggtgggat 1020
ggacctcgtc agctgcccag gccctcctct ctctactcag gaccaggag tccaggcccc 1080
agccctcct cctcagacc caggagtcca gggcccccag cctcctccc tcagaccgg 1140
gagtcaggc cccagcccc tctcctcct gaccaggag tccaggcccc agccctcct 1200
ccctcagacc cgggagtcca gggcccccag cctcctccc tcagaccagc gagtccaggc 1260
cccagtcct cctccctcag acccaggagt ccagg                                     1295

```


<210> 49
 <211> 575
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7675588CB1

<400> 49
 ccgggaggtc tgggtgttgg ctgggtccatt ccaggacagc tacatctttg gcagacctgg 60
 tgctgacaga accagctctg atcaggaggc cagtacttcc taaaatggga ctctcaggac 120
 ttctgccaat cctggtacca ttcatacctt tgggggacat ccaggaaacct gggcacgctg 180
 aaggcatcct tggcaagccg tgtcccaaaa tcaaagtgga atgcgaagtg gaagaaatag 240
 accagtgtac caaaccaga gattgcccag aaaacatgaa gtgttgcccg ttcagccgtg 300
 gaaagaaatg tttagacttc agaaaggtca gccttacttt ataccataag gaggagcttg 360
 aataacctcc aggatttggc tcataatcca ggcctctctc cacgtgtgcc tgattgatgc 420
 tccaaattgg cttccacggg ccaaaccttg gctgttccag aaactgaacc ccaggaattg 480
 cttacacact ttcttccagc gtagcatctc ttcaaacaca atgctcttcc ccttgaccac 540
 ttctcagtat gaaactctat gtcttcgggt cgttg 575

<210> 50
 <211> 1062
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6244077CB1

<220>
 <221> unsure
 <222> 33-34
 <223> a, t, c, g, or other

<400> 50
 ataaatttag agagacgtat ggccctcgag canngaattc ggcacgagge ctccagtcce 60
 tgagaattgg tactacgaaa aggtgaactc ctgggcagaa tcttgccatg agcttgccga 120
 gtccagccag gcccttgcgt aaggggccca gaccacgggc cacttctccc cgttccatct 180
 gaccagctgg gcccttgcgc ccacctggcc tccacgttcc ctctctcttc acccacaccc 240
 ctggccatgg ctaactacta cgaagtgcgt ggctgtcagg ccagcgcttc cccggaggac 300
 atcaagaaag cctaccgcaa gctggccctt cgttggcacc ccgacaagaa ccctgacaat 360
 aaggaggagg cggagaagaa gttcaagctg gtgtctgagg cctatgaggt tctgtctgac 420
 tccaagaaac gctccctgta tgaccgtgct ggctgtgaca gctggcgggc tgggtggcggg 480
 gccagcagc cctaccacag ccccttcgac accggctaca ccttcgtaa ccctgaggac 540
 atcttccggg agtttttctg tggcctggac cctttctcct ttgagttctg ggacagccca 600
 ttcaatagtg accgtggtgg ccggggccat ggcctgaggg gggccttctc ggcaggcttt 660
 ggagaatttc cggccttcat ggaggccttc tcatccttca acatgctggg ctgcagcggg 720
 ggcagccaca ccaccttctc atccacctcc ttggggggct ccagttcttg cagctcgggg 780
 ttcaagtcgg tgatgtcgtc caccgagatg atcaatggcc acaaggtcac caccaagcgc 840
 atcgtggaga acgggcagga gcgcgtggag gtggaggag acgggcagct caagtcggtg 900
 actgtgaacg gcaaggagca gctcaaattg atggacagca agtaggcgct ggccaccggg 960
 ccctgccttc ccaccaccac caccgtgcat ggggcagcaa acacgtgggg ccgcagacat 1020
 agcctgatgg ttaataaatg tgccaagtga gttcatggca aa 1062

<210> 51
 <211> 1029
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7498404CB1

<400> 51

```

ggcttcgcgc atctggctca gttccgccat ggctccttg gaagtcagtc gtagtcctcg 60
caggtctcgc cgggagctgg aagtgcgcag tccacgacag aacaaatatt cgggtgctttt 120
acctacctac aacgagcgcg agaacctgcc gctcatcgtg tggctgctgg tgaaaagctt 180
ctccgagagt ggaatcaact atgaaattat aatcatagat gatggaagcc cagatggaac 240
aagggatggt gctgaacagt tggagaagat ctatgggtca gacagaattc ttctaagacc 300
acgagagaaa aagttgggac taggaactgc atatattcat ggaatgaaac atgccacagg 360
aaactacatc attattatgg atgctgatct ctacacccat ccaaaattta ttcttgaatt 420
tattaggaag caaaaggagg gtaattttga tattgtctct ggaactcgtc acaaaggaaa 480
tggaggtgta tatggctggg atttgaaaag aaaaataatc agattatacc gaaaagaagt 540
tctagagaaa ttaatagaaa aatgtgtttc taaaggctac gtcttccaga tggagatgat 600
tggtcgggca agacagttga attatactat tggcgagggt ccaatatcat ttgtggatcg 660
tgtttatggt gaatccaagt tgggagggaa tgaaatagta tctttcttga aaggattatt 720
gactcttttt gctactacat aaaagaaaaga tactcattta tagttacgtt catttcaggt 780
taaactatgaa agaagcctgg ttactgattt gtataaaatg tactctttaa gtataaaata 840
taaggtaagg taaatttcat gcatcttttt atgaagacca cctattttat atttcaaatt 900
aaataatttt aaagttgctg gcctaagtga caatgttctc aattttcgtt ttcatatttg 960
tgtattgaga cctataaata aatgtatat tttttttgca aaaaaaaaaa aaaaaaaaaa 1020
aaaaaaaaa

```

<210> 52

<211> 905

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7391748CB1

<400> 52

```

tggggctcaa aatataaact caggctatth atcaacttaa tctggggaag caaacctgaa 60
ggcaagtacc accctgtcat ccttagctca gagctgctga gaaagaggat acagctgagc 120
cccaggggcc toccattccc tgcattctgg ttagctgcag tcttgccctc cccgtgctgt 180
ctgcctaccc tgcagagctg gtggaccata gctcctgcag cccagaccta cctcttgctt 240
ttgcagcaat ataaatgtca ccctgggccc ccacaatata cagagacggg aaaacaccca 300
gcaacacatc actgcgcgca gagccatccg ccacctcaa tataatcagc ggaccatcca 360
gaatgacatc atgttattgc agctgagcag aagagtcaga cggaaatcga acgtgaaccc 420
agtggctctg cctagagccc aggagggact gagacccggg acgctgtgca ctgtggccgg 480
ctggggcagg gtcagcatga ggaggggaa agatacactc cgagagggtg agctgagagt 540
gcagagggat aggcagtgcc tccgcatctt cggttcctac gacccccgaa ggcagatttg 600
tgtgggggac cggcggggaa ggaaggctgc cttcaagggg gattccggag gccccctgct 660
gtgtaacaat gtggcccaag gcatcgtctc ctatggaaag tcgtcagggg ttctccaga 720
agtcttcacc aggggtctca gtttcctgcc ctggataagg acaacaatga gaagcttcaa 780
actgctggat cagatggaga cccccctgtg actgactctt cttctcgggg acacaggcca 840
gtccacagat gttgccagag ccttaataaa cgtccacaga gtataaataa aaaaaaaaaa 900
aaaaa

```

<210> 53

<211> 2667

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7499780CB1

<400> 53

```

agtcctgtct cccgcccgcg ggccgagcgc cgcccgtgcc ccgcctcccg tgcgcccggg 60
acaatcctcg ccttgtctgt ggccgaggca tctggagctt tctgtagcct ccggatacgc 120
ctttttttca gggcgtagcc ccagccaaagc tgctccccgc ggccggccgca caagcagccc 180
gagcgcctcc tttccagagc tcccctccgg agctgggata caggcgcgta gccgagatcc 240
caggatcctg ggtgctgtct gggcccgcgc ccacacatga cctcctcggg gcctggaccc 300
cggttcctgc tgctgctgcc gctgctgctg cccctgcggg cctcagcctc cgaccggccc 360

```

```

cggggcccag acccgggtcaa cccagagaag ctgctggtga tcaactgtgc cacagctgaa 420
accgaggggt acctgcgctt cctgcgctct gcgagattct tcaactacac tgtgcccagg 480
ctgggacctg gagaggagtg gcgagggggg gatgtggctc gaacagtttg tggaggacag 540
aagggtccgtt gggttaaagaa ggaaatggag aaatacgtg accgggagga tatgatcatc 600
atgtttgttg atagctacga cgtgattctg gccggcagcc ccacagagct gctgaagaag 660
ttcgtccaga gtggcagccg cctgctcttc tctgcagaga gcttctgctg gcccagtggt 720
gggctggcgg agcagtacct tgaggtgggc acggggaagc gcttctcaa ttctggtgga 780
ttcatcggtt ttgccaccac catccaccaaa atcgtgcgcc agtggaagta caaggatgat 840
gacgacgacc agctgttcta cacacggctc tacctggacc caggactgag ggagaaactc 900
agccttaac tggtatcataa gtctcggatc ttccagaacc tcaacggggc tttagatgaa 960
gtggttttaa agtttgatcg gaaccgtgtg cgtatccgga acgtggccta cgacagctc 1020
cccattgttg tccatggaaa cgggtccact aagctgcagc tcaactacct gggaaactac 1080
gtccccaatg gctggactcc tgagggaggc tgtggcttct gcaaccagga cggaggaca 1140
ctcccggggg ggcaggaggt cttccatgaa cccacatcg ctgactcctg gccgcagctc 1200
caggaccact tctcagctgt gaagctcgtg gggccggagg aggtctctgag cccaggcgag 1260
gccagggaca tggccatgga cctgtgtcgg caggaccccg agtgtgagtt ctacttcagc 1320
ctggagcgcc acgtgtctct caccaacctg cagaccctgc gtatcctcat tgaggagaa 1380
aggaaggtga tcgcccccat gctgtccgc caggccaagc tgtggtccaa cttctggggc 1440
gccctgagcc ccgatgagta ctacgcccgc tccgaggact acgtggagct ggtgcagcgg 1500
aagcagtggt gtgtgtgaa tgtaccatac atctcccagg cctatgtgat ccggggtgat 1560
accctgcgga tggagctgcc ccagagggat gtgttctcgg gcagtgcac agaccgggac 1620
atggccttct gtaagagctt tccagacaag ggcattctcc tccatctgag caatcagcat 1680
gaatttgcc ggctcctggc cactccaga tacgacacgg agcacctgca cccgacctc 1740
tggcagatct tcgacaaccc cgtcgactgg aaggagcagt acatccacga gaactacagc 1800
cgggcccctg aaggggaagg aatcgtggag cagccatgcc cggacgtgta ctggttccca 1860
ctgctgtcag aacaaatgtg tgatgagctg gtggcagaga tggagcacta cggccagtgg 1920
tcaggcggcc ggcagtgagg ttcaaggctg gctggaggct acgagaatgt gccaccctg 1980
gacatccaca tgaagcaggt ggggtacgag gaccagtggc tgcagctgct gcggacgtat 2040
gtgggccccca tgaccgagag cctgtttccc ggttaccaca ccaaggcgcg ggcggtgatg 2100
aactttgttg ttgcgtaccg gccagacgag cagcgtctc tgccggccca ccacgactca 2160
tccacctta ccctcaacgt tgccctcaac cacaagggcc tggactatga gggaggtggc 2220
tgccgcttcc tgcgtacga ctgtgtgatc tccctcccga ggaagggtg ggcactcctg 2280
caccgccggc gcctcaccca ctaccacgag gggctgcca cgacctgggg cacacgctac 2340
atcatggtgt cttttgtcga cccctgacac tcaaccactc tgccaaacct gccctgccat 2400
tgtgcctttt tagggggcct ggcgcccgtc ctgggagttg ggggatgggt ctctctgtct 2460
ccccacttcc tgagttcatg ttccgcgtgc ctgaactgaa tatgtcacct tgctcccaag 2520
acacggccct ctcaggaagc tcccggagtc cccgcctctc tccctcgccc acaggggttc 2580
gtgggcacag ggttctggg gactccccgc gtgataaatt attaattgtc cgcagctca 2640
ctctgaataa aggacagttt gtaaaaaa

```

<210> 54

<211> 2977

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7499881CB1

<400> 54

```

gtgcggagtg ccgagcggcc tcacccccaa ccgtcggccc agtcggacgg ttccgaggcg 60
ttgccgggag cggggcggcg ctctgtgtgg actcggagaa acgcggggcg tctgcctgag 120
cccgcttttc tacaagatgt ggggattttt gaagcgccct gtagtggtga cggctgacat 180
caacttgagc cttgtggccc tgactgggat ggggttactg agccggctgt ggcgactcac 240
ctaccgcggg gctgtggtgt tatttaggag gattcgatgg caattttttg tggaaacagaa 300
ttggagcaga atacagtagc aacgtgcctg tgtggtccct gcgcctgctg ccagcactcg 360
cgggggcctt gtccgtcccc atggcctacc agatagtgtt ggagctccac ttttctcatt 420
gtccggccat gggagctgct ctgttgatgc ttatcgagaa tgctctcatc actcagtaaa 480
ggctaattgt tttggaatca gtgttaatat attggccgtg ttgtccctacc 540
tgaagtcttc caactgccaa aagcacagcc ctttttctct gagctgggtg ttctggctaa 600
cactgacagg ggtcgcttgt tcctgtgcag tgggcatcaa gtacatgggt gtgttcacgt 660
acgtgctcgt gctgggtgtt gcagctgtcc atgcctggca cctgcttgga gaccagactt 720
tgtccaatgt aggtgctgat gtccagtgtc gcatgaggcc ggcctgtatg gggcagatgc 780

```

```

ggatgtcaca ggggggtctgt gtgtttctgtc acttgctcgc ccgagcagtg gcttttgcctgg 840
tcatccccgt cgtcctgtac ttactgttct tctacgtcca cttgattcta gtcttccgct 900
ctgggccccca cgaccaaatc atgtccagtg ccttccaggc cagcttagag ggaggactag 960
ctcggatcac ccagggtcag ccactggagg tggccttttg gtcccaggtc actctgagga 1020
acgtcttttg gaaacctgtg ccctgctggc ttcattecca ccaggacacc taccctatga 1080
tatatgagaa cggccgaggc agctcccacc agcaacaggt gacctgttac cccttcaaag 1140
acgtcaataa ctgggtggatt gtaaaggatc ccaggaggca ccagctggtg gtgagcagcc 1200
ctccgagacc tgtgaggcac ggggacatgg tgcagctggt ccacggcatg accacccgct 1260
ccctgaacac gcatgatgtt gcagcccccc tgagccccca ttcacaggag gtctcctgct 1320
acattgacta taacatctcc atgcccgccc agaacctctg gagactggaa atttgtgaaca 1380
gaggatctga cacagacgtc tggaaagacca tctctcaga ggtccgcttt gtgcacgtga 1440
acacttccgc tgtcttaaag ctgagcgggg ctcacctccc tgactggggg tatcggcaac 1500
tggagatcgt cgggggagaag ctgtcccggg gctaccacgg gagcacggtg tggaaacgtg 1560
aggagcacgg atacggcgcg agccaggagc agagggagcg ggaacgggag ctgcactcac 1620
ctgctcaggt ggacgtcagc aggaacctca gcttcatggc gagattctcg gagctgcagt 1680
ggaggatgct ggcgctgaga agtgatgact cggaacacaa gtacagctcc agccactgg 1740
agtgggtcac cctggacacc aatattgcct actggctgca cccagggacc agcgtctaga 1800
tccacctact tggaaacata gtgatctggg tttcgggcag cctcgctctg gccatctacg 1860
ccctgctgtc cttgtggtac ctgctccgac ggcaagaaa tgtccatgac ctcccctagg 1920
atgctggtct gcgtgggtg ctggctgggg cgctgtgtgc cgggtggctgg gcagtgaact 1980
acctcccgct cttcctgatg gagaagacac tcttctctta ccactacctg ccgcactca 2040
ccttccaaat ccttctgtc cctgtggtcc tgcagcacat cagcgaccac ctgtgcaggt 2100
cccagctcca gaggagcatc ttcagcgcgc tgggtgggtggc ctggtactcc tccgctgccc 2160
acgtgtccaa cacgctgcgc ccactcacct acggggacaa gtcactctcg ccacatgaac 2220
tcaaggccct tcgctgaaa gacagctggg acatcttgat ccgaaaacac tagaacaaga 2280
gtgtggcaaa gaacaccgt gctggggctcg ggacgaggtt gaagggtctt ggtcaatgta 2340
cgtaatgagc aggggtgggccc ccacgctggg aggacacggg ctgggctgag cagggcctct 2400
agtggaacac atgggggtct cattgaaaag ctctctgatg agcacctcct tttgtgcaaa 2460
gttaatTTTT tctcgacaat aaagatattc cgtgtcttca cccctgaact aagacacagg 2520
gagtatttca gaggccagcg taggagtcac cgacaacgaa aagccgagaa cccagggccca 2580
gcagtggagc ctcagcagac cagggcctgg tcttgctaa ttgctgcagg gtggagtttg 2640
atctggcaga cccgatctc cttcatgaac acccagcaac ctgagcaagt cccggccctg 2700
ccctcagcga gcccggcagg cgtcccggga cagctcagtg ttggagggcc acctgaacca 2760
cgagccaggg ctggggccttg catgtcattg tctatgacag cgtcaagact ggcccttggc 2820
accgtgctgt gtggaaaccc tcccctctga gactccactg agacgtggct gagtgaatc 2880
ttcctcgtca gtggtcaagg tgtgtcatcc atacagctcc atgcctttgt cttttttaaa 2940
tgtaattaaa aaaggaacca actggaaaaa aaaaaaa 2977

```

<210> 55
<211> 729
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7488579CB1

```

<400> 55
atgctcctcc ttgctcccca gatgctgaat ctgctgctgc tggcgtgccc cgtcctggcg 60
agccgcgcct acgcggcccc tgccccaggc caggccctgc agcagtgagg catcgttggg 120
ggtcaggagg cccccaggag caagtggccc tggcaggtga gcctgagagt ccacggcccc 180
tactggatgc acttctgcgg gggctccctc atccaccccc agtgggtgct gaccgcagcg 240
cactgcgtgg gaccggacgt caaggatctg gccgccctca gggtgcaact gcgggagcag 300
cacctctact accaggacca gctgctgccc gtcagcagga tcatcgtgca cccacagttc 360
tacatcatcc agaccggggc ggacatcgcc ctgctggagc tggaggagcc cgtgaacatc 420
tccagccaca tccacacggc cacgctgccc cctgcctcgg agaccttccc cccggggatg 480
ccgtgctggg tcaactggctg gggcgacgtg gacaataatg tgcacctgcc gccgccatac 540
ccgctgaagg aggtggaagt ccccgtagtg gaaaaccacc tttgcaacgc ggaatatcac 600
accggcctcc atacgggcca cagctttcaa atcgtccgcg atgacatgct gtgtgccccg 660
agcgaaaatc acgactcctg ccagggtgac tctggagggc cctggtctg caagggtgaat 720
ggcacctaa 729

```

<210> 56

<211> 1879
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7510521CB1

<400> 56
 gggtttccggg cgggcgtact atttcaaggc gcgcgcctcg tgggtggactc accgctagcc 60
 cgcagcgctc ggcttcctgg taattcttca cctcttttct cagctccctg cagcatgggt 120
 gctggggccct ccttgctgct cgcgcgcctc ctgctgcttc tctccggcga cggcgccgtg 180
 cgctgcgaca cacctgccaa ctgcacctat cttgacctgc tgggcacctg ggtcttccag 240
 gaccacaaga aaaaaaagta gtggtgtacc ttcagaagct ggatacagca tatgatgacc 300
 ttggcaattc tggccatttc accatcattt acaaccaagg ctttgagatt gtgttgaatg 360
 actacaagtg gtttgctttt ttttaagtata aagaagaggg cagcaagggtg accacttact 420
 gcaacgagac aatgactggg tgggtgcatg atgtgttggg ccggaactgg gcttgtttca 480
 ccggaaagaa ggtgggaact gcctctgaga atgtgtatgt caacacagca caccttaaga 540
 attctcagga aaagtattct aataggctct acaagtatga tcacaacttt gtgaaagcta 600
 tcaatgccat tcagaagtct tggactgcaa ctacatacat ggaatatgag actcttacc 660
 tgggagatat gattaggaga agtgggtggc acagtcgaaa aatccaagg cccaaacctg 720
 caccactgac tgctgaaata cagcaaaaaga ttttgcatth gccaatctt tgggactgga 780
 gaaatgttca tggatcaaat tttgtcagtc ctgttcgaaa ccaagcatoc tgtggcagct 840
 gctactcatt tgcttctatg ggtatgctag aagcgagaat ccgtatacta accaacaatt 900
 ctcagacccc aatcctaagc cctcaggagg ttgtgtcttg tagccagtat gctcaaggct 960
 gtgaaggcgg cttcccatat cttattgcag gaaagtacgc ccaagattht gggctggtgg 1020
 aagaagcttg cttccctac acaggcactg attctccatg caaatgaag gaagactgct 1080
 ttcgttatta ctctctgag taccactatg taggaggtht ctatggaggc tgcaatgaag 1140
 ccctgatgaa gcttgagttg gtccatcatg ggcccatggc agttgcttht gaagtatatg 1200
 atgacttctt ccactacaaa aaggggatct accaccacac tgggtctaaga gaccctthtca 1260
 acccctthtga gctgactaat catgctgttc tgcttgtggg ctatggcact gactcagcct 1320
 ctgggatgga ttactggatt gttaaaaaca gctggggcac cggctggggg gagaatggct 1380
 acttccggat ccgcagagga actgatgagt gtgcaattga gagcatagca gtggcagcca 1440
 caccaattcc taaattgtag ggtatgcctt ccagtatttc ataatgatct gcatcagttg 1500
 taaaggggaa ttggtatatt cacagactgt agactthtca cagcaatctc agaagcttac 1560
 aaatagatth ccatgaagat atthgtcttc agaattaaaa ctgcccttaa tthtaataata 1620
 cctthtcaatc ggccactggc catthththt taagtattca attaatggg aaththtctg 1680
 aagatggtca gctatgaagt aatagagtht gcttaatcat ttgtaattca aacatgctat 1740
 atthththta atcaatgtga aaacatagac ttatththta attgtaccaa tcacaagaaa 1800
 ataatggcaa taattatcaa aactththta atagatgctc atatththta aataaagtht 1860
 taaaaataaa aaaaaaaaaa 1879

